

WEEK 9 PART 3: BOUNDARY VALUE PROBLEMS

We are used to initial value problems where we are given initial data. What if we are given boundary data instead? There are many applications where things are happening for a long period of time and we don't know what happened in the beginning, but we do know something about the boundary. The usual problems are solved in a similar fashion to Initial Value Problems. We do however have a bit more theory.

Definition 1. The boundary values (for a second order ODE) $y(a)$, $y(b)$, $y'(a)$, and/or $y'(b)$ are said to be homogeneous if any two of the above boundary data are zero.

We also have eigenvalue problems for BVPs. Recall that for matrices the eigenvalue problems were of the form $Ax = \lambda x$, where we solve for the "eigenvalue", λ . For BVPs of a second order ODE, we consider our linear operator to be $L = d^2/dx^2$ (for matrices the linear operator is the matrix A). So we wish to solve the problem $Ly = \lambda y$; i.e. $y'' + \lambda y = 0$. Here the y_n 's corresponding to λ_n 's are called eigenfunctions (similar to eigenvectors in the matrix case). We notice that eigenvalue problems are only for homogeneous boundary data.

Definition 2. The boundary value problem

$$y'' + \lambda y = 0; \quad (\text{with homogeneous boundary conditions}), \quad (1)$$

is called an eigenvalue problem. And the nontrivial (i.e. $y_n \neq 0$) solutions y_n corresponding to λ_n are the eigenfunctions of the corresponding eigenvalues.

Now lets do some boundary value problems,

Ex: $y'' + y = 0$; $y'(0) = 1$, $y(L) = 0$.

Solution: The characteristic polynomial gives us

$$r^2 + 1 = 0 \Rightarrow r = \pm i \Rightarrow y = A \cos t + B \sin t \Rightarrow y' = -A \sin t + B \cos t.$$

Then our first boundary condition gives $y'(0) = B = 1$, and

$$y(L) = A \cos L + \sin L = 0 \Rightarrow A = -\tan L; \quad L \neq (2k+1)\frac{\pi}{2}, \quad k = 0, \pm 1, \pm 2, \dots$$

However, if $\cos L = 0$, $\sin L = 0$, but this is clearly false because $\sin x \neq 0$ when $\cos x = 0$ and vice-versa, so the BVP has no solution if $L = (2k+1)\frac{\pi}{2}$.

Ex: $y'' + \lambda y = 0$; $y'(0) = y'(\pi) = 0$.

Solution:

(i) If $\lambda > 0$, let $\lambda = \mu^2$. Then

$$r = \pm i\mu \Rightarrow y = A \cos \mu t + B \sin \mu t \Rightarrow y' = -A\mu \sin \mu t + B\mu \cos \mu t$$

From the first boundary condition we get $y'(0) = B\mu = 0 \Rightarrow B = 0$ because $\lambda > 0$. From the second B.C. we get $y'(\pi) = -A\mu \sin \mu\pi = 0$. Since we don't want trivial solutions if we can avoid them we can't have $A = 0$, so we require $\sin \mu\pi = 0$ then $\mu = n\pi$ where $n = 1, 2, \dots$, so our eigenfunctions for the corresponding eigenvalues are

$$y_n = \cos n\pi t; \quad \lambda_n = n^2, \quad n = 1, 2, \dots$$

(ii) If $\lambda < 0$, let $\lambda = -\mu^2$. Then

$$r = \pm \mu \Rightarrow y = c_1 e^{\mu t} + c_2 e^{-\mu t} = A \cosh \mu t + B \sinh \mu t \Rightarrow y' = A \sinh \mu t + B \cosh \mu t.$$

The B.C.'s give $y'(0) = B\mu = 0 \Rightarrow B = 0$ and $y'(\pi) = A \sinh \mu\pi = 0$, but \sinh is only zero at zero and $\mu \neq 0$ since $\lambda < 0$, so we have $A = 0$. Then $y \equiv 0$, so unfortunately we get a trivial solution. (iii) If $\lambda = 0$, $y = c_1 x + c_0 \Rightarrow y' = c_1$, then applying the B.C.'s give $y'(0) = c_1 = 0$ and $y'(\pi) = 0$ automatically. Then our eigenvalue and eigenfunction are

$$y_0 = 1, \quad \lambda_0 = 0.$$

Notice I left out the constants. It is up to you if you want to include it or not.

Complete Solution:

$$y = c_0 + \sum_{n=1}^{\infty} A_n \cos n\pi t$$

Ex: $y'' + \lambda y = 0$; $y'(0) = y(L) = 0$

Solution:

(i) If $\lambda > 0$, let $\lambda = \mu^2$, then

$$y = A \cos \mu t + B \sin \mu t \Rightarrow y' = -A\mu \sin \mu t + B\mu \cos \mu t.$$

Notice how we have the same exact general solution! You do enough of these problems and you can go straight to the solution and it's derivative without having to do the characteristic polynomial. Now, from the B.C.'s we get $y'(0) = B\mu = 0 \Rightarrow B = 0$ and $y(L) = A \cos \mu L = 0$. So we require $\mu = (2n - 1)\pi/2L$ where $n = 1, 2, 3, \dots$, then our eigenvalues and eigenfunctions are

$$y_n = A_n \cos \left((2n - 1) \frac{\pi}{2} t \right); \lambda_n = (2n - 1)^2 \frac{\pi^2}{4}, n = 1, 2, 3, \dots$$

(ii) If $\lambda < 0$, let $\lambda = -\mu^2$, then

$$y = A \cosh \mu t + B \sinh \mu t \Rightarrow y' = A\mu \sinh \mu t + B\mu \cosh \mu t.$$

From the B.C.'s we get $y'(0) = B\mu = 0 \Rightarrow B = 0$ and $y(L) = A \cosh \mu L = 0 \Rightarrow A = 0$, again it's the trivial solution $y \equiv 0$.

(iii) If $\lambda = 0$, $y = c_1 x + c_0 \Rightarrow y' = c_1$, from the B.C.'s we get $y'(0) = c_1 = 0$ and $y(L) = c_0 = 0$, so again we have the trivial solution $y \equiv 0$.

Complete Solution:

$$y = \sum_{n=1}^{\infty} A_n \cos \left((2n - 1) \frac{\pi}{2} t \right)$$