

The static and dynamic extension of DFT to superfluid fermionic systems:

SLDA, ASLDA and TDSLDA

Aurel Bulgac
University of Washington

Collaborators:

Yongle YU (now at Wuhan Institute of Physics and Mathematics)

Michael M. FORBES (now at LANL)

Sukjin YOON (UW)

Kenneth J. ROCHE (ORNL)

Yuan Lung LUO (UW)

Piotr MAGIERSKI (Warsaw and UW)

Ionel STETCU (UW)

Funding: DOE grants No. DE-FG02-97ER41014 (UW NT Group)
DE-FC02-07ER41457 (SciDAC-UNEDF)

Outline:

- **Extension of DFT to superfluid systems - SLDA**
- **Extension of SLDA to asymmetric systems - ASLDA**
- **Extension of SLDA to time dependent phenomena – TDSLDA**
- **SLDA for nuclei (and vortices in neutron stars)**
- **SLDA for unitary gas (and vortices in a unitary gas), derivation of EDF from *ab initio* calculations and its validation**
- **Derivation of ASLDA and its verification**
- **New states of matter, new mechanisms of pairing**
- **Dynamics of superfluid fermionic systems and new type of collective modes**
- **Why TDSLDA is better than QRPA?**

Very brief/skewed summary of DFT

Kohn-Sham theorem

$$H = \sum_i^N T(i) + \sum_{i<j}^N U(ij) + \sum_{i<j<k}^N U(ijk) + \dots + \sum_i^N V_{ext}(i)$$

$$H\Psi_0(1,2,\dots,N) = E_0\Psi_0(1,2,\dots,N)$$

$$n(\vec{r}) = \langle \Psi_0 | \sum_i^N \delta(\vec{r} - \vec{r}_i) | \Psi_0 \rangle$$

Injective map
(one-to-one)

$$\Psi_0(1,2,\dots,N) \Leftrightarrow V_{ext}(\vec{r}) \Leftrightarrow n(\vec{r})$$

$$E_0 = \min_{n(\vec{r})} \int d^3r \left\{ \frac{\hbar^2}{2m} \tau(\vec{r}) + \varepsilon[n(\vec{r})] + V_{ext}(\vec{r})n(\vec{r}) \right\}$$

$$n(\vec{r}) = \sum_i^N |\varphi_i(\vec{r})|^2, \quad \tau(\vec{r}) = \sum_i^N |\vec{\nabla} \varphi_i(\vec{r})|^2$$

Universal functional of particle density alone
Independent of external potential

How to construct and validate an *ab initio* EDF?

- ❑ Given a many body Hamiltonian determine the properties of the infinite homogeneous system as a function of density
- ❑ Extract the energy density functional (EDF)
- ❑ Add gradient corrections, if needed or known how (?)
- ❑ Determine in an *ab initio* calculation the properties of a select number of wisely selected finite systems
- ❑ Apply the energy density functional to inhomogeneous systems and compare with the *ab initio* calculation, and if lucky declare Victory!

Extended Kohn-Sham equations

Position dependent mass

$$E_{gs} = \int d^3r \left\{ \frac{\hbar^2}{2m^*[n(\vec{r})]} \tau(\vec{r}) + \varepsilon[n(\vec{r})]n(\vec{r}) \right\}$$
$$n(\vec{r}) = \sum_{i=1}^N |\psi_i(\vec{r})|^2 \quad \tau(\vec{r}) = \sum_{i=1}^N |\vec{\nabla} \psi_i(\vec{r})|^2$$
$$-\vec{\nabla} \frac{\hbar^2}{2m^*[n(\vec{r})]} \vec{\nabla} \psi_i(\vec{r}) + U(\vec{r})\psi_i(\vec{r}) = \varepsilon_i \psi_i(\vec{r})$$

Normal Fermi systems only!

However, not everyone is normal!

Superconductivity and superfluidity in Fermi systems

- ✓ Dilute atomic Fermi gases $T_c \approx 10^{-12} - 10^{-9} \text{ eV}$
- ✓ Liquid ^3He $T_c \approx 10^{-7} \text{ eV}$
- ✓ Metals, composite materials $T_c \approx 10^{-3} - 10^{-2} \text{ eV}$
- ✓ Nuclei, neutron stars $T_c \approx 10^5 - 10^6 \text{ eV}$
- QCD color superconductivity $T_c \approx 10^7 - 10^8 \text{ eV}$

units (1 eV \approx 10⁴ K)

SLDA - Extension of Kohn-Sham approach to superfluid Fermi systems

$$E_{gs} = \int d^3r \varepsilon(n(\vec{r}), \tau(\vec{r}), \nu(\vec{r}))$$

$$n(\vec{r}) = 2 \sum_k |v_k(\vec{r})|^2, \quad \tau(\vec{r}) = 2 \sum_k |\vec{\nabla} v_k(\vec{r})|^2$$

$$\nu(\vec{r}) = \sum_k u_k(\vec{r}) v_k^*(\vec{r})$$

$$\begin{pmatrix} T + U(\vec{r}) - \mu & \Delta(\vec{r}) \\ \Delta^*(\vec{r}) & -(T + U(\vec{r}) - \mu) \end{pmatrix} \begin{pmatrix} u_k(\vec{r}) \\ v_k(\vec{r}) \end{pmatrix} = E_k \begin{pmatrix} u_k(\vec{r}) \\ v_k(\vec{r}) \end{pmatrix}$$

Mean-field and pairing field are both local fields!
(for sake of simplicity spin degrees of freedom are not shown)

There is a little problem! The densities ν and τ diverges!

The SLDA (renormalized) equations

$$E_{gs} = \int d^3r \left\{ \varepsilon_N \left[n(\vec{r}), \tau(\vec{r}) \right] + \varepsilon_S \left[n(\vec{r}), \nu(\vec{r}) \right] \right\}$$

$$\varepsilon_S \left[n(\vec{r}), \nu(\vec{r}) \right] \stackrel{\text{def}}{=} -\Delta(\vec{r})\nu_c(\vec{r}) = g_{\text{eff}}(\vec{r}) |\nu_c(\vec{r})|^2$$

$$\begin{cases} [h(\vec{r}) - \mu]u_i(\vec{r}) + \Delta(\vec{r})v_i(\vec{r}) = E_i u_i(\vec{r}) \\ \Delta^*(\vec{r})u_i(\vec{r}) - [h(\vec{r}) - \mu]v_i(\vec{r}) = E_i v_i(\vec{r}) \end{cases} \quad \begin{cases} h(\vec{r}) = -\vec{\nabla} \frac{\hbar^2}{2m(\vec{r})} \vec{\nabla} + U(\vec{r}) \\ \Delta(\vec{r}) = -g_{\text{eff}}(\vec{r})\nu_c(\vec{r}) \end{cases}$$

$$\frac{1}{g_{\text{eff}}(\vec{r})} = \frac{1}{g[n(\vec{r})]} - \frac{m(\vec{r})k_c(\vec{r})}{2\pi^2\hbar^2} \left\{ 1 - \frac{k_F(\vec{r})}{2k_c(\vec{r})} \ln \frac{k_c(\vec{r}) + k_F(\vec{r})}{k_c(\vec{r}) - k_F(\vec{r})} \right\}$$

$$n_c(\vec{r}) = 2 \sum_{E_i \geq 0}^{E_c} |v_i(\vec{r})|^2, \quad \nu_c(\vec{r}) = \sum_{E_i \geq 0}^{E_c} v_i^*(\vec{r})u_i(\vec{r})$$

$$E_c + \mu = \frac{\hbar^2 k_c^2(\vec{r})}{2m(\vec{r})} + U(\vec{r}), \quad \mu = \frac{\hbar^2 k_F^2(\vec{r})}{2m(\vec{r})} + U(\vec{r})$$

Position and momentum dependent running coupling constant
Observables are independent of cut-off energy (when chosen properly).

**Superfluid Local Density Approximation (SLDA)
for a unitary Fermi gas**

The renormalized SLDA energy density functional

Only this combination is cutoff independent

$$\varepsilon(\vec{r}) = \left[\alpha \frac{\tau_c(\vec{r})}{2} - \Delta(\vec{r})v_c(\vec{r}) \right] + \beta \frac{3(3\pi^2)^{2/3} n^{5/3}(\vec{r})}{5}$$

$$n(\vec{r}) = 2 \sum_k \left| v_k(\vec{r}) \right|^2, \quad \tau_c(\vec{r}) = 2 \sum_{E < E_c} \left| \vec{\nabla} v_k(\vec{r}) \right|^2, \quad v_c(\vec{r}) = \sum_{E < E_c} u_k(\vec{r}) v_k^*(\vec{r})$$

$$\frac{1}{g_{eff}(\vec{r})} = \frac{n^{1/3}(\vec{r})}{\gamma} - \frac{k_c(\vec{r})}{2\pi^2 \alpha} \left[1 - \frac{k_0(\vec{r})}{2k_c(\vec{r})} \ln \frac{k_c(\vec{r}) + k_0(\vec{r})}{k_c(\vec{r}) - k_0(\vec{r})} \right]$$

$$E_c + \mu = \alpha \frac{k_c^2(\vec{r})}{2} + U(\vec{r}), \quad \mu = \alpha \frac{k_0^2(\vec{r})}{2} + U(\vec{r})$$

$$U(\vec{r}) = \beta \frac{(3\pi^2)^{2/3} n^{2/3}(\vec{r})}{2} - \frac{|\Delta(\vec{r})|^2}{3\gamma n^{2/3}(\vec{r})} + V_{ext}(\vec{r})$$

$$\Delta(\vec{r}) = -g_{eff}(\vec{r})v_c(\vec{r})$$

No free parameters, EDF is fully determined by *ab initio* calculations


Bulgac, Phys. Rev. A 76, 040502(R) (2007)

Parameters defining SLDA functional for a unitary gas:

$$\left\{ \begin{array}{l} \xi_s = \frac{5E}{3N \varepsilon_F} = 0.42(2) \\ \eta = \frac{\Delta}{\varepsilon_F} = 0.504(24) \\ \zeta = \frac{\mu}{\varepsilon_F} = 0.42(2) \end{array} \right. \Rightarrow \left\{ \begin{array}{l} \alpha = 1.14 \\ \beta = -0.553 \\ \frac{1}{\gamma} = -0.0906 \end{array} \right.$$

Bonus!

Quasiparticle spectrum in homogeneous matter



solid/dotted blue line

- SLDA, homogeneous GFMC due to Carlson et al

red circles

- GFMC due to Carlson and Reddy

dashed blue line

- SLDA, homogeneous MC due to Juillet

black dashed-dotted line

- meanfield at unitarity

Extra Bonus!


The normal state has been also determined in GFMC

$$\xi_N = \frac{5E}{3N\varepsilon_F} = 0.55(2)$$

SLDA functional predicts

$$\xi_N = \alpha + \beta = 0.59$$

Fermions at unitarity in a harmonic trap



GFMC - Chang and Bertsch, Phys. Rev. A 76, 021603(R) (2007)

FN-DMC - von Stecher, Greene and Blume, PRL 99, 233201 (2007)

PRA 76, 053613 (2007)

TABLE I: Table I. The energies $E(N)$ calculated within the GFMC [14], FN-DMC [15] and SLDA. When two numbers are present the first was calculated as the expectation value of the Hamiltonian/functional, while the second is the value obtained using the virial theorem, namely $E(N) = m\omega^2 \int d^3r n(\mathbf{r})r^2$ [23].

N	E_{GFMC}	E_{FN-DMC}	E_{SLDA}
1	1.5		1.37
2	2.01/1.95	2.002	2.33/2.34
3	4.28/4.19		4.62/4.62
4	5.10	5.069	5.52/5.56
5	7.60		7.98/8.02
6	8.70	8.67	9.07/9.14
7	11.3		11.83/11.91
8	12.6/11.9	12.57	12.94/13.06
9	15.6		16.06/16.20
10	17.2	16.79	17.15/17.33
11	19.9		20.36/20.56
12	21.5	21.26	21.63/21.88
13	25.2		24.96/25.23
14	26.6/26.0	25.90	26.32/26.65
15	30.0		29.78/30.14
16	31.9	30.92	31.21/31.62
17	35.4		34.81/35.26
18	37.4	36.00	36.27/36.78
19	41.1		40.02/40.58
20	43.2/40.8	41.35	41.51/42.12
21	46.9		45.42/46.10
22	49.3		46.92/47.64

**NB Particle projection
neither required nor
needed in SLDA!!!**


SLDA - Extension of Kohn-Sham approach to superfluid Fermi systems

$$E_{gs} = \int d^3r \left\{ \varepsilon(n(\vec{r}), \tau(\vec{r}), \nu(\vec{r})) + \leftarrow \text{universal functional} \right. \\ \left. V_{ext}(\vec{r})n(\vec{r}) + \Delta_{ext}(\vec{r})\nu(\vec{r}) + \Delta_{ext}^*(\vec{r})\nu^*(\vec{r}) \right\} \\ \text{(independent of external potential)}$$

$$n(\vec{r}) = 2 \sum_k |\mathbf{v}_k(\vec{r})|^2, \quad \tau(\vec{r}) = 2 \sum_k |\vec{\nabla} \mathbf{v}_k(\vec{r})|^2$$

$$\nu(\vec{r}) = \sum_k \mathbf{u}_k(\vec{r}) \mathbf{v}_k^*(\vec{r})$$

$$\begin{pmatrix} T + U(\vec{r}) - \mu & \Delta(\vec{r}) \\ \Delta^*(\vec{r}) & -(T + U(\vec{r}) - \mu) \end{pmatrix} \begin{pmatrix} \mathbf{u}_k(\vec{r}) \\ \mathbf{v}_k(\vec{r}) \end{pmatrix} = E_k \begin{pmatrix} \mathbf{u}_k(\vec{r}) \\ \mathbf{v}_k(\vec{r}) \end{pmatrix}$$



GFMC - Chang and Bertsch, Phys. Rev. A 76, 021603(R) (2007)

FN-DMC - von Stecher, Greene and Blume, PRL 99, 233201 (2007)

PRA 76, 053613 (2007)

- **Agreement between GFMC/FN-DMC and SLDA extremely good, a few percent (at most) accuracy**

Why not better?

A better agreement would have really signaled big troubles!

- **Energy density functional is not unique, in spite of the strong restrictions imposed by unitarity**
- **Self-interaction correction neglected
smallest systems affected the most**
- **Absence of polarization effects
spherical symmetry imposed, odd systems mostly affected**
- **Spin number densities not included
extension from SLDA to SLSD(A) needed
ab initio results for asymmetric system needed**
- **Gradient corrections not included, ... but very likely small!!!**

Challenges towards implementation of SLDA in nuclei

Towards a universal nuclear density functional

S. A. Fayans

Kurchatov Institute Russian Science Center, 123182 Moscow, Russia

The total energy density of a nuclear system is represented as

$$\varepsilon = \varepsilon_{\text{kin}} + \varepsilon_v + \varepsilon_s + \varepsilon_{\text{Coul}} + \varepsilon_{s'} + \varepsilon_{\text{anom}}, \tag{1}$$

where ε_{kin} is the kinetic energy term which, since we are constructing a Kohn–Sham type functional, is taken with the free operator $t=p^2/2m$, i.e., with the effective mass $m^*=m$; all the other terms are discussed below.

The volume term in (1) is chosen to be in the form



$$\varepsilon_v = \frac{2}{3} \epsilon_F^0 \rho_0 \left[a_+^v \frac{1 - h_+^v x_+^\sigma}{1 + h_+^v x_+^\sigma} x_+^2 + a_-^v \frac{1 - h_-^v x_-^\sigma}{1 + h_-^v x_-^\sigma} x_-^2 \right].$$

Here and in the following $x_\pm = (\rho_n \pm \rho_p)/2\rho_0$, $\rho_{n(p)}$ is the neutron (proton) density, $2\rho_0$ is the equilibrium density of symmetric nuclear matter with



The surface part in Eq. (1) is meant to describe the finite-range and nonlocal in-medium effects which may presumably be incorporated phenomenologically within the EDF framework in a localized form by introducing a dependence on density gradients. It is taken as follows:

$$\varepsilon_s = \frac{2}{3} \epsilon_F^0 \rho_0 \frac{a_+^s r_0^2 (\nabla x_+)^2}{1 + h_+^s x_+^\sigma + h_+^s r_0^2 (\nabla x_+)^2}, \tag{3}$$


with $h_\pm^s = h_\pm^v$, a_\pm^s and h_\pm^s the two free parameters. Such a form is obtained by adding




Baldo, Schuck, and Vinas, arXiv:0706.0658



Gandolfi et al. arXiv:0805.2513



Gezerlis and Carlson, PRC 77, 032801 (2008)



Bulgac et al. arXiv:0801.1504, arXiv:0803.3238

Let us summarize some of the ingredients of the SLDA in nuclei

Energy Density (ED) describing the normal system

ED contribution due to superfluid correlations

$$E_{gs} = \int d^3r \left\{ \varepsilon_N[\rho_n(\vec{r}), \rho_p(\vec{r})] + \varepsilon_S[\rho_n(\vec{r}), \rho_p(\vec{r}), \nu_n(\vec{r}), \nu_p(\vec{r})] \right\}$$
$$\left\{ \begin{array}{l} \varepsilon_N[\rho_n(\vec{r}), \rho_p(\vec{r})] = \varepsilon_N[\rho_p(\vec{r}), \rho_n(\vec{r})] \\ \varepsilon_S[\rho_n(\vec{r}), \rho_p(\vec{r}), \nu_n(\vec{r}), \nu_p(\vec{r})] = \varepsilon_S[\rho_p(\vec{r}), \rho_n(\vec{r}), \nu_p(\vec{r}), \nu_n(\vec{r})] \end{array} \right.$$

Isospin symmetry

(Coulomb energy and other relatively small terms not shown here.)

Let us consider the simplest possible ED compatible with nuclear symmetries and with the fact that nuclear pairing correlations are relatively weak.

$$\varepsilon_S[\rho_p, \rho_n, \nu_p, \nu_n] = g_0 \underbrace{|\nu_p + \nu_n|^2}_{\text{like } \rho_p + \rho_n} + g_1 \underbrace{|\nu_p - \nu_n|^2}_{\text{like } \rho_p - \rho_n}$$

g_0 and g_1 could depend as well on ρ_p and ρ_n

Let us stare at the anomalous part of the ED for a moment, ... or two.

SU(2) invariant

$$\begin{aligned}\mathcal{E}_S[v_p, v_n] &= g_0 |v_p + v_n|^2 + g_1 |v_p - v_n|^2 \\ &= g [|v_p|^2 + |v_n|^2] + g' [v_p^* v_n + v_n^* v_p] \\ g &= g_0 + g_1 \quad g' = g_0 - g_1\end{aligned}$$

?

NB I am dealing here with s-wave pairing only (S=0 and T=1)!

The last term could not arise from a two-body bare interaction.

In the end one finds that a suitable superfluid nuclear EDF has the following structure:





Isospin symmetric

$$\mathcal{E}_S[\nu_p, \nu_n] = g(\rho_p, \rho_n)[|\nu_p|^2 + |\nu_n|^2] + f(\rho_p, \rho_n)[|\nu_p|^2 - |\nu_n|^2] \frac{\rho_p - \rho_n}{\rho_p + \rho_n}$$

where $g(\rho_p, \rho_n) = g(\rho_n, \rho_p)$

and $f(\rho_p, \rho_n) = f(\rho_n, \rho_p)$

The same coupling constant for both even and odd neutron/proton numbers!!!



A single universal parameter for pairing in nuclei!

Yu and Bulgac , Phys. Rev. Lett. 90, 222501 (2003)

Bulgac and Yu, Phys. Rev. Lett. 88, 042504 (2002)

Until now we kept the numbers of spin-up and spin-down equal.



What happens when there are not enough partners for everyone to pair with?

(In particular this is what one expects to happen in color superconductivity, due to the heavier strange quark)


What theory tells us?

Green – Fermi sphere of spin-up fermions
Yellow – Fermi sphere of spin-down fermions

If $|\mu_{\uparrow} - \mu_{\downarrow}| < \frac{\Delta}{\sqrt{2}}$ the same solution as for $\mu_{\uparrow} = \mu_{\downarrow}$



LOFF/FFLO solution (1964)
Pairing gap becomes a spatially varying function
Translational invariance broken




Muether and Sedrakian (2002)
Translational invariant solution
Rotational invariance broken

What we think is happening in spin imbalanced systems?

Induced P-wave superfluidity

Two new superfluid phases where before they were not expected




One Bose superfluid coexisting with one P-wave Fermi superfluid

Two coexisting P-wave Fermi superfluids

Bulgac, Forbes, and Schwenk, Phys. Rev. Lett. 97, 020402 (2006)

What happens at unitarity?




Predicted quantum first phase order transition, subsequently observed in MIT experiment, Shin *et al.* *Nature*, **451**, 689 (2008)

Red points with error bars – subsequent DMC calculations for normal state due to Lobo *et al.*, *PRL* **97**, 200403 (2006)

$$E(n_a, n_b) = \frac{3 (6\pi^2)^{2/3} \hbar^2}{5 \cdot 2m} \left[n_a g \left(\frac{n_b}{n_a} \right) \right]^{5/3}, \quad n_a \geq n_b$$

Bulgac and Forbes, *PRA* **75**, 031605(R) (2007)

A refined EOS for spin unbalanced systems



Red line: Larkin-Ovchinnikov phase

Black line: normal part of the energy density

Blue points: DMC calculations for normal state, Lobo et al, PRL 97, 200403 (2006)

Gray crosses: experimental EOS due to Shin, Phys. Rev. A 77, 041603(R) (2008)



$$E(n_a, n_b) = \frac{3 (6\pi^2)^{2/3} \hbar^2}{5 \cdot 2m} \left[n_a g \left(\frac{n_b}{n_a} \right) \right]^{5/3}$$

**Bulgac and Forbes,
Phys. Rev. Lett. 101, 215301 (2008)**

How this new refined EOS for spin imbalanced systems was obtained?

Through the use of the (A)SLDA , which is an extension of the Kohn-Sham LDA to superfluid systems

A Unitary Fermi Supersolid: the Larkin-Ovchinnikov phase



Bulgac and Forbes
Phys. Rev. Lett. 101, 215301 (2008)

$$P[\mu_a, \mu_b] = \frac{2}{30\pi^2} \left(\frac{2m}{\hbar^2} \right)^{3/2} \left[\mu_a h \left(\frac{\mu_b}{\mu_a} \right) \right]^{5/2}$$

Time Dependent Phenomena and Formalism

The time-dependent density functional theory is viewed in general as a reformulation of the exact quantum mechanical time evolution of a many-body system when only single-particle properties are considered.

A.K. Rajagopal and J. Callaway, *Phys. Rev. B* 7, 1912 (1973)

V. Peuckert, *J. Phys. C* 11, 4945 (1978)

E. Runge and E.K.U. Gross, *Phys. Rev. Lett.* 52, 997 (1984)

<http://www.tddft.org>

$$\left\{ \begin{array}{l} [h(\vec{r}, t) + V_{ext}(\vec{r}, t) - \mu]u_i(\vec{r}, t) + [\Delta(\vec{r}, t) + \Delta_{ext}(\vec{r}, t)]v_i(\vec{r}, t) = i\hbar \frac{\partial u_i(\vec{r}, t)}{\partial t} \\ [\Delta^*(\vec{r}, t) + \Delta_{ext}^*(\vec{r}, t)]u_i(\vec{r}, t) - [h(\vec{r}, t) + V_{ext}(\vec{r}, t) - \mu]v_i(\vec{r}, t) = i\hbar \frac{\partial v_i(\vec{r}, t)}{\partial t} \end{array} \right.$$

Full 3D implementation of TD-SLDA is a petaflop problem and is almost complete.

Bulgac and Roche, J. Phys. Conf. Series 125, 012064 (2008)

Lots of contributions due to Yu, Yoon, Luo, Magierski, and Stetcu

New issues arising in formulating and implementing a TD-DFT:


In ground states currents vanish, but they are present in excited states.

The dependence of the EDF on some currents can be established from general principles, e.g. Galilean invariance:

$$\tau(\vec{r}) \Rightarrow \tau(\vec{r}) - \frac{\vec{j}^2(\vec{r})}{n(\vec{r})}$$




Not all currents and densities can be introduced into the formalism in a such a manner however.

Energy of a (unitary) Fermi system as a function of the pairing gap



Response of a unitary Fermi system to changing the scattering length with time

Tool: TD DFT extension to superfluid systems (TD-SLDA)




- All these modes have a very low frequency below the pairing gap and a very large amplitude and excitation energy as well
- None of these modes can be described either within Quantum Hydrodynamics or Landau-Ginzburg like approaches

Bulgac and Yoon, Phys. Rev. Lett. 102, 085302 (2009)

3D unitary Fermi gas confined to a 1D ho potential well (pancake)

New qualitative excitation mode of a superfluid Fermi system
(non-spherical Fermi momentum distribution)



Black solid line – Time dependence of the cloud radius

Black dashed line – Time dependence of the quadrupole moment of momentum distribution

Bulgac and Yoon, Phys. Rev. Lett. 102, 085302 (2009)


Vortex generation (show movie)

Time-Dependent Superfluid Local Density Approximation

This is a general many-body problem with direct applications, which will provide the time dependent response of superfluid fermionic systems to a large variety of external probes for both cases of small and large amplitude collective motion.

- **Nuclear physics: fission, heavy-ion collision, nuclear reactions, response electromagnetic fields, beta-decay, ...**
 - **Neutron star crust, dynamics of vortices, vortex pinning mechanism**
 - **Cold atom physics, optical lattices, ...**
 - **Condensed matter physics**
-
- **Next frontier: Stochastic TDSLDA**

Generic adiabatic large amplitude potential energy SURFACES



- In LACM adiabaticity is not a guaranteed
- Level crossings are a great source of :
 - entropy production (dissipation)
 - dynamical symmetry breaking
 - non-abelian gauge fields

**Known disease of QRPA, the particle number of excited states is ill defined
see Terasaki, Engel and Bertsch, Phys. Rev. C 77, 044311 (2008)**




FIG. 1. Particle-hole character of the lowest 2^+ solutions. The histogram displays the quantity ΔN defined in Eq. (1) for 155 nuclei in the SLy4 data set (one of which we drop—see text). The values $-2, 0, +2$ correspond to excitations of hole-hole, particle-hole, and particle-particle character, respectively.

**However in TDSLDA the particle number is not violated
and all states have a well defined average particle number.**

Plato's cave

