

















Question: What is the most favorable arrangement of these two spheres?



Answer: The energy of the system does not depend on r as long as r > 2a. NB Assuming that a liquid drop model for the fermions is accurate! This is a very strange answer! Isn't it? Something is amiss here.

Fermi sea $\lambda_{\mathbf{F}}$

Let us try to think of this situation now in quantum mechanical terms.

The dark blue region is really full of de Broglie's waves, which is in the absence of homogeneities are simple plane waves.

When inhomogeneities are present, there are a lot of scattered waves.

Also, there are some almost stationary waves, which reflect back and forth from the two tips of the empty spheres.



As in the case of a musical instrument, in the absence of damping, the stable "musical notes" correspond to stationary modes.

Problems: 1) There is a large number of such modes.
2) The tip-to-tip modes cannot be absolutely stable, as the reflected wave disperses in the rest of the space.

Fermionic Casimir effect

A force from nothing onto nothing AW

Aurel Bulgac (Seattle) Piotr Magierski (Warsaw) Andreas Wirzba (Juelich) H.B.G. Casimir (1948): two parallel uncharged metallic plates attract each other in vacuum

$$\begin{array}{rcl}
& & \frac{F^{\parallel}(L)}{A} &=& -\frac{\hbar c}{L^4} \frac{\pi^2}{240} \approx -1.3 \times 10^{-7} \frac{1}{L^4} \mathsf{N} \frac{\mu \mathsf{m}^4}{\mathsf{cm}^2} \\ & & \mathcal{E}^{\parallel}(L) &=& -\frac{\hbar c}{L^3} \frac{\pi^2}{720} A
\end{array}$$

 Origin: zero-point fluctuations of e.m. field modified by the addition of the two plates relative to free case

$$\Rightarrow$$
 change in the energy of the vacuum: $\sum \hbar \omega_k |_{\text{plates}(L)} - \sum \hbar \omega_k |_{\text{free}}$

Casimir effect: Mesoscopic manifestation of quantum fluctuations of the vacuum

- Related to van der Waals forces (however, the latter always attractive !)
- experimental confirmation in the last decade (for the sphere-plate system!)
 - S. Lamoureaux, Phys. Rev. Lett. 78 (1997);
 - U. Mohideen & A. Roy, Phys. Rev. Lett. 81 (1998);
 - 9-Feb-2001 issue of New York Times about Casimir effect in MicroElectroMechanical Systems; etc.

Many of the subsequesnt slides are borrowed from my collaborator's talk A. Wirzba

Casimir effect and vacuum energy in QFT:

- Infinite zero-point energy can be subtracted (discarded) by suitable redefinition of energy-origin (e.g. normal ordering)
- However, energy-origin can be re-defined only once (and only in homogeneous space (no b.c.'s) and without gravity)
 - ∴ Casmir effect = difference of (properly regularized) eigen-mode sums of *constrained* QFT – eigen-mode sums of *free* QFT

$$\begin{split} \varepsilon_{C} &= \lim_{V \to \infty} \frac{\mathcal{E}_{C}}{V} \quad (V : \text{quantization box}) \\ &= \lim_{\Lambda \to \infty} \lim_{V \to \infty} \underbrace{(-1)^{2S}}_{\text{statistic}} \left(\sum_{k, \nu_{\text{deg}}} \frac{1}{2} \hbar \omega_{k} |_{V,\Lambda, \mathcal{C}} - \sum_{k, \nu_{\text{deg}}} \frac{1}{2} \hbar \omega_{k} |_{V,\Lambda, \emptyset} \right) \end{split}$$

Gravity serious: "cosmological constant problem!"

The critical mass density $\rho_{\text{crit.}} = \frac{3H_0^2}{8\pi G_N} \approx h_0^2 \times 10.5 \,\text{GeVm}^{-3} = h_0^2 \times 4.8 \times 10^{-47} \,\text{GeV}^4$ compared with vacuum energy density $\rho_V = \frac{\Lambda_{cosm.}}{8\pi G_N} + \varepsilon$ where $\varepsilon = \frac{\mathcal{E}_0}{V} = \int \frac{d^3k}{(2\pi)^3} \frac{1}{2} \hbar \omega_k e^{-\hbar \omega_k / \Lambda_{UV}} \approx \frac{3\hbar c}{2\pi^2} \left(\frac{\Lambda_{UV}}{\hbar c}\right)^4$

Theory	Λ_{UV}	Scale [GeV]	$\varepsilon = \frac{1}{2} \int \hbar \omega''$	$\varepsilon/ ho_{ m crit.}$
ToE	M_P	1.2×10^{19}	$10^{76} { m ~GeV^4}$	10^{123}
	String	5×10^{17}	10^{72} GeV^4	10^{120}
GUT	$SU(5)_{SUSY}$	2×10^{16}	10^{65} GeV^4	10^{112}
	SU(5) _{min.}	10 ¹⁵	10^{60} GeV^4	10^{107}
	$SUSY_{break}$	10^{3}	10 ¹² GeV ⁴	10^{59}
EW	Higgs vacuum	$2.5 imes 10^2$	10^9 GeV^4	10^{56}
	Z, W	10^{2}	10^8 GeV^4	10^{55}
QCD	$\Lambda_{had.}$	1	1GeV ⁴	10^{47}
	Λ_{QCD}	0.2	$10^{-3} { m GeV^4}$	10^{44}
EM	$\hbar \omega_{\rm phot.}$	1 eV	10-36 GeV4	10 ¹¹
?	10-3 eV	$\sim 40K$	10-46 GeV4	1

Original Casimir effect:

Experiments: (< 12 until 1998)

- Early efforts: relied on cantilever or spring balances
 J.T.G. Overbeek & M.J. Sparnaay (1954); B.V. Deriagin & I.I. Abrikosova (1957): (glass or quartz!).

 M.J. Sparnaay (1958): first exp. with two *metal* plates; sensitivity: (0.1 1) ×10⁻⁸ N and 17 mV.
 Outcome: exponent n in the 1/Lⁿ Casimir -force determined as 4 ±1.

 P.H.G.M. van Blokland & J.T.G. Overbeek (1978): sphere-plate system, ~ 50% accuracy.
- Resurgence: (1996 →): sphere-plate systems under proximity force theorem: F^{ol}(L) =2πR hcπ²/720L³
 S.K. Lamoreaux, PRL 78 (1997) & PRL 81 (1998): torsion pendulum (60 cm !) mech. hysteresis eliminated; piezzoelectrical transducers; ΔF=10⁻¹¹ N, 10% accuracy.
 - U. Mohideen & A. Roy, PRL 81 (1998) and A. Roy et al., PRD 60 (1999): atomic force microscope; 32 - 1000 nm, $\Delta F = 10^{-12}$ N and 1-2 % accuracy.
 - H.B. Chan, F. Capasso et al., Science 291 (2001): (N.Y. Times, 9-Feb.-2001) capacitance bridge; seesaw device; Microelectromech. Systems (MEMS);L_{min} ≈ 70 nm.
 Th. Ederth, PRA 62 (2000): two coated (with hydrocarbon layers) crossed cylinders, 20-100 nm.
 G. Bressi et al., PRL 88 (2002): two 1.2×1.2 mm² plates, 0.5– 3 µm, 15 % accuracy.

Ideal case:

perfect conductors without impurities at T = 0.

Reality:

finite temperature (only important for $L > 1 \,\mu\text{m}$); imperfect conductors (realistic metals, Lifshitz theory); surface roughness (geometry averaging, $\langle r^2 \rangle_{\text{s.r.}}^{1/2} \approx 3 \,\text{nm}$)

Atomic Force Microscope:







M. Bordag, U. Mohideen, V.M. Mostepanenko, Phys. Rep. 353 (2001) and U. Mohideen and A. Roy, Phys. Rev. Lett. 81 (1998).



see H.B. Chan, V.A. Aksyuk, R.N. Kleiman, D.J. Bishop, F. Capasso, Science 291 (2001); 10 PRL 87 (2001) and Phys. Rev. Focus (Nov. 2, 2001).





11

 \Rightarrow stabilization if $C = \frac{e^2}{4\pi\hbar c}$ (indepently of the radius *a*).

 However, T. Boyer (1968): force is repulsive! (Similarly, R. Balian & C. Bloch (1972-74): Casimir calculations in cavities depend on (bag-)geometry)

.:. "Generalization" of concept of Casimir energy

"Generalization" of concept of Casimir energy:

1) geometry dependence

Casimir energy = vacuum energy from the geometry-dependent part of the density of states (d.o.s)

(↔ shell correction energy in Nucl. Phys.)

d.o.s.:
$$\rho(E) \equiv \sum_{E_k} \delta(E - E_k) = \rho_0(E) + \rho_{\text{bulk}}(E) + \delta \rho_C(E, \text{geom.-dep.})$$

N.o.s.:
$$\mathcal{N}(E) \equiv \sum_{E_k} \Theta(E - E_k) = \int_0^E dE' \,\rho(E')$$

Casimir Energy: $\mathcal{E}_{C} \equiv \int dE E \,\delta\rho_{C}(E, \text{geom.-dep.}) = -\int dE \,\mathcal{N}_{C}(E, \text{geom.-dep.})$

Further "Generalization" of concept of Casimir energy:

2) matter fields

- Here: space not "filled" with *fluctuating* EM modes, but with gas of non-interacting (non-relativistic) fermions.
- Similarity : ∃ mode sums ∑ħω_k with constant degeneracy factor, (because of Pauli's exclusion principle).
- Difference : ∃ of second scale: fermi energy = chemical potential μ (at T ≈ 0) in addition to geometric size and distance scale(s).
- Concrete: Matter fields (fermions) in the space between voids
 build up a quantum pressure on the voids

∃ effective *interaction* between empty regions of space in the background of *non-interacting* fermions

Applications

In the inner crust of a neutron star:

 nuclei start to loose neutrons due to increasing pressure and density: below saturation density ⇒ chain of phases between 0.03 fm⁻³ and 0.1 fm⁻³:

 $\textbf{nuclei} \rightarrow \textbf{rods} \rightarrow \textbf{slabs} \rightarrow \textbf{tubes} \rightarrow \textbf{bubbles} \rightarrow \textbf{uniform matter}$

- Liquid drop model: *meat-ball*, *spaghetti*, *lasagne* vacua from interplay between Coulomb and surface energies with phase differences of a few keV/fm³.
 - However, neglected shell correction energy of the same order ! (see A. Bulgac & P. Magierski, Nucl. Phys. A 683 (2001) 695.)

Inside a neutron/quark star:

• Quark matter droplets immersed in hadronic matter at $ho \gg
ho_{nm}$

In the lab:

- Buckyballs immersed in an electron gas (in liquid mercury)
- or Buckyballs immersed in liquid ³He.
- (Bosonic) cavities in dilute atomic Fermi condensates.

Calculation

• Casimir energy for fermions between two impenetrable (parallel) planes at distance L:

 $\mathcal{E}_C = \mu F(k_F L)$ with $\mu = \hbar^2 k_F^2 / 2m$ as natural scale.

- For more complicated geometries → involved computations !
- However, case of immersed non-overlapping spherical voids still relatively simple:

Krein (1953) trace-formula: level density var. $\delta \rho(E) \leftrightarrow \frac{d}{dE}$ of phase shift $\frac{1}{2i} \ln \det S_n(E)$

 $\delta\rho(E) = \bar{\rho}(E) - \bar{\rho}_0(E) = \frac{1}{2\pi i} \frac{d}{dE} \operatorname{tr} \ln S_n(E) \quad \text{of the n-sphere S-matrix}$

Extract geometry-dependent Casimir fluctuations from multiple-scattering part

⇒ Calculation mapped onto a quantum mechanical *billiard* problem: *hyperbolic point-particle scattering* off *n* spheres or *n* disks



References:

- B. Eckhardt, J. Phys. A20 (1987);
- P. Gaspard & S. Rice, J. Chem. Phys. 90 (1989);
- P. Cvitanović & B. Eckhardt, Phys. Rev. Lett. 63 (1989);
- M. Henseler, A. Wirzba & T. Guhr, Ann. Phys. 258 (1997).

Digression 1: E.Beth & G.E. Uhlenbeck (1937)

predecessor of Krein (1962) formula for spherical potentials:

Idea: spherical boxes: $\lim_{R\to\infty} \left[\left(\begin{array}{c} \\ \\ \end{array} \right) - \left(\begin{array}{c} \\ \end{array} \right) \right]$

Asymptotically:

$$u_{k\ell}(r) \sim \sin\left(kr - \frac{1}{2}\,\ell\pi + \eta_{\ell}(k)\right)$$
$$u_{k\ell}^{(0)}(r) \sim \sin\left(kr - \frac{1}{2}\ell\pi\right)$$

and Dirichlet b.c.'s: $u_{k\ell}(R) = u_{k\ell}^{(0)}(R) = 0.$

 $\begin{array}{l} \Rightarrow \\ \begin{array}{l} \mathsf{EV} \text{ conditions } ((2\ell+1)\text{-fold degenerate}): \\ \\ \Rightarrow \\ \begin{array}{l} k_{\ell,n}R - \frac{1}{2}\ell\pi + \eta_{\ell}(k_{\ell,n}) &= \pi n, \quad n=0,1,2,\cdots \quad (\text{with potential}) \\ \\ k_{\ell,n}^{(0)}R - \frac{1}{2}\ell\pi &= \pi n, \quad n=0,1,2,\cdots \quad (\text{without potential}) \end{array} \end{array}$

A change of n by one unit, for ℓ fixed, leads to

$$\Delta k_{\ell} \left(R + \frac{\partial}{\partial k} \eta_{\ell}(k) \right) = \pi = \Delta k_{\ell}^{(0)} R \,,$$

such that the conditions $\bar{\rho}_{\ell}(k)\Delta k_{\ell} = \bar{\rho}_{\ell}^{(0)}(k)\Delta k_{\ell}^{(0)} = 2\ell + 1$ (note the averaging !)

induce the formula

$$\bar{\rho}_{\ell}(k) - \bar{\rho}_{\ell}^{(0)}(k) = \frac{2\ell+1}{\pi} \frac{\partial}{\partial k} \eta_{\ell}(k)$$

Digression 2: Semiclassical interpretation of Krein formula

Determinant of scattering matrix is semiclassically a sum over periodic orbits (+Weyl terms)

Consider the difference of the densities of states of two bounded reference systems :



- The container-induced periodic orbits cancel!
- However, \exists further *spurious periodic orbits* whose lengths grow with increasing *R*.
- Removal of long orbits by exponential damping or averaging:

$$\lim_{\epsilon \to 0_+} \lim_{R \to \infty} \left\{ \rho^{(n)}(k + i\epsilon; R) - \rho^{(0)}(k + i\epsilon; R) \right\}$$
$$= \left. \frac{1}{2\pi} \operatorname{Im} \frac{\mathrm{d}}{\mathrm{d}k} \ln \det \left. \mathbf{S}^{(n)}(k) \right|_{k \text{ real}}$$

Note the order of the limits!

Digression 3: Casimir energy in fermionic background

chem. potential $\mu = \hbar^2 k_F^2 / 2m$ or Fermi momentum k_F natural UV-cutoff ($\Lambda_{UV} = \mu$ or $k_{UV} = k_F$)

Casimir energy at *fixed* total number of fermions \mathcal{N} :

 \Rightarrow chem. potentials $\mu \& \mu_0$ at *finite* and *infinite* separation of the scatterers <u>differ</u>:

$$\mathcal{N} = \int_{0}^{\mu} \rho(E) dE = \int_{0}^{\mu_{0}} [\rho_{0}(E) + \rho_{W}(E)] dE$$

$$\Rightarrow \qquad \underbrace{\int_{\mu_{0}}^{\mu} \rho(E) dE}_{\mu_{0}} = -\int_{0}^{\mu_{0}} \underbrace{[\rho(E) - \rho_{0}(E) - \rho_{W}(E)]}_{\rho_{C}(E) = \frac{d}{dE} \mathcal{N}_{C}(E)} dE$$

$$\mathcal{E}_{C}|_{\mathcal{N}} = \int_{0}^{\mu} E\rho(E) dE - \int_{0}^{\mu_{0}} E[\rho_{0}(E) + \rho_{W}(E)] dE$$

$$= \int_{0}^{\mu_{0}} E\rho_{C}(E) dE + \mu_{0} \underbrace{\int_{\mu_{0}}^{\mu} \rho(E) dE}_{\mu_{0}} + \mathcal{O}(V^{-1})$$

$$= \int_{0}^{\mu_{0}} (E - \mu_{0}) \rho_{C}(E) dE = -\int_{0}^{\mu_{0}} \mathcal{N}_{C}(E) dE.$$

Grandcanonical Casimir energy at fixed $\mu = \mu_0$: $\widetilde{\mathcal{E}_C}|_{\mu} - \mu \mathcal{N}_C(\mu) = \mathcal{E}_C|_{\mathcal{N}} + \mathcal{O}(V^{-1})$

- 1. *"infinite" container:* $\rho(E) = \rho_0(E)$ (fermionic background)
- 2. n bubbles (of radii a_i) "punched out" at "infinite" separation:

 $\rho(E) = \rho_0(E) + \sum_{i=1}^n \underbrace{\rho_W(E, a_i)}_{Weyl \ term} \quad \text{(note the excluded volume !)}$

3. geometry-dependent arrangement of n bubbles:

$$\rho(E) = \rho_0(E) + \sum_{i=1}^n \rho_W(E, a_i) + \delta \rho_C(E, \{a_i\}, \{\mathbf{r}_{ij}\})$$

- 4. Krein formula (note the averaging): $\delta\rho(E) = \bar{\rho}(E) - \bar{\rho}_{0}(E) = \frac{1}{2\pi i} \frac{d}{dE} \underbrace{\operatorname{In det} S_{n}(E)}_{\operatorname{In det} S_{n}(E)}$ $\underbrace{\delta\rho(E)}_{\operatorname{In det} S_{n}(E)} = \prod_{i} \operatorname{det} S_{i}(E, a_{i}) \frac{\operatorname{det} M^{\dagger}(k^{*})}{\operatorname{det} M(k)}$ $\hookrightarrow \delta\bar{\rho}_{C}(E, \{a_{i}\}, \{\mathbf{r}_{ij}\}) = -\frac{1}{\pi} \operatorname{Im} \left(\frac{d}{dE} \operatorname{In det} M(E)\right) \qquad (\text{see A.W., Phys. Rep. 309 (1999)})$
- 6. Casimir energy: $\therefore \ \mathcal{E}_{C} = \int_{0}^{\mu} dE \left(E - \mu \right) \delta \bar{\rho}_{C} = -\int_{0}^{\mu} dE \, \overline{\mathcal{N}}_{C}$



20 *Multi-scattering* matrix for n spheres of radii a_j and distances $r_{jj'}$ (j, j'=1, 2, ..., n) $M_{lm,l'm'}^{jj'} = \delta^{jj'} \delta_{ll'} \delta_{mm'} + (1 - \delta^{jj'}) i^{2m+l'-l} \sqrt{4\pi (2l+1)(2l'+1)} \left(\frac{a_j}{a_{j'}}\right)^2 \frac{j_l(ka_j)}{h_{ij}^{(1)}(ka_{j'})}$ $\times \sum_{\bar{l}=0}^{\infty} \sum_{\bar{m}=-l'}^{l'} \sqrt{2\tilde{l}+1} \; i^{\bar{l}} \begin{pmatrix} \tilde{l} & l' & l \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \tilde{l} & l' & l \\ m-\tilde{m} & \tilde{m} & -m \end{pmatrix} D_{m',\tilde{m}}^{l'}(j,j') \, h_{\bar{l}}^{(1)}(kr_{jj'}) \, Y_{\bar{l}}^{m-\tilde{m}}(\hat{r}_{jj'}^{(j)})$ M. Henseler, A. Wirzba & T. Guhr, Ann. Phys. 258 (1997) 286. $M^{jj'}(E) \approx \delta^{jj'} - \left(1 - \delta^{jj'}\right) \underbrace{f_j(E)}_{r_{jj'}} \frac{\exp(ikr_{jj'})}{r_{jj'}} \quad (+p\text{-wave})$ In the limit of small scatterers: Two spheres of radius a at distance rin the small-scatterer limit: $\mathcal{N}_C^{\text{oo}}(E) = -\frac{1}{\pi} \operatorname{Im} \ln \det M^{\text{oo}}(E) \approx \nu_{\text{deg}} \frac{a^2}{\pi r^2} \sin[2(r-a)k] + \mathcal{O}\left((ka)^3\right),$

whereas the semiclassical result (for the single two-bounce periodic orbit with no repetions) reads:

$$\mathcal{N}_{C,scl}^{\rm oo}(E) = \nu_{\rm deg} \frac{a^2}{4\pi r(r-2a)} \sin[\underbrace{2(r-2a)k}_{S_{po}(k)/\hbar}] \qquad (\text{Gutzwiller's trace formula})$$



N.o.S $\mathcal{N}_{C}^{oo}(E) = -\frac{1}{\pi} \operatorname{Im} \ln \det M^{oo}(E) \approx \nu_{\deg} \frac{a^2}{4\pi r(r-2a)} \sin[2(r-2a)k]$



(of longer range than molecular van der Waals forces !)

Sphere-plate geometry:
$$\mathcal{E}_C^{\text{ol}} \approx -\nu_{\text{deg}} \, \mu \frac{a}{2\pi (r-a)} j_1[2(r-a)k_F], \quad (k_F a > 1)$$

Three and four spheres:

- periodic orbit summation: ∃ of genuine three and more-body interactions
- However, 2-bounce orbit dominates in equilateral three- and four sphere systems (max. correction due to 3-bounce orbit is ~ 10 % at r≈2.5a)



Billiard analogy : difficult to make long shots, especially with many bounces

 the slightest error ruins the shot.

Dominance of two-body interactions:

the Casimir energy satisfies the rule $E_3 \approx 3E_2$ for 3 identical spheres (equilateral triangle)

and

 $E_4 \approx 6E_2 \approx 2E_3$ for 4 identical spheres (symmetric tetrahedron) if $k_F a \gg 1$.

 $(k_F a \le 1)$: corrections up to 10% and 25% for 3 & 4 spheres)



What happens at the boundary of a normal and superfluid regions?





Fermionic Casimir Effect of two superfluid grains in a normal Fermi gas



Specular reflection replaced by Andreev reflection \Rightarrow particle retro-reflected as a hole $k_p^2/2 + k_h^2/2 = k_F^2$ (particle/hole states symmetric) $k_p \sin \theta_p = k_h \sin \theta_h$ (modified Snell's law), \Rightarrow disks/spheres have focusing effect



stable orbits if
$$\frac{R}{2a} \leq \frac{k_p}{k_p - k_h}$$
 satisfied
(here $R = 6a$ and $k_p/k_h = 1.5$)

Apply

- 1. Bogoliubov-de Gennes equations with pairing field $\Delta_{1,2}(\vec{r}) = \Delta e^{\pm i\phi_{\Delta}/2}$ at circular grain 1 and 2,
- 2. mod. Krein formula $\delta N(E) = -\frac{1}{\pi} \ln \det M(E)$ with M(E) multi-scattering matrix in p-h space,
- 3. asymptotic approximation valid for large separations $R \gg a$

$$\begin{split} \delta N(E) &\approx \quad \frac{4|T_{ph}(a,\mu,\Delta)|\cos\phi_{\Delta}}{\pi^2\sqrt{k_hk_p}R}\sin[(k_p-k_h)R+\phi_{ph}] \\ &\quad -\frac{2|T_{pp}(a,\mu,\Delta)|}{\pi^2k_pR}\cos(2k_pR+\phi_{pp}) + \frac{2|T_{hh}(a,\mu,\Delta)|}{\pi^2k_hR}\cos(2k_hR-\phi_{hh}) \end{split}$$

 $T_{ph} = \left(\sum_m t_{ph}^m/2\right)^2$, $T_{pp} = \left(\sum_m (-1)^m t_{pp}^m/2\right)^2$ etc. structure functions (only dep. on single scatterer)





Borrowed from www.lsw.uni-heidelberg.de/~mcamenzi/NS_Mass.html

Quantum Corrections to the GS Energy of Inhomogeneous NM







The Casimir energy for various phases. The lattice constants are: L = 23, 25 and 28 fm respectively. u - anti-filling factor $\Box_0 - average density$

A. Bulgac and P. Magierski Nucl. Phys. 683, 695 (2001) Nucl. Phys, 703, 892 (2002) (E)

The Casimir energy for the displacement of a single void in the lattice













A. Bulgac and P. Magierski Nucl. Phys. 683, 695 (2001) Nucl. Phys. 703, 892 (2002) (E)

Deformation of the rod-like phase lattice





keV/fm³

 $\alpha\beta\sin\gamma=1$ volume conservation

A. Bulgac and P. Magierski Nucl. Phys. 683, 695 (2001) Nucl. Phys, 703, 892 (2002) (E)



FIG. 2. The total energy density (1) of *npe* matter as a function of the proton quadrupole moment $Q_p = Q_{20}^p$ (middle subfigures). The integrated proton and neutron densities (see text for definition) corresponding to nuclear configurations indicated by arrows are shown in the lower and upper subfigures.

Skyrme HF with SLy4, Magierski and Heenen, Phys. Rev. C 65, 045804 (2002)



FIG. 3. The same as in the Fig. 2 but for different densities.

Skyrme HF with SLy4, Magierski and Heenen, Phys. Rev. C 65, 045804 (2002)



FIG. 4. The energy density difference $\Delta E/V$ between nuclear phases as a function of the total density. Solid curve denotes the difference between the spherical and rodlike phase. Dotted curve denotes the difference between the spherical and slablike phase. Smaller subfigures show the energy density of *npe* matter as a function of the proton quadrupole moment for four different densities. Parameter *d* denotes the length of the cubic box.

ΔE between spherical and rod-like phases

ΔE between spherical and slab-like phases

Skyrme HF with SLy4, Magierski and Heenen, Phys. Rev. C 65, 045804 (2002)

"Spherical" phase (scc)

Rod-like phase



Size of box d = 26 fm

P. Magierski, A. Bulgac and P.-H. Heenen, nucl-ph/0112003

Rod-like phase

"Spherical" phase (bcc)



Size of box d = 23.4 fm

P. Magierski, A. Bulgac and P.-H. Heenen, nucl-ph/0112003

Slab-like phase

Bubble-like phase



Size of box d = 20.8 fm

P. Magierski, A. Bulgac and P.-H. Heenen, nucl-ph/0112003



Various contributions to the energy density as a function of the proton quadrupole moment.

Magierski, Bulgac and Heenen, Nucl.Phys. A719, 217c (2003)

