

**UW CSSS/POLS 512:
Time Series and Panel Data for the Social Sciences**

Panel Data Models with Many Time Periods

Christopher Adolph

Department of Political Science

and

Center for Statistics and the Social Sciences

University of Washington, Seattle

Plan for today

Estimating deterministic trends in panel data

Performing panel unit root tests

Estimating linear panel models with large(ish) T

Estimating Deterministic Trends

Recall that for a single time series,
we estimated the following model to capture a deterministic trend:

$$y_t = t\theta_1 + \mathbf{x}_t\boldsymbol{\beta} + \varepsilon_t$$

For panel data,
if we assume a *common* deterministic trend across units we can estimate:

$$y_{it} = t\theta_1 + \mathbf{x}_{it}\boldsymbol{\beta} + \varepsilon_{it}$$

We could also add in ARMA terms if we like,
as they capture distinct time series dynamics

As before, the question is whether we can trust $\hat{\theta}$ to estimate the trend well

With a single, short time series,
 $\hat{\theta}$ will be unbiased but often far from the truth and frequently incorrectly signed

Let's see whether assuming a common trend within a *panel* helps

Estimating Deterministic Trends

Let's see whether assuming a common trend within a panel helps


Revisit our Monte Carlo experiment to see how well $\hat{\beta}_1$ estimates β_1 in practice.

We set the true model to:

$$y_{it} = \beta_0 + \beta_1 t + \varepsilon_{it}, \quad \varepsilon_{it} \sim \mathcal{N}(0, \sigma^2)$$
$$y_{it} = 0 + 0.1t + \varepsilon_{it}, \quad \varepsilon_{it} \sim \mathcal{N}(0, 1)$$

Then, for each $t \in \{2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 50, 100\}$, we draw 100,000 datasets from this “true” model and see how well we estimate $\beta_1 = 0.1$

We repeat the experiment for $N = \{1, 2, 5, 10, 20, 50, 100\}$ and compare results




Recall that for a single time series,
we got very unreliable estimates and frequent sign errors at $T = 10$ (dotted red line)

Not biased: the mean of the distribution is centered on the truth, $\beta_1 = 0.1$,
but not reliable in any specific dataset

More data helped: vastly more efficient estimates at $T = 100$ (purple spike)


What happens if we had 100 observations, but spread across units?



What happens if we had 100 observations, but spread across units?

The dotted red line still marks the performance of trend estimation with a single time series of 10 periods

We can improve quite a bit by pooling 10 cross-sections in the same model




Note the 10 cross-sections of 10 periods case is not as efficient as 100 observations in a single series


Choices:

either assume the series has a very long stable trend (and find a long data set)


or assume different cross-sections follow the same trend (and pool across short time periods)




Let's get more systematic. Recall these Monte Carlo results for a single time series




Efficient trend estimation gets much easier with more units and fixed T




When N is really large – e.g., individual data in the 1000s – can efficiently estimate trends in very small T , provided pooling assumption is valid



Recall how frequently we got the direction of the trend wrong with a single, short series



Sign errors rapidly diminish with higher N and even modest T



But you might want to check your own case in Monte Carlos that match your assumptions

Panel Unit Root Tests

Deterministic trend estimation was hard in single time series, but easier when we pooled $N > 1$ time series together and assumed a common trend

Unit root tests were underpowered in a single time series. . .

Perhaps pooling time series and assuming degree of stationarity will help?

Panel unit root tests do exactly this, and are considerably more powerful:

Most assume balanced panels

(and R implementation in `plm::purtest` seems to impose this globally)

See also the R package `punitroots`

Panel Unit Root Tests

Generally, these tests find ways to combine the results from individual ADF tests on the different series in different ways

Most treat nonstationarity as the null hypothesis

A few well-known tests:

Im-Pesaran-Shin (2003) test: Pools the units; allows different AR processes.

LevinLinChu (2002) test: Pools the units; assumes common AR process.

Maddala-Wu (1999) test: Pools the p -values from separate ADFs; more flexible

Hadri (2000) test: Also more flexible test; null is stationarity

Although performance is better than single-series ADF, panel unit root tests make a variety of complex identifying assumptions, and those assumptions can lead to different results

Care is still indicated in using these tests:
they aren't foolproof or guaranteed to be right

Estimating Linear Panel Models

Last time, we discussed how including random and/or fixed effects changes the properties of our estimators of β

In this lecture, we'll talk about how to estimate and interpret panel models using fixed and/or random effects

And how to decide if we need (or even can use) fixed effects

We can always add random effects, but in some cases FEs either be too costly to estimate (in terms of dfs), or simply impossible to estimate

We will consider first the small N , large T case, which allows more complex time series modeling

Then the large N , small T case which raises the possibility of bias in fixed effects estimation

Finally, we consider heteroskedasticity in time or across panel structures

Estimating Fixed Effects Models

Option 1: Fixed effects or “within” estimator:

$$y_{it} - \bar{y}_i = (x_{it} - \bar{x}_i)\beta + (u_{it} - \bar{u}_i)$$

- estimating the fixed effects by differencing them out before applying least squares
- including time-invariant variables directly in x_{it} impossible here
- (rarely usable workaround: if we have an instrument for the time-invariant variable that is uncorrelated with the fixed effects; see Hausman-Taylor)
- suggests a complementary “between” estimator of \bar{y}_i on \bar{x}_i which could include time-invariant x_i ; together these models explain the variance in y_{it}
- does not actually provide estimates of the fixed effects themselves; just purges them from the model to remove omitted time-invariant variables
- to recover the fixed effects, could compute $\hat{\alpha}_i = y_{it} - x_{it}\hat{\beta}$

Estimating Fixed Effects Models

Option 2: Dummy variable estimator (sometimes called LSDV)

$$y_{it} = x_{it}\beta + \alpha_i + u_{it}$$

- also estimated by least squares (hence Least Squares Dummy Variable)
- yields estimates of α_i fixed effects (may be useful in quest for omitted variables; see if the α_i look like a variable you know)
- for large T , should be very similar to FE estimator
- not a good idea for very small T : estimates of α_i will be poor

Time-Invariant Covariates & Fixed Effects

We can't include time-invariant variables in fixed effects models

If we do, we will have perfect collinearity, and can't get estimates

That is, we will get some parameter estimates equal to NA

Never report a regression with NA parameters

The regression you tried to run was impossible. Start over with a possible one.

Time-Invariant Covariates & Fixed Effects

If we can't include time-invariant variables in a fixed effects model, does that mean time-invariant variables can never explain changes over time?

You might think so: how can a constant explain a variable?

But time-invariant variables could still effect time-varying outcomes in a special way. . .

Time-invariant variables can influence how a unit weathers time-varying *shocks* in some other variable

Example: labor market regulations (e.g. employment protection) don't change much over time

Blanchard & Wolfers found that when a negative economic shock hits, unemployment may rebound more slowly where such protections are stronger

Time-Invariant Covariates & Fixed Effects

We can model how a slow moving or time-invariant covariate conditions the effect of a quickly changing covariate on y_{it}

To estimate how a time-invariant covariate x_i mediates the effect of a shock, s_{it} , include on the RHS $x_i \times s_{it}$ and s_{it} , while omitting x_i itself

(It's okay *and necessary* to omit the x_i base term in this special case, because α_i already captures the effect of x_i)

Many theories about institutions can be tested this way

Time-Invariant Covariates & Fixed Effects

What if we want to “include” time-invariant covariates’ effect on the long term average level of y ?

We might partition the fixed effect into:

1. the portion “explained” by known time-invariant variables and
2. the portion still unexplained

Plümper & Troeger have methods to do this.

In this case, our estimates of the time-invariant effects are vulnerable to omitted variable bias from unmeasured time-invariant variables, even though time varying variables in the model are not

Thus you now need to control for *lots* of time-invariant variables directly, even hard to measure ones like culture

Estimating Random & Mixed Effects Models

Estimation of random effects is by maximum likelihood (ML) or generalized least squares (GLS)

Technically we're just adding one parameter to estimate: the variance of the random effects, σ_{α}^2

This is partitioned out of the overall variance, σ^2

Can understand this most easily by abstracting away from time series for a moment

Estimating Random & Mixed Effects Models

Recall that for linear regression, we assume homoskedastic, serially uncorrelated errors, and thus a variance-covariance matrix like this:

$$\Omega = \begin{bmatrix} \sigma^2 & 0 & 0 & 0 \\ 0 & \sigma^2 & 0 & 0 \\ 0 & 0 & \sigma^2 & 0 \\ 0 & 0 & 0 & \sigma^2 \end{bmatrix}$$

Estimating Random & Mixed Effects Models

And recall that heteroskedastic (but serially uncorrelated) errors have this variance-covariance matrix

$$\Omega = \begin{bmatrix} \sigma_1^2 & 0 & 0 & 0 \\ 0 & \sigma_2^2 & 0 & 0 \\ 0 & 0 & \sigma_3^2 & 0 \\ 0 & 0 & 0 & \sigma_4^2 \end{bmatrix}$$

Estimating Random & Mixed Effects Models

And finally, remember heteroskedastic, serially correlated errors follow this general form of variance-covariance

$$\Omega = \begin{bmatrix} \sigma_1^2 & \sigma_{12} & \sigma_{13} & \sigma_{14} \\ \sigma_{21} & \sigma_2^2 & \sigma_{23} & \sigma_{24} \\ \sigma_{31} & \sigma_{32} & \sigma_3^2 & \sigma_{34} \\ \sigma_{41} & \sigma_{42} & \sigma_{43} & \sigma_4^2 \end{bmatrix}$$

What does this matrix look like for random effects with no serial correlation?

Estimating Random & Mixed Effects Models

Define the variance of the random effect as

$$E(\alpha_i^2) = \sigma_\alpha^2 = \text{var}(\alpha_i)$$

Define the expected value of the squared white noise term as σ_ε^2

$$E(\varepsilon_{it}^2) = \sigma_\varepsilon^2 = \text{var}(\varepsilon_{it})$$

White noise is serially uncorrelated, so has covariance 0 for $t \neq s$:

$$E(\varepsilon_{it}\varepsilon_{is}) = 0 = \text{cov}(\varepsilon_{it}, \varepsilon_{is})$$

Finally, note that we assumed the white noise error and random effect are uncorrelated,

$$E(\alpha_i\varepsilon_{it}) = 0 = \text{cov}(\alpha_i, \varepsilon_{it})$$

Estimating Random & Mixed Effects Models

Thus the variance of the whole random component of the model is

$$\begin{aligned} \mathbf{E}((\alpha_i + \varepsilon_{it})(\alpha_i + \varepsilon_{it})) &= \mathbf{E}(\alpha_i^2) + 2\mathbf{E}(\alpha_i\varepsilon_{it}) + \mathbf{E}(\varepsilon_{it}^2) \\ &= \sigma_\alpha^2 + 0 + \sigma_\varepsilon^2 \\ &= \sigma_\alpha^2 + \sigma_\varepsilon^2 \end{aligned}$$

And the covariance of the whole random component is:

$$\begin{aligned} \mathbf{E}((\alpha_i + \varepsilon_{it})(\alpha_i + \varepsilon_{is})) &= \mathbf{E}(\alpha_i^2) + \mathbf{E}(\alpha_i\varepsilon_{is}) + \mathbf{E}(\alpha_i\varepsilon_{it}) + \mathbf{E}(\varepsilon_{it}\varepsilon_{is}) \\ &= \sigma_\alpha^2 + 0 + 0 + 0 \\ &= \sigma_\alpha^2 \end{aligned}$$

Estimating Random & Mixed Effects Models

If our data have a single random effect in the mean for each unit
→ serially correlated errors, but expressible using only two variances:

- the random effects variance σ_α^2
- the white noise term's variance σ_ε^2

$$\Omega = \begin{bmatrix} \sigma_\alpha^2 + \sigma_\varepsilon^2 & \sigma_\alpha^2 & \sigma_\alpha^2 & \sigma_\alpha^2 \\ \sigma_\alpha^2 & \sigma_\alpha^2 + \sigma_\varepsilon^2 & \sigma_\alpha^2 & \sigma_\alpha^2 \\ \sigma_\alpha^2 & \sigma_\alpha^2 & \sigma_\alpha^2 + \sigma_\varepsilon^2 & \sigma_\alpha^2 \\ \sigma_\alpha^2 & \sigma_\alpha^2 & \sigma_\alpha^2 & \sigma_\alpha^2 + \sigma_\varepsilon^2 \end{bmatrix}$$

Estimating Random & Mixed Effects Models

We have drastically simplified this matrix, and can now use FGLS (Feasible Generalized Least Squares) or ML to estimate it

$$\hat{\beta}_{\text{GLS}} = \left(\sum_{i=1}^N X_i' \Omega^{-1} X_i \right)^{-1} \left(\sum_{i=1}^N X_i' \hat{\Omega}^{-1} y_i \right)$$

where X_i is the $T \times K$ matrix of covariates for unit i , all times $t = 1, \dots, T$, and all K covariates

All we need are the estimates $\hat{\sigma}_\alpha^2$ and $\hat{\sigma}_\varepsilon^2$, and we can calculate $\hat{\beta}_{\text{GLS}}$

Estimating Random & Mixed Effects Models

We get $\hat{\sigma}_\varepsilon^2$ from the residuals from a LS regression:

$$\hat{\sigma}_\varepsilon^2 = \frac{1}{NT - K} \sum_{i=1}^N \sum_{t=1}^T \hat{\varepsilon}_{it\text{LS}}^2$$

(This is the usual estimator, but for NT observations)

Estimating Random & Mixed Effects Models

To get an estimator of $\hat{\sigma}_\alpha^2$, we need to adjust for the fact that we have only so many unique pairs of errors to compare:

$$\begin{aligned}\hat{\sigma}_\alpha^2 &= \mathbf{E} \left(\sum_{t=1}^{T-1} \sum_{s=t+1}^T (\alpha_i + \varepsilon_{it})(\alpha_i + \varepsilon_{is}) \right) \\ &= \mathbf{E} \sum_{t=1}^{T-1} \sum_{s=t+1}^T ((\alpha_i + \varepsilon_{it})(\alpha_i + \varepsilon_{is})) \\ &= \sum_{t=1}^{T-1} \sum_{s=t+1}^T \sigma_\alpha^2 \\ &= \sigma_\alpha^2 \sum_{t=1}^{T-1} (T - t) \\ &= \sigma_\alpha^2 ((T - 1) + (T - 2) + \dots + 2 + 1)\end{aligned}$$

Estimating Random & Mixed Effects Models

$$\begin{aligned} &= \sigma_{\alpha}^2((T - 1) + (T - 2) + \dots + 2 + 1) \\ &= \sigma_{\alpha}^2 T(T - 1)/2 \end{aligned}$$

$$\hat{\sigma}_{\alpha}^2 = \frac{1}{NT(T - 1)/2 - K} \sum_{i=1}^N \sum_{t=1}^{T-1} \sum_{s=t+1}^T \hat{\epsilon}_{it} \hat{\epsilon}_{is}$$

where in the last step we replace σ_{α}^2 with its estimator from pooled LS (the average of the products of the unique pairs of residuals)

With some algebra, this approach extends to serial correlaton of other kinds (ARMA)

For complex models, with many levels and/or hyperparameters, best to go Bayesian, set diffuse priors on the parameters, and use MCMC

Selecting Fixed Effects vs Random Effects Models

Choosing random effects when α_i is actually correlated with x_{it} will lead to omitted variable bias

Choosing fixed effects when α_i is really uncorrelated with x_{it} will lead to inefficient estimates of β (compared to random effects estimation) *and* kick out our time-invariant variables

Often in comparative we are certain there are important omitted time invariant variables (culture, unmeasured institutions, long effects of history)

So choice to include fixed effects requires nothing more than theory

Still could include random effects in addition to the fixed effects

Selecting Fixed Effects vs Random Effects Models

But if we are uncertain, or want to check against estimating unnecessary fixed effects, we can use the Hausman test for (any) fixed effects versus just having random effects

Hausman sets up the null hypothesis of random effects

Attempts to reject it in favor of fixed effects

Checks whether the random α_i 's are correlated with x_i under the null

Does this by calculating the variance-covariance matrices of regressors under FE and then just RE

Null is no correlation between these covariances

If there is no correlation, that means the regressors do not predict the random effects (ie, are uncorrelated)

Rejecting the null suggests you may need fixed effects to deal with omitted variable bias

`phptest` in `plm` library

Interpreting Random Effects Models

Usually, interest focuses on the percentage of variance explained by the random effects

And how this variance compares to that remaining in the model

Reported by your estimation routine

What if T is very small?

If T is very small (< 15 perhaps), estimating panel dynamics efficiently and without bias gets harder

In these cases, we should investigate alternatives:

1. First differencing the series to produce a stationary, hopefully white noise process
2. Including fixed effects for the time period (time dummies)
3. Checking for serial correlation after estimation (LM test)
4. Using lags of the dependent variable, while removing the bias from including lags with fixed effects by instrumenting with lagged differences (Arellano-Bond)

Example: GDP in a panel

Let's use the Przeworski et al democracy data to try out our variable intercept models

This exercise is for pedagogical purposes only; the models we fit are badly specified

We will investigate the following model:

$$\Delta^d \text{GDP}_{it} = \alpha_i + \beta_1 \text{OIL}_{it} + \beta_2 \text{REG}_{it} + \beta_3 \text{EDT}_{it} + \nu_{it}$$

- where $\nu_{it} \sim \text{ARMA}(p, q)$,
- d may be 0 or 1, and
- α_i may be fixed, random, or a mixed


Example: GDP in a panel

We first investigate the time series properties of GDP

But we have $N = 113$ countries! So we would have to look at 113 time series plots, 113 ACF plots, and 113 PACF plots


Fortunately, they do look fairly similar. . .

Country 1




GDP time series for country 1

Country 2




GDP time series for country 2

Country 3




GDP time series for country 3

Country 4




GDP time series for country 4

Country 113




GDP time series for country 113

Series GDPW[COUNTRY == currcty]




GDP ACF for country 1

Series GDPW[COUNTRY == currcty]




GDP ACF for country 2

Series GDPW[COUNTRY == currcty]




GDP ACF for country 3

Series GDPW[COUNTRY == currcty]




GDP ACF for country 4

Series GDPW[COUNTRY == currcty]




GDP ACF for country 113

Series GDPW[COUNTRY == currcty]




GDP PACF for country 1

Series GDPW[COUNTRY == currcty]




GDP PACF for country 2

Series GDPW[COUNTRY == currcty]




GDP PACF for country 3

Series GDPW[COUNTRY == currcty]




GDP PACF for country 4

Series GDPW[COUNTRY == currcty]



GDP PACF for country 113


Histogram of adftest.pvalues



Histogram of p -values from ADF tests on GDPW

What would we see if there were no unit roots?

Histogram of PPtest.pvalues



Histogram of p -values from Phillips-Peron tests on GDPW

Choosing AR(p,q) for panel

What do we think?


Clearly some heterogeneity

If had to pick one time series specification, choose ARIMA(0,1,0) or ARIMA(1,1,0)

Seems to fit many cases; guards against spurious regression


But if we're dubious about imposing a single ARIMA(p,d,q) across units, we could let them be heterogeneous

Country 1




GDPdiff time series for country 1

Country 2




GDPdiff time series for country 2

Country 3




GDPdiff time series for country 3

Country 4




GDPdiff time series for country 4

Country 113




GDP diff time series for country 113

Series GDPWdiff[COUNTRY == currcty]




GDPdiff ACF for country 1

Series GDPWdiff[COUNTRY == currcty]




GDPdiff ACF for country 2

Series GDPWdiff[COUNTRY == currcty]




GDPdiff ACF for country 3

Series GDPWdiff[COUNTRY == currcty]




GDPdiff ACF for country 4

Series GDPWdiff[COUNTRY == currcty]




GDPdiff ACF for country 113

Series GDPWdiff[COUNTRY == currcty]




GDPdiff PACF for country 1

Series GDPWdiff[COUNTRY == currcty]




GDPdiff PACF for country 2

Series GDPWdiff[COUNTRY == currcty]




GDPdiff PACF for country 3

Series GDPWdiff[COUNTRY == currcty]




GDPdiff PACF for country 4

Series GDPWdiff[COUNTRY == currcty]



GDPdiff PACF for country 113


Histogram of adftestdiff.pvalues



Histogram of p -values from ADF tests on GDPWdiff

What is this pattern consistent with?

Histogram of PPtestdiff.pvalues



Histogram of p -values from Phillips-Peron tests on GDPWdiff

Example continued in R demonstration

We will continue this example in section using the code provided

For now, let's focus on the results that emerge,
and how they depend on treating intercepts as either random or fixed by country

In particular, we want to see if fixed effects can help us with omitted time invariant variables, which are legion in this example


In the example, we will decide on an ARIMA(1,1,0) model of GDP
(What does this mean?)

We will fit three different models of the relationship between education and GDP:

1. ARIMA(1,1,0) with random country intercepts
and controls for oil producing countries and democracy
2. ARIMA(1,1,0) with fixed country intercepts
and controls for democracy
3. ARIMA(1,1,0) with "mixed" country intercepts
and controls for democracy


	Model				
	RE	FE	FE-pcse	FE	ME
Education _{<i>it</i>}	23.96	-75.56	-75.56	-86.68	-84.75
	5.59	12.16	13.48	12.45	14.56
Democracy _{<i>it</i>}	110.94	-12.90	-12.90	-26.15	-3.63
	34.63	47.69	50.58	47.97	55.22
Oil-Producer _{<i>it</i>}	-26.89	—	—	—	—
	44.84	—	—	—	—
GDP _{<i>i,t-1</i>}	0.23	0.15	0.15	0.17	0.20
		0.02	0.02	0.02	
GDP _{<i>i,t-2</i>}				-0.12	
				0.02	
σ_α	0.14	—	—	—	309.10
Fixed effects		x	x	x	x
Random effects	x				x
<i>N</i>	113	113	113	113	113
<i>T</i>	328	328	328	228	328
observed <i>N</i> × <i>T</i>	2794	2794	2794	2741	2794
AIC	43376				42112
LM test <i>p</i> -value		<0.001	<0.001	0.131	

Random effects ARIMA(1,1,0)



What does the model imply substantively, and how does this depend on model assumptions?


Random effects ARIMA(1,1,0)



Suppose we have a country with “average” characteristics, and we increase education to 1 sd above the mean


How much does the model predict education to rise over the following years?

Random effects ARIMA(1,1,0)



The above plot shows the expected change in GDP over time in the high education country relative to an average (untreated) country


Random effects ARIMA(1,1,0)



The result seems sensible, but the model:

(1) ignored many unmeasured confounders and (2) differences GDP, so we should be skeptical in both the short- and long-run

Fixed effects ARIMA(1,1,0)




Adding fixed effects completely flips the results for education

Now the results make little sense

(Suggests the model is badly identified, even with fixed effects)

Mixed effects ARIMA(1,1,0)



In a model with both RE and FE for countries, the FEs dominate, as the “Mixed” effects model shows