Homework 4. aka "The calm before the storm" *Due Friday Oct 20 at 5pm* in Prof. Ginger's mailbox. CIRCLE YOUR ANSWERS AND KEY INTERMEDIATE RESULTS *USE MAPLE WHENEVER POSSIBLE* STAPLE YOUR PAPERS TOGETHER INCLUDE ALL COMPUTER PRINTOUTS (with commentary)

You will note that this problem set is (much) shorter than usual because of the reduced time this week. I know this will disappoint many of you and I promise I'll try to make it up to you next week!

Levine Problems

4.27 vibrations in LiH and ICl

Additional Problems

1) In lecture we introduced the lowering operator as $a^- = \frac{1}{\sqrt{2\hbar\mu\omega}}(+ip + \mu\omega x)$

By arguing that energy can't be negative, we reasoned that there must be some state $|\psi_0\rangle$ for which $a|\psi_0\rangle=0$ Use this relationship to generate a first-order differential equation for ψ_0 and solve the differential equation (it should be a "simple" first-order equation) to verify that ψ_0 is the same as obtained with the power series solution.

2) Find ψ_l , the first excited state of the SHO, by explicitly applying the raising operator, a^+ to ψ_0

3) The frequencies of the three normal modes of H₂O are ω_1 =3833 cm⁻¹ ω_2 =1649 cm⁻¹ and ω_3 =3943 cm⁻¹. If we describe a vibrationally excited state by the notation (n₁n₂n₃) where n_i is the quantum number associated with the ith normal mode, what is the energy of the (121) state? What is the energy difference between the (112) and (010) state? Side note: the anharmonicity in real bonds tends to mix the normal modes over time.

4) The spatial Parity operator P satisfies the eigenvalue equation: $P \ \psi = p \psi$ where the eigenvalues of P are p=+1 (if ψ is even) and p=-1 (if ψ is odd). Only even and odd functions are eigenfunctions of P. The symmetry of the Hamiltonian has important consequences for the symmetry of the allowed wave functions that we will examine below.

4a) Show that if the Hamiltonian is a symmetric (even) function, then [P,H]=0 (hint: what is the parity of an even function times an even function, or an odd function times an odd function, i.e. $P(f_1f_2)=?$)

4b) Two operators will commute if and only if they have a simultaneous set of eigenfunctions. Use this fact, and your result from a) to justify that statement that "for a symmetric Hamiltonian, the only allowed energy eigenfunctions will be even and odd functions." The analysis of the spatial symmetry of a wavefunction is a very powerful tool that allows you to predict properties (i.e. IR and Raman activity of vibrational modes).