

Rods And Cones Use Different Proteins For Phototransduction

- Lerea, C. L., Somers, D. E., Hurley, J. B., Kloch, I. B. and Bunt-Milam, A. H. (1986) Identification of specific transducin α subunits in retinal rod and cone photoreceptors. *Science* **234**:77.
- Lerea, C. L., Bunt-Milam, A. H., Kloch, I. and Hurley, J. B. (1989) Transducin is present in red, green and blue sensitive photoreceptors in the human retina. *Neuron* **3**:367-376.
- Hamilton, S.E., Prusti, R. P., Bentley, J. K., Beavo, J. A., and Hurley, J. B. (1993) Affinities of bovine photoreceptor cGMP phosphodiesterases for rod and cone inhibitory subunits. *FEBS Lett.* **318**:157-161.
- Nathan, J., Reh, R., Ankoudinova, I., Andoudinova, G., Chang, B., Heckenlively, J. and Hurley, J.B. (2006). Scotopic and photopic visual thresholds and spatial and temporal discrimination evaluated by behavior analysis of mice in a water maze. *Photochem. Photobiol.* In press.
- Rosenzweig, D.H., Nair, K.S., Wei, J., Wang, Q., Garwin, G., Saari, J.C., Chen, C.K., Smrcka, A.V., Swaroop, A. Lem, J., Hurley, J.B., Slepak, V.Z. (2007). Subunit dissociation and diffusion determine the subcellular localization of rod and cone transducins. *J. Neurosci.* 2007. 16:5484-94.