

An Axiomatization of the Ratio/Difference Representation

JOHN M. MIYAMOTO

University of Michigan

If \geq_r and \geq_d are two quaternary relations on an arbitrary set A , a ratio/difference representation for \geq_r and \geq_d is defined to be a function f that represents \geq_r as an ordering of numerical ratios and \geq_d as an ordering of numerical differences. Krantz, Luce, Suppes and Tversky (1971, *Foundations of Measurement*. New York, Academic Press) proposed an axiomatization of the ratio/difference representation, but their axiomatization contains an error. After describing a counterexample to their axiomatization, Theorem 1 of the present article shows that it actually implies a weaker result: if \geq_r and \geq_d are two quaternary relations satisfying the axiomatization proposed by Krantz *et al.* (1971), and if $\geq_{r'}$ and $\geq_{d'}$ are the relations that are inverse to \geq_r and \geq_d , respectively, then either there exists a ratio/difference representation for \geq_r and \geq_d , or there exists a ratio/difference representation for $\geq_{r'}$ and $\geq_{d'}$, but not both. Theorem 2 identifies a new condition which, when added to the axioms of Krantz *et al.* (1971), yields the existence of a ratio/difference representation for relations \geq_r and \geq_d .

Garner (1954) suggested that one could determine a ratio scale for loudness if subjects are able to judge what stimulus is a given fraction as loud as another stimulus and if, in addition, they could partition a loudness interval into a given number of subintervals of equal subjective size. His proposal assumes that there are distinct mental operations that can be carried out on the subjective representation of loudness, the one operation being isomorphic to the calculation of numerical ratios and the second operation being isomorphic to the calculation of numerical differences. Torgerson (1961) doubted that there exist two distinct operations of loudness judgment. He proposed that even if subjects are instructed, on the one hand, to judge the magnitude of subjective ratios and, on the other hand, to judge the magnitude of subjective differences, the mental operations underlying their responses would be the same. Michael Birnbaum and his colleagues have carried out an extensive program of experimentation devoted to testing these opposing hypotheses. On the whole, their evidence supports the theory that only one mental operation underlies judgments of ratios and differences, although this conclusion has not been universally accepted.¹ Birnbaum (1978, 1982), Hagerty and Birnbaum (1978) and

I would like to thank David H. Krantz for many valuable discussions of these and related ideas. Requests for reprints should be sent to John M. Miyamoto, Department of Psychology, University of Michigan, Ann Arbor, Mi. 48104-1346.

¹ Birnbaum (1978, 1980, 1982) reviews evidence and arguments favoring the theory that only one

Veit (1978) have attempted to show that the sole underlying operation of magnitude judgment is a subtractive operation. Their arguments lie beyond the scope of the present discussion.

The purpose of this article is not to discuss these psychophysical issues, but rather to present an axiomatization of the theory that two mental operations of magnitude judgment exist. The following formulation of this theory is due to Krantz *et al.* (1971), although some of the terminology is my own. If there are distinct mental operations of magnitude judgment that are isomorphic, respectively, to the calculation of ratios and of differences, then subjects ought to order pairs of stimuli in two different ways depending on whether they are asked to judge the magnitude of subjective ratios or the magnitude of subjective differences. Furthermore, these two orderings of stimulus pairs ought to possess a common numerical representation as orderings, respectively, of ratios and of differences. This common representation will be called a ratio/difference representation and is described formally in Definition 1. As a notational convention, if y and z denote elements of a set A , then yz denotes the ordered pair (y, z) .

DEFINITION 1. Let \geq_r and \geq_d be quaternary relations on an arbitrary set A . Let Re^+ denote the positive real numbers and let $f: A \rightarrow \text{Re}^+$ be any function. Then, f is a ratio/difference representation for \geq_r and \geq_d iff for all $w, x, y, z \in A$

- (i) $wx \geq_r yz$ iff $f(w)/f(x) \geq f(y)/f(z)$,
- (ii) $wx \geq_d yz$ iff $f(w) - f(x) \geq f(y) - f(z)$.

Thus, a ratio/difference representation is a function f that simultaneously represents an ordering \geq_r as an ordering of numerical ratios and an ordering \geq_d as an ordering of numerical differences.

Before discussing the ratio/difference representation, it will be helpful to set down several conventions. First, $>_r$ and $>_d$ denote the strict inequalities and \sim_r and \sim_d denote the equivalence relations corresponding to \geq_r and \geq_d , respectively. Furthermore, there is a trivial case that it is convenient to exclude, namely, if $wx \sim_r yz$ and $wx \sim_d yz$ for all $w, x, y, z \in A$, then any constant, positive real function is a ratio/difference representation for \geq_r and \geq_d . Since this case is uninteresting, it will be assumed throughout this essay that \geq_r and \geq_d are nondegenerate in the sense that $st >_r uv$ and $wx >_d yz$ for some $s, t, u, v, w, x, y, z \in A$.

Krantz *et al.* (1971) propose an axiomatization of the ratio/difference represen-

mental operation underlies judgments of subjective ratios and subjective differences. The theory originates with Torgerson (1961). Studies supporting this theory include Birnbaum and Elmasian (1978), Birnbaum and Mellers (1978), Birnbaum and Veit (1974), Hagerty and Birnbaum (1978), Mellers, Davis, and Birnbaum (1984), Schneider, Parker, Farrell, and Kanow (1976) and Veit (1978, 1980). The theory that there are distinct judgments of subjective ratio and difference applying to a single mental representation has been defended by Rule and Curtis (1980) and Rule, Curtis, and Mullin (1981). Parker, Schneider, and Kanow (1975) provide evidence that distinct judgments of subjective ratio and subjective difference can be performed on the continuum of perceived line lengths. Marks (1974) and Stevens (1957, 1971) develop views that do not fall into either camp.

tation and construct a purported proof that the axiomatization is sufficient to establish the existence of the representation. Next, I will state their axiomatization, discuss its interpretation, and then describe a counterexample to the sufficiency of the axiomatization. Later, an analysis will be given that locates the error in the attempted proof of the representation theorem appearing in Krantz *et al.* (1971). The following definition makes use of the concept of an algebraic difference structure which is defined in Krantz *et al.* (1971, Definition 4.3).

DEFINITION 2. Let \geq_r and \geq_d be quaternary relations on an arbitrary set A . Then \geq_r and \geq_d satisfy generalized ratio/difference compatibility iff $(A \times A, \geq_r)$ and $(A \times A, \geq_d)$ are both algebraic difference structures and for all $x, y, z, x', y', z' \in A$,

- (i) $xy \geq_r xx$ iff $xy \geq_d xx$,
- (ii) if $xx' \sim_r yy' \sim_r zz'$, then $xy \geq_d yz$ iff $x'y' \geq_d y'z'$,
- (iii) \geq_r and \geq_d are not identical orderings.

Generalized ratio/difference compatibility is proposed by Krantz *et al.* (1971) as an axiomatization of the ratio/difference representation, although they do not use this terminology (cf. the hypotheses of their Theorem 4.3, p. 154).

The assumption that $(A \times A, \geq_r)$ and $(A \times A, \geq_d)$ are both algebraic difference structures is quite plausible, since the existence of a ratio/difference representation implies that both structures have difference representations (the logarithm of the ratio representation is a difference representation). Intuitively, condition (i) asserts that \geq_r and \geq_d determine the same set of positive intervals. Condition (ii) is rather complicated in appearance, but it implies the more easily interpreted condition,

- (ii') if $xx' \sim_r yy' \sim_r zz'$, then $xy \sim_d yz$ iff $x'y' \sim_d y'z'$.

Note that if $uv \sim_d vw$, then v may be regarded as the \geq_d midpoint between u and w , i.e., the midpoint in the \geq_d ordering. Thus, condition (ii') asserts that if xx' , yy' , and zz' are all in the same \geq_r equivalence class, then y is the \geq_d midpoint between x and z iff y' is the \geq_d midpoint between x' and z' . Condition (ii) is just an ordinal version of (ii'). Krantz *et al.* (1971) show that conditions (i) and (ii) of Definition 2 are necessary for the ratio/difference representation. To see that condition (iii) must hold, suppose that \geq_r and \geq_d were identical. If f were a ratio/difference representation for \geq_r and \geq_d , then $\log f$ would be a difference representation for \geq_r , so $\log f$ and f would be difference representations for \geq_d . By the uniqueness of the algebraic difference representation for $(A \times A, \geq_d)$, there must be constants α, β such that $\log f = \alpha f + \beta$. It can be shown that $f(A)$ is dense in an interval of real numbers² and consequently there exist $x, y, z \in A$ such that $f(x) \neq f(y) \neq f(z) \neq f(x)$. Since a linear relation between $\log f$ and f cannot be obtained if the range of f has more than two values, \geq_r and \geq_d cannot be identical.

² The set $f(A)$ is dense in an interval of real numbers iff for every x and z in that interval, $x < z$ implies that there exists $y \in f(A)$ such that $x < y < z$. The proof that $f(A)$ is dense in an interval of reals is given in Krantz *et al.* (1971, p. 159).

An important feature of generalized ratio-difference compatibility is that if it holds for a pair of relations \geq_r and \geq_d , then it must also hold for the corresponding inverse relations. In other words, define relations \geq_{r^*} and \geq_{d^*} by the conditions

$$\begin{aligned} wx \geq_{r^*} yz & \quad \text{iff} \quad yz \geq_r wx, \\ wx \geq_{d^*} yz & \quad \text{if} \quad yz \geq_d wx, \end{aligned}$$

for every $w, x, y, z \in A$. I claim that if \geq_r and \geq_d satisfy generalized ratio/difference compatibility, then \geq_{r^*} and \geq_{d^*} must also satisfy generalized ratio/difference compatibility. Assuming that \geq_r and \geq_d satisfy generalized ratio/difference compatibility, it is easy to check that $(A \times A, \geq_{r^*})$ and $(A \times A, \geq_{d^*})$ are algebraic difference structures and that \geq_{r^*} and \geq_{d^*} satisfy conditions (i) and (iii) of Definition 2. As for condition (ii), the pairs xx' , yy' and zz' are in the same \geq_r equivalence class iff they are in the same \geq_{r^*} equivalence class. So the antecedent of (ii) holds with respect to \geq_r iff it holds with respect to \geq_{r^*} . The consequent of (ii) asserts that $xy \geq_d yz$ iff $x'y' \geq_d y'z'$, but this is equivalent to asserting that $xy \geq_{d^*} yz$ iff $x'y' \geq_{d^*} y'z'$. So if the consequent of (ii) holds with respect to \geq_d , it also holds with respect to \geq_{d^*} .

Thus, the definition of generalized ratio/difference compatibility possesses a kind of symmetry with respect to the relations \geq_r and \geq_d , and the inverse relations \geq_{r^*} and \geq_{d^*} . This symmetry is the crucial defect in the conjecture that generalized ratio/difference compatibility implies the existence of a ratio/difference representation. Consider the following counterexample to this conjecture. Let $A = \text{Re}^+$. Define relations R and D by the conditions: for any $w, x, y, z \in \text{Re}^+$,

$$\begin{aligned} wx R yz & \quad \text{if} \quad w/x \geq y/z, \\ wx D yz & \quad \text{if} \quad w - x \geq y - z, \end{aligned}$$

where $/$ and $-$ denote division and subtraction of real numbers. Obviously, R and D satisfy generalized ratio/difference compatibility and the identity function $I: \text{Re}^+ \rightarrow \text{Re}^+$ is a ratio/difference representation for R and D . Let R^* and D^* denote the relations inverse to R and D . By a previous argument, R^* and D^* must satisfy generalized ratio/difference compatibility.

Suppose that generalized ratio/difference compatibility were sufficient to imply the existence of a ratio/difference representation, and let f be the ratio/difference representation for R^* and D^* . Since D^* is inverse to D , $-f$ must be a difference representation for D . Hence $-f = \eta I + \lambda$ for $\eta \in \text{Re}^+$ and $\lambda \in \text{Re}$. Similarly, since R^* is inverse to R , $1/f$ must be a ratio representation for R . Hence, $1/f = \alpha I^\beta$ for some $\alpha, \beta \in \text{Re}^+$. But then, for every $x \in \text{Re}^+$,

$$-\eta x - \lambda = -\eta I(x) - \lambda = f(x) = 1/\alpha I(x)^\beta = 1/\alpha x^\beta.$$

Hence, $-\eta \alpha x^{\beta+1} - \lambda \alpha x^\beta = 1$ for all $x \in \text{Re}^+$. Obviously, this identity cannot be satisfied within the constraints on η , λ , α , and β . Therefore, there cannot exist a ratio/difference representation f for R^* and D^* , so generalized ratio/difference

compatibility is not sufficient to imply the existence of a ratio/difference representation.

It should now be plausible that a more general statement of the counterexample also holds. Namely, if \geq_r and \geq_d are any relations satisfying generalized ratio/difference compatibility, then their respective inverses \geq_{r^*} and \geq_{d^*} also satisfy generalized ratio/difference compatibility. If generalized ratio/difference compatibility were sufficient to imply the existence of a ratio/difference representation, then both pairs of relations, \geq_r and \geq_d , and \geq_{r^*} and \geq_{d^*} , would possess ratio/difference representations. The essence of the counterexample is to show that this cannot be. At most one pair of relations, \geq_r and \geq_d , or the inverses \geq_{r^*} and \geq_{d^*} , could have a ratio/difference representation. Theorem 1 asserts that *exactly one pair of relations*, \geq_r and \geq_d , or \geq_{r^*} and \geq_{d^*} , has a ratio/difference representation if either pair satisfies generalized ratio/difference compatibility.

THEOREM 1. *Let \geq_r and \geq_d be quaternary relation on an arbitrary set A , and let \geq_{r^*} and \geq_{d^*} be the respective inverse relations. If \geq_r and \geq_d satisfy generalized ratio/difference compatibility, then either there exists a ratio/difference representation f for \geq_r and \geq_d , or there exists a ratio/difference representation g for \geq_{r^*} and \geq_{d^*} , but not both. Moreover, if f or g exists, then it is a ratio scale.*

The proof of Theorem 1 is based on Lemma 1 stating the solution to a functional equation.

LEMMA 1. *Let I_1 be a nonempty interval of real numbers and let I_2 be a nonempty interval of positive real numbers. If $H: I_2 \rightarrow I_1$ is a continuous, strictly increasing function satisfying*

$$H^{-1} \left[\frac{H(tx) + H(ty)}{2} \right] = t \cdot H^{-1} \left[\frac{H(x) + H(y)}{2} \right] \quad (1)$$

for all $x, y \in I_2$ and for any $t \in \text{Re}^+$ such that $tx, ty \in I_2$, then either (i) or (ii) holds:

- (i) $H(x) = \alpha + \eta \log x$ for all $x \in I_2$, where $\alpha \in \text{Re}$, $\eta \in \text{Re}^+$,
- (ii) $H(x) = \delta \alpha x^{\delta \theta} - \beta$ for all $x \in I_2$, where $\alpha, \theta \in \text{Re}^+$, $\beta \in \text{Re}$ and $\delta = +1$ or -1 .

Aczel (1966, Sect. 3.1.3) proves Lemma 1 for the case where $I_2 = \text{Re}^+$. Krantz *et al.* (1971, Sect. 4.5.3) develop the present generalization of the lemma, except that their proof appears to rule out the possibility in solution (ii) that $\delta = -1$. That this is an oversight can be established by checking that $H(x) = -x^{-1}$ is a function satisfying (1). The proof of Lemma 1 will be given in the Appendix because it is rather lengthy. Readers who wish to skip the proof may nevertheless want to note that the error in the proof of Krantz *et al.* (1971) occurs on p. 163 where it is inferred that a function called γ_1 is strictly increasing. As shown in the Appendix, γ_1 can either be strictly decreasing or strictly increasing.

The proof of Theorem 1 presented here is very close to the proof of Theorem 4.3 in Krantz *et al.* (1971, pp. 158–163), except for the modifications resulting from the reformulation of Lemma 1. To abbreviate the many references to Krantz *et al.* (1971), this work will be referred to as Fnd in the remainder of this article.

Proof of Theorem 1. Let ϕ_1 be a difference representation for $(A \times A, \geq_d)$ and let ϕ_2 be a ratio representation for $(A \times A, \geq_r)$. Notice for future reference that $-\phi_1$ must be a difference representation for $(A \times A, \geq_d)$ and $1/\phi_2$ must be a ratio representation for $(A \times A, \geq_r)$. Let $R_i = \phi_i(A)$, $i = 1, 2$. Define $h: R_2 \rightarrow R_1$ by $h[\phi_2(x)] = \phi_1(x)$ for any $x \in A$. The function h is well defined because condition (i) of Definition 2 implies that $\phi_2(x) = \phi_2(y)$ iff $xy \sim_r xx$ iff $xy \sim_d xx$ iff $\phi_1(x) = \phi_1(y)$.

It can be prove tht there exist intervals $I_1 \subseteq \text{Re}$ and $I_2 \subseteq \text{Re}^+$ and a strictly increasing, continuous function $H: I_2 \rightarrow I_1$ such that R_i is a dense subset of I_i ($i = 1, 2$), $H[\phi_2(x)] = h[\phi_2(x)]$ for all $x \in A$, and

$$H^{-1} \left[\frac{H(tu) + H(tv)}{2} \right] = t \cdot H^{-1} \left[\frac{H(u) + H(v)}{2} \right]$$

for all $u, v \in I_2$ and $t \in \text{Re}^+$ for which $tu, tv \in I_2$ (see Fnd, pp. 159–160, for proofs of these properties). Lemma 1 states the possible solutions of this functional equation.

Solution (i) of Lemma 1 implies that $\phi_1(x) = H[\phi_2(x)] = \alpha + \eta \cdot \log \phi_2(x)$ for all $x \in A$. Hence ϕ_1 is a difference representation for $(A \times A, \geq_r)$, because ϕ_2 is a ratio representation for that structure. Since ϕ_1 is also a difference representation for $(A \times A, \geq_d)$, the orderings \geq_r and \geq_d must be identical, contrary to assumption. Therefore, solution (i) must be excluded.

Solution (ii) of Lemma 1 implies that $\phi_1(x) = H[\phi_2(x)] = \delta\alpha\phi_2(x)^{\delta\theta} - \beta$ for every $x \in A$. If $\delta = +1$, then $\phi_1(x) + \beta = \alpha\phi_2(x)^\theta$ for all $x \in A$. Therefore, $f = \phi_1 + \beta = \alpha\phi_2^\theta$ is the desired ratio/difference representation for \geq_r and \geq_d . If f' is any other ratio/difference representation for \geq_r and \geq_d , then $f' = \gamma f^\lambda$ for $\gamma, \lambda \in \text{Re}^+$ because f' and f are both ratio representations for \geq_r , and $f' = \pi f + \omega$ for $\pi \in \text{Re}^+$ and $\omega \in \text{Re}$ because f' and f are both difference representations for \geq_d . Hence $\gamma f^\lambda - \pi f - \omega \equiv 0$ (where \equiv signifies “is constantly equal to”), so $\lambda = 1$, $\gamma = \pi$, and $\omega = 0$. Thus, $f' = \pi f$ so f is a ratio scale. If $\delta = -1$, then $-\phi_1(x) - \beta = \alpha[1/\phi_2(x)]^\theta$ for all $x \in A$. Since $-\phi_1$ is a difference representation for \geq_d , and $1/\phi_2$ is a ratio representation for \geq_r , we have $g = -\phi_1 - \beta = \alpha[1/\phi_2]^\theta$ as the ratio/difference representation for \geq_r and \geq_d . The proof that g is a ratio scale is the same as the proof for f .

There cannot exist both a ratio/difference representation f for \geq_r and \geq_d and a ratio/difference representation g for \geq_r and \geq_d , for suppose they both existed. Then $1/g$ would be a ratio representation for \geq_r and $-g$ would be a difference representation for \geq_d . By the uniqueness of these representations, $f = -\alpha g - \beta = \eta(1/g)^\lambda$ for $\alpha, \eta, \lambda \in \text{Re}^+$ and $\beta \in \text{Re}$. Hence

$$-\alpha g^{\lambda+1} - \beta g^\lambda - \eta \equiv 0. \tag{2}$$

Since $g(A)$ is dense in a nonempty interval of real numbers (see Fnd, p. 159 for the proof), there are no $\alpha, \lambda \in \text{Re}^+$ for which (2) is satisfied. So f and g cannot both exist. Q.E.D.

It is clear from Theorem 1 that the ratio/difference representation can be axiomatized by supplementing generalized ratio/difference compatibility with conditions that distinguish between the case where a representation exists for \geq_r and \geq_d , and the case where a representation exists for \geq_{r^*} and \geq_{d^*} . Such conditions will now be described. Suppose that \geq_r and \geq_d satisfy generalized ratio/difference compatibility, that ϕ is a ratio representation for \geq_r and that ψ is a difference representation for \geq_d . Consider the possible relations between ϕ and ψ . There exists a ratio/difference representation f for \geq_r and \geq_d iff f is a ratio representation for \geq_r and f is a difference representation for \geq_d , i.e., iff $\alpha\phi^\beta = f = \lambda\psi + \tau$ for $\alpha, \beta, \lambda \in \text{Re}^+$ and $\tau \in \text{Re}$. Therefore, the existence of a ratio/difference representation f is equivalent to the existence of constants $\eta = \alpha/\lambda \in \text{Re}^+$ and $\mu = -\tau/\lambda \in \text{Re}$ such that

$$\psi = \eta\phi^\beta + \mu. \tag{3}$$

By the same reasoning, there exists a ratio/difference representation g for \geq_{r^*} and \geq_{d^*} iff $\alpha\phi^{-\beta} = g = -\lambda\psi + \tau$ for $\alpha, \beta, \lambda \in \text{Re}^+$ and $\tau \in \text{Re}$, and this is equivalent to

$$\psi = -\eta\phi^{-\beta} + \mu \tag{4}$$

for $\eta = \alpha/\lambda \in \text{Re}^+$ and $\mu = \tau/\lambda \in \text{Re}$. Furthermore, it can be shown that

$$\psi = \eta[\log \phi] + \mu \tag{5}$$

for $\eta \in \text{Re}^+$ and $\mu \in \text{Re}$ is equivalent to the hypothesis that, with the exception of condition (iii), all assumptions of generalized ratio/difference compatibility are satisfied, i.e., \geq_r and \geq_d are identical orderings, $(A \times A, \geq_r)$ is an algebraic difference structure and conditions (i) and (ii) of Definition 2 hold.

Fagot (1963) points out that there is a simple way to distinguish between (3), (4), and (5). To describe the appropriate diagnostic property, it will be helpful to introduce some temporary existential assumptions. Later, these assumptions will be dropped once the basic ideas have been explained. Suppose that for every $w, z \in A$, there exist a \geq_r midpoint and a \geq_d midpoint in the sense that there exist $u, v \in A$ such that $wu \sim_r uz$ and $wv \sim_d vz$. If this is the case, one can define binary operations $|_r$ and $|_d$ on A by selecting for each $w, z \in A$, elements $w|_r z$ and $w|_d z$ satisfying

$$w(w|_r z) \sim_r (w|_r z)z, \tag{6}$$

$$w(w|_d z) \sim_d (w|_d z)z. \tag{7}$$

By definition, $w|_r z$ is a \geq_r midpoint between w and z , while $w|_d z$ is a midpoint between w and z . Given a ratio representation ϕ for \geq_r and a difference representation ψ for \geq_d , the relations (6) and (7) imply that $\phi(w)/\phi(w|_r z) = \phi(w|_r z)/\phi(z)$ and $\psi(w) - \psi(w|_d z) = \psi(w|_d z) - \psi(z)$. Thus, $\phi(w|_r z) = [\phi(w)\phi(z)]^{1/2}$ and

$\psi(w|_d z) = [\psi(w) + \psi(z)]/2$. In other words, $\phi(w|_r z)$ is the geometric mean of $\phi(w)$ and $\phi(z)$, while $\psi(w|_d z)$ is the arithmetic mean of $\psi(w)$ and $\psi(z)$. Fagot (1963) points out that (3)–(5) can be distinguished by the conditions:

(a) Equation (3) holds iff $\phi(w|_d z) > \phi(w|_r z)$ for all $w, z \in A$ such that $\phi(w) \neq \phi(z)$.

(b) Equation (4) holds iff $\phi(w|_d z) < \phi(w|_r z)$ for all $w, z \in A$ such that $\phi(w) \neq \phi(z)$.

(c) Equation (5) holds iff $\phi(w|_d z) = \phi(w|_r z)$ for all $w, z \in A$.

Thus (3)–(5) are distinguished by whether for every pair $w, z \in A$ of nonequivalent stimuli, ϕ applied to the \geq_d midpoint of w and z is strictly greater than, strictly less than, or equal to the geometric mean of $\phi(w)$ and $\phi(z)$.

Since (3) implies that a ratio/difference representation exists for \geq_r and \geq_d , the equivalence in (a) shows that the ratio/difference representation can be axiomatized by augmenting generalized ratio/difference compatibility with a qualitative axiom guaranteeing that $\phi(w|_d z) > \phi(w|_r z)$ for all $w, z \in A$ such that $\phi(w) \neq \phi(z)$. The following postulate formalizes this property:

(d) For all $w, z \in A$, if $wz \not\sim_r zz$, then $(w|_d z)(w|_r z) >_r zz$.

If (d) is added to Definition 2, and if \geq_r and \geq_d midpoints exist for every pair of elements, then \geq_r and \geq_d possess a ratio/difference representation. Although this statement should be qualified in ways described below, one can say heuristically that the error in the axiomatization stated in Krantz *et al.* (1971) is that it fails to assert that \geq_d midpoints of nonequivalent stimuli must strictly exceed corresponding \geq_r midpoints.

Rather than to include (d) in an axiomatization of the ratio/difference representation, however, it is preferable to weaken (d) in ways that increase both the generality and the empirical testability of the axiomatization. Assuming that \geq_r and \geq_d satisfy generalized ratio/difference compatibility, either (3) or (4) must hold. But if there exists even one pair $w, z \in A$ such that $\phi(w) \neq \phi(z)$ and $\phi(w|_d z) > \phi(w|_r z)$, then (b) implies that (4) is false. But then, (3) must be true, so a ratio/difference representation exists for \geq_r and \geq_d . This argument suggests the postulate:

(e) There exist $w, z \in A$ such that $(w|_d z)(w|_r z) >_r zz$ and $wz \not\sim_r zz$.

If (e) is added to the assumptions of Definition 2, then \geq_r and \geq_d possess a ratio/difference representation. The advantage of (e) over (d) is that it only requires that a \geq_d midpoint exceed a \geq_r midpoint for one pair of nonequivalent stimuli, and hence, its empirical verification is simpler than that of (d).

The only shortcoming of (e) is that it requires that a \geq_r midpoint and a \geq_d midpoint exist for at least one pair $w, z \in A$ such that $wz \not\sim_d zz$. Since this existential claim might be difficult to establish in certain applications of the theory, it would be preferable to weaken it. To see the appropriate weakening, appeal must be made to the following technical point. It is proven in Fnd (p. 159) that if $(A \times A, \geq_r)$ and

$(A \times A, \geq_d)$ are algebraic difference structures, and if \geq_r and \geq_d possess a ratio/difference representation, then there exist intervals I_r and I_d of real numbers such that $\phi(A)$ is dense in I_r and $\psi(A)$ is dense in I_d .³ Therefore, even if midpoints $w|_r z$ and $w|_d z$ do not exist, there must exist elements that are arbitrarily close to the midpoint locations in the sense that for any $\varepsilon > 0$ there exist $p, q \in A$ such that $\varepsilon > |\log \phi(p) - \log[\phi(w)\phi(z)]^{1/2}|$ and $\varepsilon > |\psi(q) - [\psi(w) + \psi(z)]/2|$. It should be plausible that (e) can be reformulated using elements that are close to if not precisely at the \geq_r and \geq_d midpoints of w and z . The desired condition is stated in:

(f) There exist $w, x, y, z \in A$ such that $wx \geq_d xz$, $wy \leq_r yz$, $wz \not\leq_r zz$ and $xy >_r xx$.

Intuitively, the elements x and y of (f) may be thought of as having been chosen to satisfy $\varepsilon > [\psi(w) + \psi(z)]/2 - \psi(x) > 0$ and $\varepsilon > \log \phi(y) - \log[\phi(w)\phi(z)]^{1/2} > 0$ for some very small ε . In other words, x is close to and below the \geq_d midpoint of w and z , and y is close to and above the \geq_r midpoint of w and z .

Theorem 2 asserts that if $(A \times A, \geq_r)$ and $(A \times A, \geq_d)$ are algebraic difference structures such that \geq_r and \geq_d possess a ratio/difference representation, then (f) must be satisfied. Indeed, for any choice of $w, z \in A$ such that $wz \not\leq_r zz$, there exist $x, y \in A$ satisfying (f). Conversely, if \geq_r and \geq_d satisfy (f) and generalized ratio/difference compatibility, then a ratio/difference representation exists for \geq_r and \geq_d . Before stating the representation theorem, it will be useful to state and prove a lemma showing that (f) implies condition (iii) of Definition 2. This lemma permits the substitution of (f) for condition (iii) of Definition 2 in the axiomatization of the ratio/difference representation.

LEMMA 2. *Let $(A \times A, \geq_d)$ and $(A \times A, \geq_r)$ be algebraic difference structures satisfying conditions (i) and (ii) of Definition 2. If there exist $w, x, y, z \in A$ such that $wx \geq_d xz$, $wy \leq_r yz$, $wz \not\leq_r zz$ and $xy >_r xx$, then \geq_d and \geq_r are not identical.*

Proof. Since $(A \times A, \geq_d)$ is an algebraic difference structure, there exists a difference representation f for \geq_d . Using the difference representation, it readily follows that if $ab >_d a'b'$, then $ba <_d b'a'$, and if $ab \geq_d a'b'$ and $bc >_d b'c'$, then $ac >_d a'c'$, for any $a, b, c, a', b', c' \in A$. Let $w, x, y, z \in A$ satisfy $wx \geq_d xz$, $wy \leq_r yz$ and $xy >_r xx$. By (i), $xy >_d xx$. Now $wx \geq_d xz$ and $xy >_d xx$ imply $wy >_d wx$, and $zx \geq_d zx$ and $xy >_d xx$ imply $zy >_d zx$, i.e., $xz >_d yz$. Hence $wy >_d wx \geq_d xz >_d yz$. But then $wy >_d yz$ and $wy \leq_r yz$, so \geq_d and \geq_r are not identical. Q.E.D.

The ratio/difference representation can now be axiomatized using conditions (i) and (ii) of Definition 2 and the condition formulated in Lemma 2.

THEOREM 2. *Let \geq_r and \geq_d be quaternary relations on an arbitrary set A such that $(A \times A, \geq_r)$ and $(A \times A, \geq_d)$ are both algebraic difference structures. Then there exists a ratio/difference representation f for \geq_r and \geq_d iff the following conditions (i)–(iii) hold: for any $x, y, z, x', y', z' \in A$,*

³ See footnote 2 for the definition of "dense in an interval of real numbers."

- (i) $xy \geq_r xx$ iff $xy \geq_d xx$,
- (ii) if $xx' \sim_r yy' \sim_r zz'$, then $xy \geq_d yz$ iff $x'y' \geq_d y'z'$,
- (iii) there exist $w, x, y, z \in A$ such that $wx \geq_d xz$, $wy \leq_r yz$, $wz \not\leq_d zz$ and $xy >_r xx$.

Proof. First, suppose that f is a ratio/difference representation for \geq_r and \geq_d . It is proven in Fnd (p. 153) that the existence of f implies conditions (i) and (ii) of Theorem 2. To prove condition (iii), first note that we are assuming throughout this essay that \geq_d is nondegenerate in the sense that $ab >_d a'b'$ for some $a, b, a', b' \in A$. If $ab \sim_d bb \sim_d b'b' \sim_d a'b'$, then $ab \sim_d a'b'$, contradicting $ab >_d a'b'$. Since $bb \sim_d b'b'$, either $ab \not\leq_d bb$ or $a'b' \not\leq_d b'b'$. In either case, there exist $w, z \in A$ such that $wz \not\leq_d zz$. Choose any such w and z , and let $s, t \in \text{Re}^+$ satisfy $s = [f(w) + f(z)]/2$ and $t = [f(w)f(z)]^{1/2}$. Then $4s^2 = f(w)^2 + 2f(w)f(z) + f(z)^2$ and $t^2 = f(w)f(z)$, so $4s^2 - 4t^2 = [f(w) - f(z)]^2 > 0$. But then $s - t = (s^2 - t^2)/(s + t) > 0$, so $s > t$. Let $\varepsilon = s - t$.

We know from the proof of Theorem 1 that $f(A)$ is a dense subset of an interval of positive real numbers (see Fnd, p. 159). Therefore there exist $x, y \in A$ such that $\varepsilon/2 > s - f(x) \geq 0$ and $\varepsilon/2 > f(y) - t \geq 0$. Hence $\varepsilon > s - f(x) + f(y) - t$. Since $s - t = \varepsilon$, $0 > f(y) - f(x)$, i.e., $f(x)/f(y) > 1$. Hence $xy >_r xx$. From the choice of s and x , $2f(x) \leq f(w) + f(z)$. Thus, $f(w) - f(x) \geq f(x) - f(z)$, i.e., $wx \geq_d xz$. From the choice of t and y , $f(y)^2 \geq f(w)f(z)$, so $f(w)/f(y) \leq f(y)/f(z)$, i.e., $wy \leq_r yz$. Hence we have proven the existence of $w, x, y, z \in A$ satisfying $wx \geq_d xz$, $wy \leq_r yz$, $wz \not\leq_d zz$ and $xy >_r xx$.

Conversely, suppose conditions (i)–(iii) of Theorem 2 hold. By Lemma 2, condition (iii) of Definition 2 follows from condition (iii) of Theorem 2. Since condition (i) and (ii) of Definition 2 and Theorem 2 are identical, generalized ratio/difference compatibility is satisfied by \geq_r and \geq_d . By Theorem 1, either there exists a ratio/difference representation f for \geq_r and \geq_d , or there exists a ratio/difference representation g for \geq_{r^*} and \geq_{d^*} , but not both. Choose $w, x, y, z \in A$ satisfying (iii) of Theorem 2. If g exists, then $g(w) - g(x) \leq g(x) - g(z)$, and $g(w)/g(y) \geq g(y)/g(z)$. Hence $2g(x) \geq g(w) + g(z)$ and $g(w)g(z) \geq g(y)^2$. Since $g(w) \neq g(z)$, we have

$$4g(x)^2 - 4g(y)^2 \geq g(w)^2 + 2g(w)g(z) + g(z)^2 - 4g(w)g(z) \geq [g(w) - g(z)]^2 > 0.$$

Therefore $[g(x) + g(y)][g(x) - g(y)] > 0$. Since $g(x) + g(y) > 0$, we have $g(x) - g(y) > 0$, i.e., $g(x)/g(y) > 1$. Hence $xy >_{r^*} xx$, or equivalently, $xy <_r xx$. But this contradicts $xy >_r xx$. Therefore g does not exist, so f exists. Q.E.D.

Since under the hypotheses of Theorem 2, conditions (i)–(iii) of that theorem are necessary and sufficient for the existence of a ratio/difference representation, they may be regarded as defining a property called “ratio/difference compatibility.”

DEFINITION 3. Let \geq_r and \geq_d be quaternary relations on an arbitrary set A .

Then, we say that \succcurlyeq_r and \succcurlyeq_d satisfy ratio/difference compatibility iff $(A \times A, \succcurlyeq_r)$ and $(A \times A, \succcurlyeq_d)$ are algebraic difference structures and conditions (i)–(iii) of Theorem 2 are satisfied.

The terminology makes sense because ratio/difference compatibility is a special case of generalized ratio/difference compatibility and Theorem 2 establishes that ratio/difference compatible relations possess a ratio/difference representation.

The following corollary points out that if there exists a pair of nonequivalent elements for which \succcurlyeq_r and \succcurlyeq_d midpoints exist, then there is a simple test for whether a ratio/difference representation exists for \succcurlyeq_r and \succcurlyeq_d , or for \succcurlyeq_{r^*} and \succcurlyeq_{d^*} .

COROLLARY 1. *Let \succcurlyeq_r and \succcurlyeq_d be quaternary relations on an arbitrary set A such that \succcurlyeq_r and \succcurlyeq_d satisfy generalized ratio/difference compatibility. Let \succcurlyeq_{r^*} and \succcurlyeq_{d^*} be the respective inverse relations to \succcurlyeq_r and \succcurlyeq_d . Suppose that $w, x, y, z \in A$ are any elements satisfying $wx \sim_d xz$, $wy \sim_r yz$ and $wz \not\sim_d zz$. Then, $xy \not\sim_r xx$ and*

- (i) $xy >_r xx$ iff there exists a ratio/difference representation f for \succcurlyeq_r and \succcurlyeq_d ,
- (ii) $xy <_r xx$ iff there exists a ratio/difference representation g for \succcurlyeq_{r^*} and \succcurlyeq_{d^*} .

Proof. Let $w, x, y, z \in A$ satisfy $wx \sim_d xz$, $wy \sim_r yz$ and $wz \not\sim_d zz$. Note that $wx \sim_{d^*} xz$, $wy \sim_{r^*} yz$ and $wz \not\sim_{d^*} zz$, by definition of \succcurlyeq_{r^*} and \succcurlyeq_{d^*} . By Theorem 1, either there exists a ratio/difference representation for \succcurlyeq_r and \succcurlyeq_d , or there exists a ratio/difference representation for \succcurlyeq_{r^*} and \succcurlyeq_{d^*} , but not both. If f exists, then $f(w) - f(x) = f(x) - f(z)$, $f(w)/f(y) = f(y)/f(z)$ and $f(w) - f(z) \neq 0$. Hence $2f(x) = f(w) + f(z)$ and $f(y)^2 = f(w)f(z)$, so

$$4f(x)^2 - 4f(y)^2 = [f(w) - f(z)]^2 > 0. \tag{8}$$

Since $f(x) + f(y) > 0$, the left side of (8) can be factored to yield $f(x) - f(y) > 0$. Thus $f(x)/f(y) > 0$, so $xy >_r xx$. Conversely, if $xy >_r xx$, then w, x, y, z satisfy (iii) of Theorem 2. Since \succcurlyeq_r and \succcurlyeq_d satisfy generalized ratio/difference compatibility, the remaining assumptions of Theorem 2 are satisfied. Therefore there exists a ratio/difference representation f for \succcurlyeq_r and \succcurlyeq_d by Theorem 2. A completely analogous argument shows that g exists iff $xy >_{r^*} xx$, i.e., iff $xy <_r xx$. Since f or g exists, we have $xy \not\sim_r xx$ in either case. Q.E.D.

According to Corollary 1, if \succcurlyeq_r and \succcurlyeq_d satisfy generalized ratio/difference compatibility and if it is possible to find a \succcurlyeq_d midpoint x and a \succcurlyeq_r midpoint y for some pair of nonequivalent stimuli, then it is easy to establish whether \succcurlyeq_r and \succcurlyeq_d , or \succcurlyeq_{r^*} and \succcurlyeq_{d^*} possess a ratio/difference representation. If x is strictly greater than y , a ratio/difference difference representation exists for \succcurlyeq_r and \succcurlyeq_d . If x is strictly less than y , a ratio/difference representation exists for \succcurlyeq_{r^*} and \succcurlyeq_{d^*} . Furthermore, it must be the case that either x is strictly greater than y or x is strictly less than y . Earlier, the heuristic remark was made that the axiomatization of the ratio/difference representation in Fnd falls short of being sufficient because it omits the requirement that the

\geq_d midpoints of nonequivalent elements must be strictly greater than the corresponding \geq_r midpoints. This remark is valid in any case where it can be established that there exists at least one pair of nonequivalent elements for which a \geq_r midpoint and a \geq_d midpoint both exist. If no such pair exists, the weaker condition (iii) of Theorem 2 will do in its stead, although its verbal formulation is not as simple to state.

APPENDIX

The proof of Lemma 1 given here parallels the logic of the proof given in Fnd (pp. 160–163), with the exception that a correction is substituted at the point where that proof goes wrong. The present proof also differs from that of Fnd in that certain constructions are explicitly formalized here, whereas they are only informally sketched in Fnd. To facilitate comparisons between the present proof and that of Fnd, equations will be numbered in the manner (k/n) to indicate that the equation is the k th equation of the present essay and the n th equation of the relevant sections of Fnd (pp. 152–154; 158–163). For example, Eq. (1) will henceforth be referred to as Eq. (1/5) since it is the first equation of this article but Eq. (5) in the indicated sections of Fnd. Equations only appearing in the present article will be numbered in the usual manner, consecutively with the other equations.

Proof of Lemma 1. It is routine to check that functions of the forms (i) and (ii) are continuous, strictly increasing, and satisfy (1/5). We must show that these are the only functions that satisfy (1/5).

For $t \in \text{Re}^+$, define $I_2(t)$ and $I_1(t)$ by

$$I_2(t) = \{x \in I_2 : tx \in I_2\},$$

$$I_1(t) = \{H(x) \in I_1 : x \in I_2(t)\}.$$

If there exist a greatest element x^* and a least element x_* in I_2 , let $Z = \{x^*/x_*, x_*/x^*\}$. If x^* or x_* does not exist, let $Z = \emptyset$, where \emptyset denotes the empty set. Define $T = \{s \in \text{Re}^+ : I_2(s) \neq \emptyset\} - Z$. It is routine to show that T is an interval of real numbers, $s \in T$ iff $s^{-1} \in T$, and for any $s \in T$, $I_2(s)$ and $I_1(s)$ are intervals of real numbers. (The reason for the fussy definition of T is that if $s = x^*/x_*$, then $I_2(s) = \{x_*\}$ is not an interval. So x^*/x_* and x_*/x^* must be excluded from T if we are to have that $I_2(s)$ is an interval for all $s \in T$.)

By a neighborhood of unity, we simply mean an interval containing 1. The proof depends on constructing a neighborhood of unity having certain desirable properties (namely, equations (9), (11/11), and (12/12)). This neighborhood of unity (denoted U) can be constructed as follows. Since I_2 is a positive interval, we can choose $a, b, c, d \in I_2$ such that $a > b > c > d$ and $a/b = c/d$. Let $\mu = \sqrt{ab}$ and $\pi = \sqrt{cd}$. Since $a > \mu > \pi > d$, we must have $\mu, \pi \in I_2$. Choose $\xi \in \text{Re}^+$ such that $\xi^2 = a/b = c/d$.

Then $\xi\mu = a$, $\xi^{-1}\mu = b$, $\xi\pi = c$ and $\xi^{-1}\pi = d$. Hence $\mu, \pi \in I_2(\xi) \cap I_2(\xi^{-1})$. Moreover, if $\xi^{-1} < \alpha < \xi$ and $x \in I_2(\xi) \cap I_2(\xi^{-1})$, then $\xi^{-1}x < \alpha x < \xi x$ so $x \in I_2(\alpha)$. Therefore $I_2(\xi) \cap I_2(\xi^{-1}) \subseteq I_2(\alpha)$ for every α such that $\xi^{-1} < \alpha < \xi$. Define $U = \{t \in T: \xi^{-1} < t^2 < \xi\}$. Note that if $s, t \in U$, then $\xi^{-1} < st < \xi$. Hence, $st \in T$ and $I_2(\xi) \cap I_2(\xi^{-1}) \subseteq I_2(st)$. Also, if $t \in U$, then $\xi^{-1} < \xi^{-1/2} < t < \xi^{1/2} < \xi$ so $I_2(\xi) \cap I_2(\xi^{-1}) \subseteq I_2(t)$. We have established that there exist $\mu, \pi \in I_2$ such that $\mu \neq \pi$ and

$$\mu, \pi \in I_2(\xi) \cap I_2(\xi^{-1}) \subseteq I_2(s) \cap I_2(t) \cap I_2(st) \tag{9}$$

for every $s, t \in U$.

It can be shown that (1/5) implies that

$$H(tx) = \gamma_1(t)H(x) + \gamma_2(t) \tag{10/10}$$

for all $t \in T$ and $x \in I_2(t)$, where γ_2 is some real valued function and γ_1 is a positively valued function (Fnd, p. 161). Furthermore, using (9), it can be shown that γ_1 and γ_2 satisfy the relations

$$\gamma_1(st) = \gamma_1(s)\gamma_1(t), \tag{11/11}$$

$$\gamma_2(t)[\gamma_1(s) - 1] = \gamma_2(s)[\gamma_1(t) - 1] \tag{12/12}$$

for every $s, t \in U$ [Fnd, p. 162]. For the sake of completeness, the derivation of (10/10)–(12/12) given in Fnd will be repeated here.

For any $t \in T$, define $f_t: I_1(t) \rightarrow I_1$ by $f_t(u) = H[t \cdot H^{-1}(u)]$ for every $u \in I_1(t)$. Since f_t is a composition of continuous functions, it is continuous. For any $u, v \in I_1(t)$, let $x = H^{-1}(u)$ and $z = H^{-1}(v)$; by definition of $I_1(t)$, $x, z, tx, tz \in I_2$ so by (1/5) we have

$$f_t \left[\frac{u+v}{2} \right] = \frac{f_t(u) + f_t(v)}{2}. \tag{13/8}$$

Therefore f_t satisfies Jensen's equation; its only continuous solutions on an interval $I_1(t)$ have the form $f_t(u) = \gamma_1 u + \gamma_2$ (Aczel, 1966, Sect. 2.1.4). Here, γ_1 must be positive since f_t is increasing. Noting that γ_1 and γ_2 may depend on t and that $u = H(x)$ for some $x \in I_2(t)$, we have $f_t[H(x)] = \gamma_1(t)H(x) + \gamma_2(t)$. Applying the definition of f_t , we have (10/10). Now choose any $s, t \in U$. We know that $t\mu, t\pi \in I_2(s)$ and $s\mu, s\pi \in I_2(t)$ for the μ, π satisfying (9) because $\mu, \pi \in I_2(s) \cap I_2(t) \cap I_2(st)$. From (10/10) we have

$$\begin{aligned} \gamma_1(st)[H(\mu) - H(\pi)] &= H(st\mu) - H(st\pi) \\ &= \gamma_1(s)[H(t\mu) - H(t\pi)] \\ &= \gamma_1(s)\gamma_1(t)[H(\mu) - H(\pi)]. \end{aligned}$$

Since H is strictly increasing, $H(\mu) - H(\pi) \neq 0$. Therefore (11/11) holds. Furthermore, repeated application of (10/10) yields

$$\begin{aligned} H(st\mu) &= \gamma_1(s)[\gamma_1(t)H(\mu) + \gamma_2(t)] + \gamma_2(s) \\ &= \gamma_1(t)[\gamma_1(s)H(\mu) + \gamma_2(s)] + \gamma_2(t). \end{aligned}$$

Multiplying out this last equation and rearranging terms yields (12/12).

The proof now splits into two cases. First, suppose $\gamma_1 \equiv 1$ in U . Then Eq. (10/10) implies that

$$H(tx) = H(x) + \gamma_2(t) \quad (14/13)$$

for any $t \in U$, $x \in I_2(t)$. For arbitrarily chosen $s, t \in U$, we may not have $st \in U$. Nevertheless for the $\mu, \pi \in I_2$ satisfying (9), Eqs. (14/13) and (10/10) imply

$$\begin{aligned} H(\mu) + \gamma_2(t) + \gamma_2(s) &= H(t\mu) + \gamma_2(s) \\ &= H(st\mu) \\ &= \gamma_1(st)H(\mu) + \gamma_2(st), \end{aligned}$$

and similarly,

$$H(\pi) + \gamma_2(t) + \gamma_2(s) = \gamma_1(st)H(\pi) + \gamma_2(st). \quad (15)$$

Hence, $H(\mu) - H(\pi) = \gamma_1(st)[H(\mu) - H(\pi)]$. Since $H(\mu) \neq H(\pi)$, $\gamma_1(st) = 1$. Combining this with (15) yields

$$\gamma_2(st) = \gamma_2(s) + \gamma_2(t) \quad (16/14)$$

for any $s, t \in U$.

Second, suppose $\gamma_1 \not\equiv 1$ in U . It is claimed in Fnd (p. 162) that in this case, $\gamma_1(t) = 1$ iff $\gamma_2(t) = 0$, but this statement is too strong. If $\gamma_1(t) = 1$, then (12/12) implies that $\gamma_2(t)[\gamma_1(s) - 1] = 0$. Since $s \in U$ may be chosen such that $\gamma_1(s) \neq 1$, it follows that $\gamma_2(t) = 0$. Therefore, $\gamma_1(t) = 1$ implies that $\gamma_2(t) = 0$. The converse, however, is not true. Even if $\gamma_2(t) = 0$, if $\gamma_2(s) = 0$ for all $s \in U$, then $\gamma_1(t)$ need not equal 1. For example, if H is the identity function, then H satisfies (1/5), and $H(tx) = \gamma_1(t)H(x) + \gamma_2(t)$, where γ_1 is the identity function and $\gamma_2 \equiv 0$. But then for $t \neq 1$, $\gamma_2(t) = 0$ but $\gamma_1(t) \neq 1$. This error does not lead to invalid inferences in the remainder of the proof presented in Fnd.

If $\gamma_1 \not\equiv 1$, choose $s, t \in U$ such that $\gamma_1(s), \gamma_1(t) \neq 1$. Separating variables in (12/12) yields

$$\gamma_2(t)/[\gamma_1(t) - 1] = \gamma_2(s)/[\gamma_1(s) - 1] = \beta,$$

where $\beta \neq 0$ iff $\gamma_2(t) \neq 0$ for some $t \in U$ such that $\gamma_1(t) \neq 1$. Thus, $\gamma_2(t) = \beta[\gamma_1(t) - 1]$ for all $t \in U$ (including those t for which $\gamma_1(t) = 1$). Substituting this in (10/10) yields

$$H(tx) + \beta = \gamma_1(t)[H(x) + \beta] \tag{17/15}$$

for all $t \in U$ and $x \in I_2(t)$.

It is important to examine the sign of $H(x) + \beta$. First suppose that $H(x) + \beta = 0$ for some $x \in I_2$. Since I_2 is an interval, there exists $y \in I_2$ such that $y < x$ or $x < y$. If $y < x$, choose $t \in U$ such that $y < tx < x$. Then $tx \in I_2$, so (17/15) implies that $H(tx) + \beta = \gamma_1(t)[H(x) + \beta] = 0 = H(x) + \beta$, contradicting the assumption that H is strictly monotone increasing. Similarly, $x < y$ leads to a contradiction of the same assumption. Hence, $H(x) + \beta \neq 0$ for every $x \in I_2$. But now, if $H(x) + \beta < 0$ and $H(y) + \beta > 0$ for some $x, y \in I_2$, then there exists $z \in I_2$ such that $H(z) + \beta = 0$ because H is continuous and I_2 is an interval. Since this is impossible, $H(x) + \beta > 0$ for all $x \in I_2$ or $H(x) + \beta < 0$ for all $x \in I_2$.

To see that there exists H satisfying (1/5) for which $H(x) + \beta$ is always negative, let $H(x) = -x^{-1}$ for $x \in \text{Re}^+$. Then H is a continuous, strictly increasing function satisfying (1/5), $\beta = 0$ and $H(x) + \beta < 0$ for all $x \in \text{Re}^+$. On the other hand, if H is the identity function on Re^+ , H satisfies (1/5), $\beta = 0$ and $H(x) + \beta > 0$ for all $x \in \text{Re}^+$.

Note that Eqs. (11/11) and (16/14) are satisfied by all $s, t \in U$. As pointed out in Fnd, these equations are variants of Cauchy's equation whose respective solutions in any neighborhood of unity are

$$\gamma_1(t) = t^\theta, \tag{18/16}$$

$$\gamma_2(t) = \eta \log t \tag{19/17}$$

for some real θ and η (see, also, Aczel, 1966, Sect. 2.1.4). If $\gamma_1 \equiv 1$ in U and thus (16/14) holds, then γ_2 must be strictly increasing because H is strictly increasing and (14/13) holds. Thus, η in (19/17) must be positive.

The critical error in the derivation in Fnd occurs at this point. It is asserted in Fnd that if $\gamma_1 \not\equiv 1$ in U , then γ_1 is strictly increasing because H is strictly increasing and (17/15) holds. But this inference is valid only if $H(x) + \beta$ is necessarily positive, and as previously noted, this condition is not satisfied. Since $H(x) + \beta$ can either be always positive or always negative, γ_1 can either be strictly increasing or strictly decreasing. Hence, if $\gamma_1 \not\equiv 1$, the parameter θ of (18/16) can either be strictly positive or strictly negative depending on the sign of $H(x) + \beta$ (for any $x \in I_2$).

The remainder of the proof is essentially the same as the proof given in Fnd. Suppose $\gamma_1 \not\equiv 1$ in U . Then from (18/16) and (17/15) we have

$$H(tx) + \beta = t^\theta [H(x) + \beta] \tag{20/19}$$

for all $t \in U$ and $x \in I_2(t)$. Choose an arbitrary x_0 in the interior of I_2 . For any $x \in I_2$ such that $x/x_0 \in U$, (20/19) yields

$$H(x) + \beta = H[(x/x_0)x_0] + \beta = (x/x_0)^\theta [H(x_0) + \beta] = ax^\theta \tag{21}$$

where $\alpha = x_0^{-\theta}[H(x_0) + \beta]$. To show that (21) actually holds for all $x \in I_2$, let $V = \{x \in I_2 : x > x_0 \text{ and } H(x) + \beta \neq \alpha x^\theta\}$. If V is nonempty, let v be the greatest lower bound of V . Choose $y \in V$ and $x \in I_2 - V$ such that $y > v \geq x \geq x_0$ and $y/x \in U$. Since $H(x) + \beta = \alpha x^\theta$, (20/19) implies that

$$H(y) + \beta = (y/x)^\theta [H(x) + \beta] = \alpha y^\theta$$

contradicting the choice of y . Hence $V = \emptyset$. Similarly, let $W = \{x \in I_2 : x < x_0 \text{ and } H(x) + \beta \neq \alpha x^\theta\}$. If W is nonempty, let w be the least upper bound of W . Choose $y \in W$ and $x \in I_2 - W$ such that $y < w \leq x \leq x_0$ and $y/x \in U$. Repeating the previous argument shows that $H(y) + \beta = \alpha y^\theta$, contradicting the choice of y . Hence $W = \emptyset$. Therefore (21) holds for all $x \in I_2$. But this shows that $H(x) = \alpha x^\theta - \beta$ for all $x \in I_2$. Since H is increasing, the definition of α and (21) imply that $\alpha > 0$ iff $H(x_0) + \beta > 0$ iff $\theta > 0$. Let $\delta = +1$ or -1 depending on whether $H(x_0) + \beta > 0$ or < 0 . Then we can stipulate that α and θ be positive, and $H(x) = \delta \alpha x^{\delta\theta} - \beta$.

If $\gamma_1 \equiv 1$ in U , then from (14/23) and (19/17) we have

$$H(tx) = H(x) + \eta \log t$$

for $t \in U$, $x \in I_2(t)$. Again choosing an arbitrary x_0 in the interior of I_2 and any $x \in I_2$ such that $x/x_0 \in U$,

$$\begin{aligned} H(x) &= H[(x/x_0) x_0] = H(x_0) + \eta[\log x/x_0] \\ &= \eta \log x + \alpha \end{aligned} \tag{22}$$

where $\alpha = H(x_0) - \eta \log x_0$. But the same argument used to show that (21) holds for all $x \in I_2$ when $\gamma_1 \neq 1$ in U proves that (22) holds for all $x \in I_2$ when $\gamma_1 \equiv 1$ in U .

Q.E.D.

REFERENCES

- ACZEL, J. (1966). *Lectures on functional equations and their applications*. New York: Academic Press.
- BIRNBAUM, M. H. (1978). Differences and ratios in psychological measurement. In N. J. Castellan, Jr. & F. Restle (Eds.), *Cognitive theory* (Vol. 3). Hillsdale, N. J.: Erlbaum.
- BIRNBAUM, M. H. (1980). Comparison of two theories of "ratio" and "difference" judgments. *Journal of Experimental Psychology: General*, **109**, 304-319.
- BIRNBAUM, M. H. (1982). Controversies in psychological measurement. In B. Wegener (Ed.), *Social attitudes and psychophysical measurement* (pp. 401-485). Hillsdale, N. J.: Erlbaum.
- BIRNBAUM, M. H., & ELMASIAN, R. (1978). Loudness "ratios" and "differences" involve the same psychophysical operation. *Perception & Psychophysics*, **23**, 403-408.
- BIRNBAUM, M. H., & MELLERS, B. A. (1978). Measurement and the mental map. *Perception & Psychophysics*, **23** (5), 403-408.
- BIRNBAUM, M. H., & VEIT, C.T. (1974). Scale convergence as a criterion for rescaling: Information integration with difference, ratio and averaging tasks. *Perception & Psychophysics*, **15**, 7-15.
- FAGOT, R. F. (1963). On the psychophysical law and estimation procedures in psychophysical scaling. *Psychometrika*, **28**, 145-160.

- GARNER, W. R. (1954). A technique and a scale for loudness measurement. *Journal of the Acoustical Society of America*, **26** (1), 73-88.
- HAGERTY, M., & BIRNBAUM, M. H. (1978). Nonmetric tests of ratio vs subtractive theories of stimulus comparison. *Perception & Psychophysics*, **24**, 121-129.
- KRANTZ, D. H., LUCE, R. D., SUPPES, P., & TVERSKY, A. (1971). *Foundations of measurement* (Vol. 1) New York: Academic Press.
- MARKS, L. E. (1974). On scales of sensation: Prolegomena to any future psychophysics that will be able to come forth as science. *Perception & Psychophysics*, **16**, 358-376.
- MELLERS, B. A., DAVIS, D. M., & BIRNBAUM, M. H. (1984). Weight of evidence supports one operation for "ratios" and "differences" of heaviness. *Journal of Experimental Psychology: Human Perception & Performance*, **10**, 216-230.
- PARKER, S., SCHNEIDER, B., & KANOW, G. ((1975). Ratio scale measurement of the perceived length of lines. *Journal of Experimental Psychology: Human Perception and Performance*, **104**, 195-204.
- RULE, S. J., & CURTIS, D. W. (1980). Ordinal properties of subjective ratios and differences: Comment on Veit. *Journal of Experimental Psychology: General*, **109**, 296-300.
- RULE, S. J., CURTIS, D. W. & MULLIN, L. C. (1981). Subjective ratios and differences in perceived heaviness. *Journal of Experimental Psychology: Human Perception and Performance*, **7**, 459-466.
- SCHNEIDER, B., PARKER, S., FARRELL, G., & KANOW, G. (1976). The perceptual basis of loudness ratio judgments. *Perception & Psychophysics*, **19**, 309-320.
- STEVENS, S. S. (1957). On the psychophysical law. *Psychological Review*, **64**, 153-181.
- STEVENS, S. S. (1971). Issues in psychophysical measurement. *Psychological Review*, **78**, 426-450.
- TORGERSOHN, W. S. (1961). Distances and ratios in psychophysical scaling. *Acta Psychologica*, **19** (5), 201-205.
- VEIT, C. T. (1978). Ratio and subtractive processes in psychophysical judgment. *Journal of Experimental Psychology: General*, **107**, 81-107.
- VEIT, C. T. (1980). Analyzing "ratio" and "difference" judgments: A reply to Rule and Curtis. *Journal of Experimental Psychology: General*, **109**, 301-303.

RECEIVED: September 21, 1983