

## Statistics 521, Problem Set 10

Wellner; 11/28/2007

**Reading:** Shorack, PfS, Chapter 10, pages 219 - 258.

**Reminder:** Final exam, 2:30 - 4:20, Wednesday, December 12.

**Due:** Wednesday, December 5, 2007.

1. Let  $K \geq 3$  be a prime and let  $X$  and  $Y$  be independent random variables that are uniformly distributed on  $\{0, 1, \dots, K - 1\}$ . For  $0 \leq n < K$  let  $Z_n = (X + nY) \bmod(K)$ . Show that  $Z_0, \dots, Z_{K-1}$  are *pairwise independent*; that is, each pair is independent, but if we know the values of two of the variables, then we know the values of all the variables.
2. Show that if  $X_n$  is any sequence of random variables, there are constants  $c_n \rightarrow \infty$  so that  $X_n/c_n \rightarrow_{a.s.} 0$ .
3. Show that if  $P(A_n) \rightarrow 0$  and  $\sum_{n=1}^{\infty} P(A_n \cap A_{n+1}^c) < \infty$ , then  $P(A_n \text{ i.o.}) = 0$ .
4. Let  $X_1, X_2, \dots$  be independent. Show that  $\sup X_n < \infty$  almost surely if and only if  $\sum_n P(X_n > M) < \infty$  for some  $M < \infty$ .
5. Let  $X_1, X_2, \dots$  be independent with  $P(X_n = 1) = p_n$  and  $P(X_n = 0) = 1 - p_n$ . Show that: (i)  $X_n \rightarrow_p 0$  if and only if  $p_n \rightarrow 0$ , and  $X_n \rightarrow_{a.s.} 0$  if and only if  $\sum_n p_n < \infty$ .
6. Suppose that  $X_1, X_2, \dots$  are independent with  $P(X_n > x) = x^{-5}$  for all  $x \geq 1$  and  $n = 1, 2, \dots$ . Show that  $\limsup_{n \rightarrow \infty} (\log X_n) / \log n = c$  almost surely for some number  $c$ , and find  $c$ .

7. **Optional bonus problem:** Suppose  $U(\omega) = \omega$  for

$$(\Omega, \mathcal{A}, P) = ((0, 1], \mathcal{B}_{(0,1]}, \lambda)$$

where  $\lambda$  is Lebesgue measure (restricted to  $(0, 1]$ ). Thus  $U \sim \text{Uniform}(0, 1)$ . Define

$$T(\omega) = \begin{cases} 2\omega, & 0 < \omega \leq 1/2, \\ 2\omega - 1, & 1/2 < \omega \leq 1, \end{cases} \quad X_1(\omega) = \begin{cases} 0, & 0 < \omega \leq 1/2, \\ 1, & 1/2 < \omega \leq 1, \end{cases}$$

and, for  $i \geq 2$ ,

$$X_i(\omega) = X_1(T^{i-1}\omega).$$

It follows that

$$\sum_{i=1}^n \frac{X_i(\omega)}{2^i} < \omega \leq \sum_{i=1}^n \frac{X_i(\omega)}{2^i} + \frac{1}{2^n}$$

and the  $X_i$ 's give the diadic (non-terminating expansion) representation of  $U$ :

$$U(\omega) = \sum_{i=1}^{\infty} \frac{X_i(\omega)}{2^i}.$$

Show that  $X_1, X_2, \dots$  are independent random variables.

[Hint: see Billingsley, *Probability Theory and Measure*, pages 1-5 and A31, page 572.]