

Statistics 521, Problem Set 5 Solutions

Wellner; 10/31/2007

1. PfS, Exercise 3.4.2, page 48: Show that $\rho = 1$ if and only if $X - \mu_X = a(Y - \mu_Y)$ for some $a > 0$; and $\rho = -1$ if and only if $X - \mu_X = a(Y - \mu_Y)$ for some $a < 0$. Thus ρ measure linear dependence, not dependence.

Solution: The “if” direction is easy: if $X - \mu_X = a(Y - \mu_Y)$ with $a > 0$, then $Cov(X, Y) = E(X - \mu_X)(Y - \mu_Y) = aE(Y - \mu_Y)^2 = aVar(Y)$ and $Var(X) = a^2Var(Y)$, which yields $\rho = 1$. Similarly, if $X - \mu_X = a(Y - \mu_Y)$ with $a < 0$, then $\rho = -1$. Conversely, suppose that $\rho^2 = 1$. Then $|Cov(X, Y)|^2 = Var(X)Var(Y)$, and hence, by the if and only if condition for equality in the Cauchy-Schwarz inequality, $\frac{|X - \mu_X|}{\sigma_X} = \frac{|Y - \mu_Y|}{\sigma_Y}$ a.s., or equivalently

$$|X - \mu_X| = \frac{\sigma_X}{\sigma_Y}|Y - \mu_Y|. \quad (1)$$

To separate out what is going on in the two cases $\rho = 1$ and $\rho = -1$, consider first $\rho = 1$. Then we have equality throughout the system of inequalities given by

$$\begin{aligned} Cov(X, Y) &= E(X - \mu_X)(Y - \mu_Y) \\ &\leq |E[(X - \mu_X)(Y - \mu_Y)]| \\ &\leq E[|(X - \mu_X)(Y - \mu_Y)|] \\ &\leq \sqrt{Var(X)Var(Y)}. \end{aligned}$$

Equality in the third inequality implies that (1) holds. Equality in the second inequality implies that either $(X - \mu_X)(Y - \mu_Y) \geq 0$ a.s. or $(X - \mu_X)(Y - \mu_Y) \leq 0$ a.s. (to see this, write out $|EY| = E|Y|$ in terms of positive and negative parts and use problem #3, problem set 4). But equality in the first inequality implies that $E\{[(X - \mu_X)(Y - \mu_Y)]^+\} \geq E\{[(X - \mu_X)(Y - \mu_Y)]^-\}$, and when combined with the preceding, this implies that $(X - \mu_X)(Y - \mu_Y) \geq 0$ a.s. Hence we conclude in this case that $(X - \mu_X)$ and $(Y - \mu_Y)$ have the same sign, and this in combination with (1) yields

$$X - \mu_X = \frac{\sigma_X}{\sigma_Y}(Y - \mu_Y),$$

with $a = \frac{\sigma_X}{\sigma_Y} > 0$.

In the case $\rho = -1$, equality in the third inequality implies that (1) holds. But now we must have $-Cov(X, Y) = |Cov(X, Y)|$ which $E\{[(X - \mu_X)(Y - \mu_Y)]^-\} \geq E\{[(X - \mu_X)(Y - \mu_Y)]^+\}$, and when combined with the consequence of the second inequality (either $(X - \mu_X)(Y - \mu_Y) \geq 0$ a.s. or $(X - \mu_X)(Y - \mu_Y) \leq 0$ a.s.), this implies that $(X - \mu_X)(Y - \mu_Y) \leq 0$ a.s.; i.e. $(X - \mu_X)$ and $(Y - \mu_Y)$ have opposite signs. This in combination with (1) yields

$$X - \mu_X = -\frac{\sigma_X}{\sigma_Y}(Y - \mu_Y) = a(Y - \mu_Y)$$

with $a = -\frac{\sigma_X}{\sigma_Y} < 0$.

2. Let $\mu_r \equiv E|X|^r$. For $r \geq s \geq t \geq 0$ we have $\mu_r^{s-t} \mu_t^{r-s} \geq \mu_s^{r-t}$.

Solution: Note that we can rewrite the inequality as

$$\mu_s \leq \mu_r^{(s-t)/(r-t)} \mu_t^{(r-s)/(r-t)}$$

where $\alpha \equiv (r-t)/(s-t)$ and $\beta \equiv (r-t)/(r-s)$ satisfy $\alpha^{-1} + \beta^{-1} = (s-t)/(r-t) + (r-s)/(r-t) = 1$. Note that $r/\alpha + t/\beta = s$. Thus we see that Hölder's inequality with the powers α and β yields

$$\begin{aligned} \mu_s = E|X|^s &= E\{|X|^{r/\alpha} |X|^{t/\beta}\} \leq \{E|X|^r\}^{1/\alpha} \{E|X|^t\}^{1/\beta} \\ &= \mu_r^{(s-t)/(r-t)} \mu_t^{(r-s)/(r-t)}. \end{aligned}$$

3. Suppose that $\epsilon_1, \dots, \epsilon_n$ are i.i.d. random variables with $P(\epsilon_i = \pm 1) = 1/2$, and let $a_i \in R, i = 1, \dots, n$. Prove the Khintchine inequalities for the case $p = 1$: for some constants A_p and B_p

$$A_p \left(\sum_{i=1}^n a_i^2 \right)^{1/2} \leq \left(E \left| \sum_{i=1}^n a_i \epsilon_i \right|^p \right)^{1/p} \leq B_p \left(\sum_{i=1}^n a_i^2 \right)^{1/2}.$$

Solution: When $p = 1$, the upper bound is easy: by Liapunov's inequality (or by Jensen's inequality with $g(x) = x^2$), $E|X| \leq (E|X|^2)^{1/2}$. Thus

$$E\left|\sum_1^n a_i \epsilon_i\right| \leq \left(E\left|\sum_1^n a_i \epsilon_i\right|^2\right)^{1/2} = \left(\sum_1^n a_i^2\right)^{1/2},$$

so the upper bound holds with $B_1 = 1$.

For the lower bound, taking $r = 4$, $s = 2$, and $t = 1$ in the previous problem (Littlewood's inequalities) yields

$$E|X|^2 \leq \{E|X|^4\}^{1/3} \{E|X|\}^{2/3},$$

or

$$\frac{\{E|X|^2\}^{3/2}}{\{E|X|^4\}^{1/2}} \leq E|X|.$$

With $X = \sum_{i=1}^n a_i \epsilon_i$ we find that $E(X^2) = \sum_{i=1}^n a_i^2$ and

$$\begin{aligned} E|X|^4 &= E\left\{\sum_{j,j',k,k'=1}^n a_j a_{j'} a_k a_{k'} \epsilon_j \epsilon_{j'} \epsilon_k \epsilon_{k'}\right\} \\ &= \sum_{i=1}^n a_i^4 + \binom{4}{2} \sum_{j < j'} a_j^2 a_{j'}^2 \\ &= \sum_{i=1}^n a_i^4 + 6 \sum_{j < j'} a_j^2 a_{j'}^2 \\ &\leq 3 \left(\sum_{i=1}^n a_i^2\right)^2. \end{aligned}$$

Hence it follows that

$$E|X| \geq \frac{\{E|X|^2\}^{3/2}}{\{E|X|^4\}^{1/2}} \geq \frac{(\sum a_i^2)^{3/2}}{\sqrt{3} \sum a_i^2} = \frac{1}{\sqrt{3}} \left(\sum_{i=1}^n a_i^2\right)^{1/2}.$$

We conclude that Khintchine's inequality holds for $p = 1$ with $A_1 = 1/\sqrt{3}$ and $B_1 = 1$. The best possible constants A_p and B_p are known for all p ; for $p = 1$ the best possible value of A_p is $1/\sqrt{2}$, and this is due to Szarek (1976), *Studia Math.* **63**, 197-208.

4. PfS, Exercise 3.5.3, page 55:

(a) Use simple functions and the MCT to show that $EY = \int_0^\infty (1 - F(y))dy$.

(b) Use the formula $EX = \int_0^\infty (1 - F(y))dy$ to show that for $X \geq 0$ and $\lambda \geq 0$ we have

$$\int_{[X \geq \lambda]} X dP = \lambda P(X \geq \lambda) + \int_\lambda^\infty P(X \geq y) dy.$$

(c) Suppose that there is a rv $Y \in \mathcal{L}_1$ such that $P(|X_n| \leq y) \leq P(Y \leq y)$ for all $y > 0$ and all $n \geq 1$. Use the result of (b) to show that this implies that $\{X_n\}$ is uniformly integrable.

Solution: (a) Suppose $Y \geq 0$. Then the simple functions

$$Y_n \equiv \sum_{k=1}^{n2^n} \frac{k-1}{2^n} 1_{[(k-1)2^{-n} \leq Y < k2^{-n}]}$$

satisfy $Y_n \nearrow Y$. For Y_n we compute

$$\begin{aligned} EY_n &= \sum_{k=1}^{n2^n} \frac{k-1}{2^n} P((k-1)2^{-n} \leq Y < k2^{-n}) \\ &= \sum_{k=1}^{n2^n} \left(\sum_1^{k-1} \frac{1}{2^n} \right) P((k-1)2^{-n} \leq Y < k2^{-n}) \\ &= \sum_{k=1}^{n2^n} \sum_1^{n2^n} \frac{1}{2^n} 1_{[j \leq k-1]} P((k-1)2^{-n} \leq Y < k2^{-n}) \\ &= \sum_{j=1}^{n2^n} \frac{1}{2^n} \sum_1^{n2^n} 1_{[j \leq k-1]} P((k-1)2^{-n} \leq Y < k2^{-n}) \\ &= \sum_{j=1}^{n2^n} \frac{1}{2^n} P(j2^{-n} \leq Y < n) = \sum_{j=1}^{n2^n} \frac{1}{2^n} P(j2^{-n} \leq Y_n) \\ &= \int_0^\infty P(Y_n \geq y) dy \end{aligned}$$

since $P(Y_n \geq n) = 0$ and $P(Y_n \in (j-1, j)/2^n) = 0$. By the MCT, the left side of the last display satisfies $EY_n \nearrow E(Y)$. Since $Y_n \nearrow$, we also

have $P(Y_n \geq y) \nearrow P(Y \geq y)$ for each fixed y . Hence the right side of the last display satisfies

$$\int_0^\infty P(Y_n \geq y)dy \nearrow \int_0^\infty P(Y \geq y)dy$$

by the MCT again. Thus we conclude that

$$E(Y) = \int_0^\infty P(Y \geq y)dy = \int_0^\infty P(Y > y)dy = \int_0^\infty (1 - F(y))dy$$

where the second equality holds since there are at most countably many points y with $P(Y > y) \neq P(Y \geq y)$.

(b) We will apply the formula in (a) to the random variable $Y = X1_{[X \geq \lambda]}$. Note that $Y = 0$ if $X < \lambda$, and $Y = X$ if $X \geq \lambda$. Hence we find that $P(Y = 0) = P(X < \lambda)$, and thus $P(Y > y) = 1 - P(X < \lambda)$ for $0 \leq y < \lambda$ while $P(Y > y) = 1 - P(X \leq y)$ for $\lambda \leq y < \infty$. Thus it follows from the formula in (a) that

$$\begin{aligned} E\{X1_{[X \geq \lambda]}\} &= E(Y) = \int_0^\infty P(Y > y)dy \\ &= \int_0^\lambda P(X \geq \lambda)dy + \int_\lambda^\infty P(X > y)dy \\ &= \lambda P(X \geq \lambda) + \int_\lambda^\infty P(X \geq y)dy; \end{aligned}$$

in the last equality we have used the fact that the number of discontinuities of $P(X \geq y)$ is at most countable, and hence of Lebesgue measure 0, and hence the two integrals are equal since the integrands differ on a set of Lebesgue measure at most 0.

(c) From (b) and the hypothesis it follows that

$$\begin{aligned} E\{|X_n|1_{[|X_n| \geq \lambda]}\} &= \lambda P(|X_n| \geq \lambda) + \int_\lambda^\infty P(|X_n| > y)dy \\ &\leq \lambda P(Y \geq \lambda) + \int_\lambda^\infty P(Y > y)dy \\ &= E\{Y1_{[Y \geq \lambda]}\} \end{aligned}$$

by (b) again in the last step. Thus we have

$$\sup_n E\{|X_n|1_{\{|X_n|\geq\lambda\}}\} \leq E\{Y1_{[Y\geq\lambda]}\} \rightarrow 0 \quad \text{as } \lambda \rightarrow \infty;$$

i.e. $\{X_n\}$ is uniformly integrable.

5. (a) Show that if $|X_n| \leq Y$ and Y is integrable, then $\{X_n\}$ is uniformly integrable.
 (b) Let $U \sim \text{Uniform}(0, 1)$, and let $X_n \equiv (n/\log n)1_{[0,1/n]}(U)$ for $n \geq 3$. Show that $\{X_n\}$ is uniformly integrable and $\int X_n dP \rightarrow 0$ even though they are not dominated by any integrable rv Y .
 (c) Let $Z_n = n1_{[0,1/n]}(U) - n1_{[1/n,2/n]}(U)$. Show that $\{Z_n\}$ is not uniformly integrable, but that $\int Z_n dP \rightarrow 0$.

Solution: (a) Since $|X_n| \leq Y$ implies that $P(|X_n| \geq y) \leq P(Y \geq y)$, this follows from the preceding exercise.

(b) Now $X_n \geq 0$, $X_n \rightarrow_p 0 \equiv X$, and $E(X_n) = 1/\log(n) \rightarrow 0 = E(X)$ as $n \rightarrow \infty$. Thus $\{X_n\}$ is uniformly integrable by Vitali's theorem. However the smallest rv above X_n for all $n \geq 3$ is the rv $Y = \sum_{k=3}^{\infty} \frac{k}{\log k} 1_{(1/(k+1), 1/k]}(U)$, and this has expectation

$$\begin{aligned} E(Y) &= \sum_{k=3}^{\infty} \frac{k}{\log(k)} \left\{ \frac{1}{k} - \frac{1}{k+1} \right\} \\ &= \sum_{k=3}^{\infty} \frac{k}{\log(k)} \frac{1}{k(k+1)} \\ &= \sum_{k=3}^{\infty} \frac{1}{(k+1)\log(k)} = \infty. \end{aligned}$$

(c) Note that $E(Z_n) = 1 - 1 = 0 \rightarrow 0$, and $Z_n \rightarrow_p 0 \equiv Z$ since, for $\epsilon \leq 1$ we have $P(|Z_n| \geq \epsilon) = P(U \leq 2/n) = 2/n \rightarrow 0$. But

$$E|Z_n| = 2 \not\rightarrow 0 = E(Z).$$

Hence by Vitali's theorem we conclude that $\{Z_n\}$ is not uniformly integrable.