

## Statistics 522, Problem Set 8 Solutions

Wellner; March 1, 2017

1. Exercise 11.6.4, page 34, Wellner, Chapter 11 notes.

Give a direct proof of the equivalence of (i) and (iv) in Proposition 2.2. (Hint: consider the functions  $\psi_\epsilon(x) \equiv \psi(x/\epsilon)$  with  $\psi$  as given on page 34.)

**Solution:** Suppose that (iv) holds. Let  $x \in C_F$  and  $\epsilon > 0$ . Suppose that  $\psi$  is defined by

$$\psi(y) \equiv \begin{cases} \frac{\int_y^1 \exp(-1/(u(1-u))) du}{\int_0^1 \exp(-1/(u(1-u))) du}, & 0 \leq y \leq 1, \\ 1, & y \leq 0, \\ 0, & y \geq 1. \end{cases}$$

Consider the function  $f_u(y) \equiv f_u(y; x, \epsilon)$  defined by

$$f_u(y) \equiv \psi_{\epsilon, x}(y) = \begin{cases} 0, & y \geq x + \epsilon, \\ 1, & y \leq x, \\ \psi((y - x)/\epsilon), & x \leq y \leq x + \epsilon. \end{cases}$$

Then  $f_u \in C^\infty(\mathbb{R})$  and it satisfies

$$1_{(-\infty, x]}(y) \leq f_u(y) \leq 1_{(-\infty, x + \epsilon]}(y),$$

and hence it follows that

$$\begin{aligned} F_n(x) &= E1_{(-\infty, x]}(X_n) \leq E f_u(X_n) \leq E1_{(-\infty, x + \epsilon]}(X_n) = F_n(x + \epsilon), \quad \text{and} \\ F(x) &= E1_{(-\infty, x]}(X) \leq E f_u(X) \leq E1_{(-\infty, x + \epsilon]}(X) = F(x + \epsilon). \end{aligned}$$

Therefore

$$\limsup_n F_n(x) \leq \limsup_n E f_u(X_n) = E f_u(X) \leq E1_{(-\infty, x + \epsilon]}(X) = F(x + \epsilon).$$

Letting  $\epsilon \searrow 0$  and using right-continuity of  $F$ , this yields

$$\limsup_n F_n(x) \leq F(x). \tag{1}$$

Similarly the function  $f_l(y) \equiv f_l(y; x, \epsilon) \equiv \psi((y - (x - \epsilon))/\epsilon)$  satisfies  $f_l \in C^\infty(\mathbb{R})$  and

$$1_{(-\infty, x-\epsilon]}(y) \leq f_l(y) \leq 1_{(-\infty, x]}(y),$$

and hence

$$F(x - \epsilon) \leq Ef_l(X) = \lim_n Ef_l(X_n) = \liminf_n Ef_l(X_n) \leq \liminf_n F_n(x).$$

Letting  $\epsilon \searrow 0$  and using  $x \in C_F$  (here is where  $x \in C_F$  is used!), this yields

$$F(x) \leq \liminf_n F_n(x). \quad (2)$$

Combining (1) and (2) yields  $F_n \rightarrow_d F$ ; i.e. (i) holds.

For the reverse implication, note that (i) implies (ii) by the Helly-Bray Theorem 3.5.1; (i.e.  $Ef(X_n) \rightarrow Ef(X)$  for all  $f \in C_b(\mathbb{R})$ ), and since  $C^\infty(\mathbb{R}) \subset C_b(\mathbb{R})$  (iv) holds.

2. Exercise 9.2.4, PfS Course Notes, page 199. (Exercise 11.8.4, page 293, PfS 2000.)

Suppose that  $\log X \sim N(0, 1)$ .

(i) Show that the density of  $X$  is given by  $f_X(x) = x^{-1} \exp(-(\log x)^2/2)/\sqrt{2\pi}$  for  $x > 0$  (and 0 otherwise).

(ii) For each  $a \in [-1, 1]$  consider the random variable  $Y_a$  with density

$$f_a(y) = f_X(y)(1 + a \sin(2\pi \log y)) \quad \text{for } y > 0.$$

Show that  $EX^k = EY_a^k$  for all integers  $k \geq 1$  and  $a \in [-1, 1]$ .

**Solution:** (i) Since  $X \stackrel{d}{=} e^Z$  where  $Z \sim N(0, 1)$ , it is clear that  $X > 0$  with probability 1. Hence we compute, for  $x > 0$ ,

$$F_X(x) = P(X \leq x) = P(e^Z \leq x) = P(Z \leq \log x) = \Phi(\log x)$$

where  $\Phi(z) \equiv \int_{-\infty}^z \phi(y)dy$  is the standard normal distribution function and  $\phi(z) \equiv (2\pi)^{-1/2} \exp(-z^2/2)$  is the standard normal density function. Thus

$$f_X(x) = F'_X(x) = \phi(\log x) \cdot x^{-1} \quad \text{for } x > 0.$$

(ii) Now by changing variables to  $y = \log x$  we get

$$\begin{aligned}
EX^k &= \int_0^\infty x^k x^{-1} (2\pi)^{-1/2} \exp(-(\log x)^2/2) dx \\
&= \int_{-\infty}^\infty e^{ky} \phi(y) dy \quad \text{where } \phi(z) = (2\pi)^{-1/2} e^{-z^2/2} \\
&= e^{k^2/2} \int_{-\infty}^\infty \phi(z - k) dz = e^{k^2/2}.
\end{aligned}$$

On the other hand, by the same change of variables,

$$\begin{aligned}
EY_a^k &= \int_0^\infty x^k x^{-1} (2\pi)^{-1/2} \exp(-(\log x)^2/2) (1 + a \sin(2\pi \log x)) dx \\
&= \int_{-\infty}^\infty e^{ky} \phi(y) (1 + a \sin(2\pi y)) dy \\
&= e^{k^2/2} \int_{-\infty}^\infty \phi(y - k) (1 + a \sin(2\pi y)) dy \\
&= e^{k^2/2} \int_{-\infty}^\infty \phi(z) (1 + a \sin(2\pi(z + k))) dz \\
&= e^{k^2/2} \int_{-\infty}^\infty \phi(z) (1 + a \sin(2\pi z)) dz
\end{aligned}$$

by using

$$\begin{aligned}
\sin(2\pi(z + k)) &= \sin(2\pi z) \cos(2\pi k) + \cos(2\pi z) \sin(2\pi k) \\
&= \sin(2\pi z) \cdot 1 + \cos(2\pi z) \cdot 0 \\
&= \sin(2\pi z)
\end{aligned}$$

$$\begin{aligned}
&= e^{k^2} \left( 1 + a \int_{-\infty}^\infty \sin(2\pi z) \phi(z) dz \right) \\
&= e^{k^2/2} \quad \text{since } \sin \text{ is odd and } \phi \text{ is even} \\
&= e^{k^2/2}.
\end{aligned}$$

3. Exercise 11.6.7, page 34, Wellner, Chapter 11 notes: Suppose that  $X$  and  $Y$  are independent random vectors in  $\mathbb{R}^k$  and  $X$  and  $W$  are independent random vectors in  $\mathbb{R}^k$  with  $\mu \equiv E(Y) = E(W)$  and  $\Sigma = E(X - \mu)(X - \mu)^T = E(W - \mu)(W - \mu)^T$ . Then for each  $f \in C^3(\mathbb{R})$  there is a constant  $C_f$  such that

$$|Ef(X + Y) - Ef(X + W)| \leq C_f (E\|Y\|^3 + E\|W\|^3)$$

where  $\|v\| \equiv \sum_{i=1}^k |v_i|$  for  $v = (v_1, \dots, v_k) \in \mathbb{R}^k$ .

**Solution:** First the needed multivariate Taylor's theorem with remainder. (Thanks to Mathias Hudoba de Badyn for pointing out a nice source for this, namely G. Folland's course notes for Math 425: <https://www.math.washington.edu/folland/Math425/taylor2.pdf>.) Suppose that  $f : \mathbb{R}^k \rightarrow \mathbb{R}$  with  $f \in C^3(\mathbb{R}^k)$ . Then for  $x, y \in \mathbb{R}^k$

$$f(x + y) = f(x) + y^T \nabla f(x) + \frac{1}{2} y^T H(x) y + R_{x,2}(y)$$

where

$$\begin{aligned} \nabla f(x) &= \left( \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_k} \right)^T \equiv \text{the gradient of } f \text{ at } x, \\ H(x) &\equiv \nabla^2 f(x) = \left( \frac{\partial^2 f}{\partial x_i \partial x_j} \right)_{i,j=1}^k \equiv \text{the Hessian of } f \text{ at } x, \\ R_{x,2}(y) &\equiv \sum_{|\alpha|=3} \partial^\alpha f(x + \theta y) \frac{y^\alpha}{\alpha!} \text{ for some } 0 \leq \theta \leq 1, \end{aligned}$$

and where, for  $\alpha = (\alpha_1, \dots, \alpha_k) \in \mathbb{N}^k$ ,

$$\begin{aligned} |\alpha| &\equiv \sum_{j=1}^k \alpha_j, & \alpha! &= \prod_{i=1}^k \alpha_i!, \\ x^\alpha &\equiv x_1^{\alpha_1} \cdots x_k^{\alpha_k} \text{ for } x \in \mathbb{R}^k, \\ \partial^\alpha f(x) &= \partial_1^{\alpha_1} \partial_2^{\alpha_2} \cdots \partial_k^{\alpha_k} f = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \cdots \partial x_k^{\alpha_k}}. \end{aligned}$$

Thus if  $|\partial^3 f(x)| \leq M$  for all  $x \in \mathbb{R}^k$  (and all  $\alpha$  with  $|\alpha| = 3$ ), it follows from the multinomial formula

$$(x_1 + x_2 + \cdots + x_k)^m = \sum_{|\alpha|=m} \frac{m!}{\alpha!} x^\alpha$$

with  $m = 3$ , that

$$|R_{x,2}(y)| \leq \frac{M}{3!} \|y\|^3$$

where  $\|y\|_1 = \sum_{j=1}^k |y_j|$  is the  $\ell_1$  norm of  $y$ .

Now back to the problem at hand. From the Taylor expansion we find, using the independence of  $X$  and  $Y$  and then the hypothesized equality of first and second moments, that

$$\begin{aligned}
Ef(X + Y) &= Ef(X) + E[Y^T \nabla f(X)] + \frac{1}{2} E[Y^T H(X) Y] + E[R_{X,2}(Y)] \\
&= Ef(X) + E[Y^T] E \nabla f(X) + \frac{1}{2} E[YY^T] \cdot EH(X) + E[R_{X,2}(Y)] \\
&= Ef(X) + E[W^T] E \nabla f(X) + \frac{1}{2} E[WW^T] \cdot EH(X) + E[R_{X,2}(Y)] \\
&= Ef(X + W) - E[R_{X,2}(W)] + E[R_{X,2}(Y)].
\end{aligned}$$

The second equality above is based on a computation via the trace operation as follows: with  $H \equiv H(X)$  and using the fact that equal means and covariance (matrices) for  $Y$  and  $W$  holds if and only if  $Y$  and  $W$  have equal means and equal second moment matrices (so  $E(YY^T) = E(WW^T)$ ),

$$\begin{aligned}
&E[Y^T H(X) Y] \\
&= E[\text{tr}(Y^T H Y)] = E[\text{tr}(Y Y^T \cdot H)] \\
&= \text{tr}(E[YY^T] \cdot H) \\
&= \text{tr}(E[YY^T] \cdot E(H)) \quad \text{by independence of } Y \text{ and } X \\
&= E[W^T H(X) W] \quad \text{by the same argument as for } Y \text{ above.}
\end{aligned}$$

This yields

$$\begin{aligned}
|Ef(X + Y) - Ef(X + W)| &\leq E|R_{X,2}(Y)| + E|R_{X,2}(W)| \\
&\leq \frac{M}{6} \{E\|Y\|^3 + E\|W\|^3\}
\end{aligned}$$

as claimed. Note that this can be formulated in terms of the usual Euclidean norm  $|v|_2 \equiv \{\sum_{j=1}^k v_j^2\}^{1/2}$  since the  $\ell_p$  norms on  $\mathbb{R}^k$  are related by  $|v|_p \leq |v|_{p'} \leq k^{1/p' - 1/p} |v|_p$  if  $1 \leq p' \leq p$ . In particular,  $|v|_1 \leq k^{1/2} |v|_2$ .

4. Exercise 11.6.9, page 35, Wellner, Chapter 11 notes. Use the Cramér - Wold device to prove the multivariate CLT from the classical (one-dimensional) CLT.

**Solution:** Suppose that  $\underline{X}_1, \dots, \underline{X}_n$  are i.i.d with  $E(\underline{X}) = \underline{\mu}$  and  $E((\underline{X} - \underline{\mu})(\underline{X} - \underline{\mu})^T) \equiv \Sigma$ . Let  $\underline{a} \in \mathbb{R}^k$  and consider  $Y_n \equiv \underline{a}^T \sqrt{n}(\overline{\underline{X}}_n - \underline{\mu})$ . Then since

$$\underline{V}_n \equiv \sqrt{n}(\overline{\underline{X}}_n - \underline{\mu}) = n^{-1/2} \sum_{i=1}^n (\underline{X}_i - \underline{\mu}),$$

we see that

$$\begin{aligned} Y_n &= \underline{a}^T \underline{V}_n = \underline{a}^T \sqrt{n}(\overline{\underline{X}}_n - \underline{\mu}) \\ &= n^{-1/2} \sum_{i=1}^n \underline{a}^T (\underline{X}_i - \underline{\mu}) \\ &\equiv n^{-1/2} \sum_{i=1}^n Z_i = n^{1/2} \overline{Z}_n \end{aligned}$$

where  $Z_i \equiv \underline{a}^T (\underline{X}_i - \underline{\mu})$  are i.i.d. with mean 0 and variance

$$\begin{aligned} \text{Var}(Z_1) &= \text{Var}(\underline{a}^T (\underline{X}_1 - \underline{\mu})) \\ &= \underline{a}^T E\{(\underline{X}_1 - \underline{\mu})(\underline{X}_1 - \underline{\mu})^T\} \underline{a} \\ &= \underline{a}^T \Sigma \underline{a}. \end{aligned}$$

Thus the one-dimensional (Lindeberg) CLT yields

$$Y_n \rightarrow_d Y \sim N_1(0, \underline{a}^T \Sigma \underline{a}) \stackrel{d}{=} \underline{a}^T \underline{V}$$

where  $\underline{V} \sim N_k(0, \Sigma)$ . That is,  $\underline{a}^T \underline{V}_n \rightarrow_d \underline{a}^T \underline{V}$  for every  $\underline{a} \in \mathbb{R}^k$ . By the Cramér-Wold device this implies that  $\underline{V}_n \rightarrow_d \underline{V} \sim N_k(0, \Sigma)$ .