

References, 581 Notes

Chapter 4, Section 5: EM

References

- BOYLES, R. A. (1983). On the convergence of the EM algorithm. *J. Roy. Statist. Soc. Ser. B* **45** 47–50.
- DEMPSTER, A. P., LAIRD, N. M. and RUBIN, D. B. (1977). Maximum likelihood from incomplete data via the EM algorithm. *J. Roy. Statist. Soc. Ser. B* **39** 1–38. With discussion.
- FINCH, S. J., MENDELL, N. R. and THODE, H. C. (1989). Probabilistic measures of adequacy of a numerical search for a global maximum. *J. Amer. Statist. Assoc.* **84** 1020–1023.
- GROENEBOOM, P. (1996). Lectures on inverse problems. In *Lectures on probability theory and statistics (Saint-Flour, 1994)*, vol. 1648 of *Lecture Notes in Math.* Springer, Berlin, 67–164.
- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2008). The support reduction algorithm for computing non-parametric function estimates in mixture models. *Scand. J. Statist.* **35** 385–399.
- GROENEBOOM, P. and WELLNER, J. A. (1992). *Information bounds and nonparametric maximum likelihood estimation*, vol. 19 of *DMV Seminar*. Birkhäuser Verlag, Basel.
- JAMSHIDIAN, M. and JENNRICH, R. I. (1993). Conjugate gradient acceleration of the EM algorithm. *J. Amer. Statist. Assoc.* **88** 221–228.
- JAMSHIDIAN, M. and JENNRICH, R. I. (1997). Acceleration of the EM algorithm by using quasi-Newton methods. *J. Roy. Statist. Soc. Ser. B* **59** 569–587.
- JEWELL, N. P. (1982). Mixtures of exponential distributions. *Ann. Statist.* **10** 479–484.
- KAWAKATSU, H. and LARGEY, A. G. (2009). EM algorithms for ordered probit models with endogenous regressors. *Econom. J.* **12** 164–186.

- LAIRD, N. (1978). Nonparametric maximum likelihood estimation of a mixed distribution. *J. Amer. Statist. Assoc.* **73** 805–811.
- LANGE, K. and WEEKS, D. E. (1989). Efficient computation of lod scores: genotype elimination, genotype redefinition, and hybrid maximum likelihood algorithms. *Ann. Human Genetics* **53** 67–83.
- LOUIS, T. A. (1982). Finding the observed information matrix when using the EM algorithm. *J. Roy. Statist. Soc. Ser. B* **44** 226–233.
- MASSÉ, J.-C. (1997). Nonparametric maximum likelihood estimation in a non-locally compact setting. *Metrika* **46** 123–145.
- MCLACHLAN, G. J. and KRISHNAN, T. (2008). *The EM algorithm and extensions*. 2nd ed. Wiley Series in Probability and Statistics, Wiley-Interscience [John Wiley & Sons], Hoboken, NJ.
- MEILIJSON, I. (1989). A fast improvement to the EM algorithm on its own terms. *J. Roy. Statist. Soc. Ser. B* **51** 127–138.
- MENG, X.-L. and RUBIN, D. B. (1993). Maximum likelihood estimation via the ECM algorithm: a general framework. *Biometrika* **80** 267–278.
- MENG, X.-L. and VAN DYK, D. (1998). Fast EM-type implementations for mixed effects models. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **60** 559–578.
- VAN DYK, D. A., MENG, X.-L. and RUBIN, D. B. (1995). Maximum likelihood estimation via the ECM algorithm: computing the asymptotic variance. *Statist. Sinica* **5** 55–75.
- VARADHAN, R. and ROLAND, C. (2008). Simple and globally convergent methods for accelerating the convergence of any EM algorithm. *Scand. J. Statist.* **35** 335–353.
- VARDI, Y. (1996). Network tomography: estimating source-destination traffic intensities from link data. *J. Amer. Statist. Assoc.* **91** 365–377.
- VARDI, Y. and LEE, D. (1993). From image deblurring to optimal investments: maximum likelihood solutions for positive linear inverse problems. *J. Roy. Statist. Soc. Ser. B* **55** 569–612. With discussion.
- VARDI, Y., SHEPP, L. A. and KAUFMAN, L. (1985). A statistical model for positron emission tomography. *J. Amer. Statist. Assoc.* **80** 8–37. With discussion.

- VARDI, Y. and ZHANG, C.-H. (1999). Reconstruction of binary images via the EM algorithm. In *Discrete tomography*. Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 297–316.
- WELLNER, J. A. and ZHAN, Y. (1997). A hybrid algorithm for computation of the nonparametric maximum likelihood estimator from censored data. *J. Amer. Statist. Assoc.* **92** 945–959.
- WU, C.-F. J. (1983). On the convergence properties of the EM algorithm. *Ann. Statist.* **11** 95–103.