

Statistics 582, Problem Set 8 Solutions

Wellner; 3/4/2009

1. (a) Show that the logistic distribution with location parameter θ having density

$$p_\theta(x) = \frac{\exp(x - \theta)}{(1 + \exp(x - \theta))^2} = \frac{1}{2(1 + \cosh(x - \theta))}$$

has monotone likelihood ratio.

(b) Unfortunately the result of (a) does not carry over to a sample of size n . If X_1, \dots, X_n are i.i.d. P_θ with density p_θ as in (a), then there is no $T(\underline{X})$ for which the MLR property holds. Nevertheless we can look for locally best tests. Find the locally best test of $H_0 : \theta = 0$ versus $H_1 : \theta > 0$. How would you carry out this test?

Solution: (a) Let $\theta' > \theta$. Then the ratio of densities is given by

$$\begin{aligned} \frac{p_{\theta'}(x)}{p_\theta(x)} &= \frac{e^{x-\theta'}}{(1 + e^{x-\theta'})^2} \cdot \frac{(1 + e^{x-\theta})^2}{e^{x-\theta}} \\ &= e^{\theta-\theta'} \left(\frac{1 + e^{x-\theta}}{1 + e^{x-\theta'}} \right)^2. \end{aligned}$$

This is a monotone increasing function of x if and only if its logarithm is a monotone increasing function of x . The logarithm is given by

$$\begin{aligned} \log \left(\frac{p_{\theta'}(x)}{p_\theta(x)} \right) &\equiv R(x; \theta, \theta') \equiv R(x) \\ &= \theta - \theta' + 2 \log \left(\frac{1 + e^{x-\theta}}{1 + e^{x-\theta'}} \right) \\ &= \theta - \theta' + 2 \left\{ \log(1 + e^{x-\theta}) - \log(1 + e^{x-\theta'}) \right\}, \end{aligned}$$

where R has derivative (with respect to x) R' given by

$$\begin{aligned} R'(x) &= 2 \left\{ \frac{e^{x-\theta}}{1 + e^{x-\theta}} - \frac{e^{x-\theta'}}{1 + e^{x-\theta'}} \right\} \\ &= \frac{2e^x}{(1 + e^{x-\theta})(1 + e^{x-\theta'})} \cdot (e^{-\theta} - e^{-\theta'}) \\ &> 0. \end{aligned}$$

Thus the family $\{p_\theta\}$ has monotone likelihood ratio in $T(x) = x$.

(b) As in Example 6.1.5, the locally best test is the one-sided score test, reject if $S_n(\theta_0) > k$ where

$$S_n(\theta_0) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \dot{\mathbf{i}}_\theta(X_i; \theta_0).$$

Straightforward calculation yields

$$\dot{\mathbf{i}}_\theta(x) = 2 \left\{ \frac{1}{1 + e^{-(x-\theta)}} - 1 \right\} = 2 \{F(x - \theta) - 1\}$$

where $F(x) = 1/(1 + e^{-x})$ is the standard logistic distribution function. Thus for $\theta_0 = 0$ the test statistic is

$$S_n(0) = n^{-1/2} \sum_{i=1}^n 2 \left\{ \frac{1}{1 + e^{-X_i}} - 1 \right\},$$

and we reject for large values of $S_n(0)$. Since $S_n(0) \rightarrow_d N(0, 1/3)$ under $\theta_0 = 0$, taking the constant k to be $3^{-1/2}z_\alpha$ leads to approximate size α for large n .

2. Continuation of problem 2, problem set 7:

(a) For P_0 and P_1 as given in problem 1 of problem set # 7, compute $d_{TV}(P_0, P_1)$, $H(P_0, P_1)$, and the affinity $\rho(P_0, P_1) = \int \sqrt{p_0 p_1} d\mu$.

(b) For the product laws P_{0n} and P_{1n} (corresponding to observation of X_1, \dots, X_n i.i.d. P_0 or P_1 respectively) compute $\rho(P_{0n}, P_{1n})$ and $H(P_{0n}, P_{1n})$ for $n = 20, 50, 100$.

(c) What does this imply about the test, ϕ_n say, based on X_1, \dots, X_n which minimizes the sum of risks?

Solution: First,

$$d_{TV}(P_0, P_1) = (1/2)\{.18 + .12 + .12 + .18\} = (1/2)(.6) = .3.$$

Furthermore,

$$\rho(P_0, P_1) = \sqrt{(.18)(.36)} + \sqrt{(.06)(.18)} + \sqrt{(.36)(.24)} + \sqrt{(.40)(.22)} = 0.949068$$

so that $H^2(P_0, P_1) = 1 - \rho(P_0, P_1) = .0509318\dots$, and $H(P_0, P_1) = 0.225681\dots$

Note that the inequalities of problem xxx are indeed satisfied:

$$H^2(P_0, P_1) \leq d_{TV}(P_0, P_1) \leq H(P_0, P_1)(1 + \rho(P_0, P_1))^{1/2} \leq \sqrt{2}H(P_0, P_1),$$

which in this case becomes:

$$.0509318 < .3 < 0.225681(1 + .0509318\dots)^{1/2} = .315071\dots < .319161\dots$$

For $n = 20, 50, 100$ we have

n	$\rho(P_0^n, P_1^n)$	$H(P_0^n, P_1^n)$	$H^2(P_0^n, P_1^n)$
1	0.949068	0.225681	0.0509318
20	0.351519	0.805283	0.648481
50	0.073260	0.962673	0.926739
100	0.00536713	0.997313	0.994633

Since the test $\phi = \phi(\underline{X})$ which minimizes the sum of risks has

$$\begin{aligned} E_0\phi(\underline{X}) + E_1(1 - \phi(\underline{X})) &= \int p_0(\underline{x}) \wedge p_1(\underline{x}) d\mu(\underline{x}) \\ &\leq \rho(P_0^n, P_1^n) \\ &= \rho^n(P_0, P_1) \rightarrow 0. \end{aligned}$$

From the table above we see that this happens quite rapidly.

3. Consider the Locally Most Powerful test ϕ for testing $H : \theta \leq 0 \equiv \theta_0$ versus $K : \theta > 0 = \theta_0$ in Example 6.1.5.

(a) Suggest two different approximations to the power of this test, one for local alternatives (of the form $\theta_n = t/\sqrt{n}$ with $t > 0$), and the other for fixed alternatives, $\theta > 0$.

(b) What is the behavior of each of these two approximations for large values of θ ? Which of them shows that the power function decreases to 0 as $\theta \rightarrow \infty$? Why?

Solution: (a) The test is “reject H if $\sqrt{n}\bar{Y}_n > 2^{-1/2}z_\alpha$ ” where $Y_i \equiv 2X_i/(1+X_i^2)$ are i.i.d. and $X_i \sim \text{Cauchy}(\theta, 1)$. Thus under P_θ , by using contour integration and Cauchy’s formula, or by using Mathematica, Maple, or your favorite symbolic manipulation program,

$$\begin{aligned} m(\theta) \equiv E_\theta Y_i &= \int_{-\infty}^{\infty} \frac{2x}{1+x^2} p_\theta(x) dx = \int_{-\infty}^{\infty} \frac{2x}{1+x^2} \frac{1}{\pi} \frac{1}{1+(x-\theta)^2} dx \\ &= \frac{2\theta}{4+\theta^2}, \end{aligned}$$

and

$$\begin{aligned} \sigma^2(\theta) &\equiv \text{Var}_\theta(Y_i) = E_\theta Y_i^2 - m^2(\theta) \\ &= \frac{2(4+3\theta^2)}{(4+\theta^2)^2} - \left(\frac{2\theta}{4+\theta^2} \right)^2. \end{aligned}$$

For local alternatives $\theta = \theta_n = t/\sqrt{n}$, we have

$$\begin{aligned} \text{Power}(\theta_n) &= P_{\theta_n}(\sqrt{n}\bar{Y}_n > 2^{-1/2}z_\alpha) \\ &= P_{\theta_n}(\sqrt{n}(\bar{Y}_n - m(\theta_n)) \geq 2^{-1/2}z_\alpha - \sqrt{n}(m(\theta_n) - m(0))) \\ &\rightarrow P(2^{-1/2}Z \geq 2^{-1/2}z_\alpha - m'(0)t) \end{aligned}$$

where

$$\begin{aligned} m'(0) &= \int_{-\infty}^{\infty} \frac{2x}{1+x^2} \frac{d}{d\theta} p_{\theta}(x) |_{\theta=0} dx \\ &= \int_{-\infty}^{\infty} \dot{l}_{\theta}(x; 0) \dot{l}_{\theta}(x; 0) p_{\theta}(x; 0) dx = I(\theta) = 1/2. \end{aligned}$$

Hence we have

$$\text{Power}(\theta_n) \rightarrow P(Z > z_{\alpha} - 2^{-1/2}t) = 1 - \Phi(z_{\alpha} - 2^{-1/2}t).$$

This approximation to the power function increases monotonically from α at $t = 0$ to 1 at $t = \infty$ (effectively when $t > 2^{1/2} \cdot 4$). Note that this result is very much in qualitative agreement with corollary 4.2.4 from Statistics 581.

(b) For fixed alternatives $\theta > 0$ we have

$$\begin{aligned}
 \text{Power}(\theta) &= P_{\theta}(\sqrt{n}\bar{Y} > 2^{-1/2}z_{\alpha}) \\
 &= P_{\theta}(\sqrt{n}(\bar{Y}_n - m(\theta)) > 2^{-1/2}z_{\alpha} - \sqrt{nm}(\theta)) \\
 &\doteq P(Z > (2^{-1/2}z_{\alpha} - \sqrt{nm}(\theta))/\sigma(\theta)) \\
 &= 1 - \Phi((2^{-1/2}z_{\alpha} - \sqrt{nm}(\theta))/\sigma(\theta)).
 \end{aligned}$$

This approximation to the power function is completely determined by the behavior of the functions $m(\theta)$ and $\sigma(\theta)$. Inspection of the function $(2^{-1/2}z_{\alpha} - \sqrt{nm}(\theta))/\sigma(\theta)$ shows that it first decreases (as it should if the power is to increase), but then it reaches a minimum and increases thereafter, and hence this approximation to the power decreases to zero just as we argued that it must in class. See the attached plot for $\alpha = .05$ and $n = 3, 6, 9, 12, 15$. the two different approximations for various sample sizes. Note that the two approximations agree for θ 's close to 0, but the local approximation is always monotone increasing, while the approximations with θ fixed show the approximate power decreasing to 0 as $\theta \rightarrow \infty$ as we know it must. [Also note that we still have not computed the exact (finite n) power functions; it would be interesting to know how close our approximations are to the "exact" power. I'm betting that they are reasonably good.]

4. Let X_1, \dots, X_n be a sample of size n from the uniform distribution $U(0, \theta)$. Sufficiency reduces the problem to $T = \max X_i$.
 - (a) Find the class of all Neyman-Pearson best tests of $H_0 : \theta = \theta_0$ versus $H_1 : \theta = \theta_1$, where $\theta_1 > \theta_0$.
 - (b) Find the subclass of the tests that are independent of θ_1 . These are UMP tests of H_0 versus $H_1' : \theta > \theta_0$.
 - (c) Show that the test $\phi(t) = 1\{t > \theta_0\} + \alpha 1\{t \leq \theta_0\}$ is UMP of size α for testing $H_0' : \theta \leq \theta_0$ versus $H_1' : \theta > \theta_0$ but that ϕ is not admissible.
 - (d) Show that $\phi(t) = 1\{[t > \theta_0] \cup [t \leq b]\}$ where $b = \theta_0 \alpha^{1/n}$ is a UMP test of size α for testing $H_0 : \theta = \theta_0$ versus $\theta \neq \theta_0$.

Solution: (a) For testing $\theta = \theta_0$ versus $\theta = \theta_1 > \theta_0$, the class of all NP tests is given by tests of the form

$$\phi(t) = \begin{cases} 1, & \text{if } \theta_1^{-n} 1_{[0, \theta_1]}(t) > k \theta_0^{-n} 1_{[0, \theta_0]}(t) \\ \gamma(t), & \text{if } \theta_1^{-n} 1_{[0, \theta_1]}(t) = k \theta_0^{-n} 1_{[0, \theta_0]}(t) \\ 0, & \text{if } \theta_1^{-n} 1_{[0, \theta_1]}(t) < k \theta_0^{-n} 1_{[0, \theta_0]}(t) \end{cases}.$$

Thus for $k = (\theta_0/\theta_1)^n$ the NP tests are of the form

$$\phi(t) = \begin{cases} 1, & \text{if } \theta_0 < t \leq \theta_1 \\ \gamma(t), & \text{if else;} \end{cases}$$

for $k > (\theta_0/\theta_1)^n$ the NP tests are of the form

$$\phi(t) = \begin{cases} 1, & \text{if } \theta_0 < t \leq \theta_1 \\ \gamma(t), & \text{if else} \\ 0, & \text{if } 0 \leq t \leq \theta_0; \end{cases}$$

and for $k < (\theta_0/\theta_1)^n$ the NP tests are of the form

$$\phi(t) = \begin{cases} 1, & \text{if } 0 \leq t \leq \theta_1 \\ \gamma(t), & \text{if else.} \end{cases}$$

(b) The subclass of these tests that do not depend on θ_1 is the class of tests ϕ with

$$\phi(t) = \begin{cases} 1, & \text{if } \theta_0 < t < \infty \\ \gamma(t), & \text{if else} \end{cases}$$

where $E_{\theta_0}\gamma(T) = \alpha$.

(c) The test $\phi(t) = 1\{t > \theta_0\} + \alpha 1\{t \leq \theta_0\}$ is of the form of the tests in (b) with $\gamma(t) = \alpha$ for $0 \leq t \leq \theta_0$, and hence is UMP of size α for testing $\theta = \theta_0$ versus $\theta > \theta_0$. To see that it is UMP for testing $\theta \leq \theta_0$ versus $\theta > \theta_0$, we first compute its power function to confirm that it is of size α for the composite null hypothesis $\theta \leq \theta_0$. The power is

$$\begin{aligned} \beta_\phi(\theta) &= E_\theta\phi(T) = P_\theta(T > \theta_0) + \alpha P_\theta(T \leq \theta_0) \\ &= \{1 - (\frac{\theta_0}{\theta})^n\} 1_{(\theta_0, \infty)}(\theta) + \alpha 1_{[0, \theta_0]}(\theta) + \alpha \left(\frac{\theta_0}{\theta}\right)^n 1_{(\theta_0, \infty)}(\theta) \\ &= \{1 - (\frac{\theta_0}{\theta})^n(1 - \alpha)\} 1_{(\theta_0, \infty)}(\theta) + \alpha 1_{[0, \theta_0]}(\theta). \end{aligned}$$

Thus we see that $\beta_\phi(\theta) = \alpha$ for $\theta \leq \theta_0$, and ϕ is of size α for $\theta \leq \theta_0$. Since the class of size α tests for testing $\theta = \theta_0$ is a larger class than the class of size α tests for testing $\theta \leq \theta_0$, and since ϕ is UMP in the larger class, it is also UMP in the smaller class. Hence ϕ is UMP for testing $\theta \leq \theta_0$ versus $\theta > \theta_0$. But the competing test $\phi_0(t) = 1\{t > (1 - \alpha)^{1/n}\theta_0\}$ has power function

$$\begin{aligned} \beta_{\phi_0}(\theta) &= E_\theta\phi_0(T) = P_\theta(T > \theta_0(1 - \alpha)^{1/n}) \\ &= 1 - \{(1 - \alpha) \left(\frac{\theta_0}{\theta}\right)^n 1_{(\theta_0(1 - \alpha)^{1/n}, \infty)}(\theta) + 1_{[0, \theta_0(1 - \alpha)^{1/n}]}(\theta)\}, \end{aligned}$$

so the power function of the test ϕ_0 is strictly below that of the test ϕ on the set $[0, \theta_0)$. Hence ϕ is inadmissible and the test ϕ_0 is also UMP.

(d) The test $\phi(t) = 1 - 1_{(\theta_0\alpha^{1/n}, \theta_0]}(t)$ is of size α for testing $\theta = \theta_0$ versus $\theta \neq \theta_0$ since

$$E_{\theta_0}\phi(T) = P_{\theta_0}(T \leq \theta_0\alpha^{1/n}) = \left(\frac{\theta_0\alpha^{1/n}}{\theta_0}\right)^n = \alpha.$$

Furthermore it is of the form of the class of all UMP tests for testing $\theta = \theta_0$ versus $\theta > \theta_0$, hence it UMP among the size α tests for $\theta > \theta_0$. For testing $\theta = \theta_0$ versus $\theta = \theta_1 < \theta_0$, the NP Pearson tests of the form $\phi_1(t) = \gamma(t)1_{[0, \theta_0]}(t)$ are most powerful of their size. But the test ϕ is of this form (with $\gamma(t) = 1_{[0, \theta_0\alpha^{1/n}]}(t)$), is in this class and does not depend on $\theta_1 < \theta_0$. Hence ϕ is UMP for testing $\theta = \theta_0$ versus $\theta \neq \theta_0$. [This is an unusual situation in which we get “something for free” from the structure of the uniform distributions. Usually two-sided tests are *not* UMP!]

5. Let X and Y be random variables with joint density

$$p_{X,Y}(x, y) = \lambda\mu \exp(-\lambda x - \mu y)1_{(0, \infty)}(x)1_{(0, \infty)}(y).$$

- (a) Find a UMP unbiased test of size $\alpha = .2$ for testing $H_0 : \lambda \leq \mu + 1$ versus $H_1 : \lambda > \mu + 1$.
- (b) Find a UMP unbiased test of size $\alpha = .2$ for testing $H_0 : \lambda = \mu$ versus $H_1 : \lambda \neq \mu$.
- (c) Find a UMP unbiased test of size $\alpha = .2$ for testing $H_0 : \lambda \geq 2\mu$ versus $H_1 : \lambda < 2\mu$.
- (d) What happens when X_1, \dots, X_m are i.i.d. Exponential(λ) and Y_1, \dots, Y_n are i.i.d. Exponential(μ)?

Solution: When $X \sim \exp(\lambda)$ and $Y \sim \exp(\mu)$ we have

$$p_{\lambda, \mu}(x, y) = \lambda\mu \exp(-\lambda x - \mu y)1_{(0, \infty)}(x)1_{(0, \infty)}(y).$$

(a) For testing $H : \lambda \leq \mu + 1$ versus $K : \lambda > \mu + 1$ we rewrite the density as follows:

$$\begin{aligned} p_{\lambda, \mu}(x, y) &= \lambda\mu \exp(-\lambda x - \mu y) \\ &= \lambda\mu \exp((\lambda - \mu)y - \lambda(x + y)) \\ &= \lambda\mu \exp(\theta U(x, y) + \xi T(x, y)) \end{aligned}$$

where $\theta \equiv \lambda - \mu$, $U(x, y) \equiv y$, $\xi \equiv -\lambda$, and $T(x, y) \equiv x + y$. Since $\lambda \leq \mu + 1$ is equivalent to $\lambda - \mu = \theta \leq 1 \equiv \theta_0$, our theory for exponential families applies, and the UMP unbiased test of H versus K is given by

$$\phi(X, Y) = \begin{cases} 1 & \text{if } Y > c_\alpha(T) \\ \gamma(T) & \text{if } Y = c_\alpha(T) \\ 0 & \text{if } Y < c_\alpha(T) \end{cases}$$

where c_α and $\gamma(\alpha)$ satisfy $E\{\phi(X, Y)|T\} = \alpha$. In this case, the conditional distribution of Y given $T = X + Y$ on the boundary $\Theta_B = \{(\lambda - 1, \lambda) : \lambda \geq 1\}$ is given by

$$f_{Y|T}(y|t) = \frac{e^y}{e^t - 1} 1_{[0, t]}(y).$$

Therefore $1 - F_{Y|T}(y|t) = 1 - (e^y - 1)/(e^t - 1)$ and for $\alpha = .2$ the critical point for the conditional test is given by

$$c_\alpha(T) = \log(\exp(T) - (\exp(T) - 1)/5) = \log((4/5)\exp(T) + 1/5), \quad \gamma(T) = 0.$$

(b) For testing $H : \lambda = \mu$ versus $K : \lambda \neq \mu$, the same rewrite of the density as in (a) works. Now we have $\lambda = \mu$ is equivalent to $\mu - \lambda = 0 \equiv \theta_0$, and $\lambda \neq \mu$ is equivalent to $\mu - \lambda \neq 0 \equiv \theta_0$, so our theory for exponential families applies, and the UMP unbiased test of H versus K is given by

$$\phi(X, Y) = \begin{cases} 1 & \text{if } Y > c_2(T) \text{ or } Y < c_1(T) \\ \gamma_i(T) & \text{if } Y = c_1(T) \text{ or } Y = c_2(T) \\ 0 & \text{if } c_1(T) < Y < c_2(T) \end{cases}$$

where the c_1 , c_2 , γ_1 and γ_2 are determined so that $E\{\phi(X, Y)|T\} = \alpha$. In this case the conditional distribution of Y given T on $\Theta_B = \{(\lambda, \lambda) : \lambda \geq 0\}$ is Uniform(0, T), and hence the conditional distribution of Y/T given T is Uniform(0, 1), and this is independent of T . Hence the UMPU test of H versus K of size .2 is given by “reject H if $Y/T < .1$ or $Y/T > .9$ ”.

(c) For testing $H : \lambda \geq 2\mu$ versus $K : \lambda < 2\mu$, we need a somewhat different rewrite of the joint density. Now

$$\begin{aligned} p_{\lambda, \mu}(x, y) &= \lambda\mu \exp(-\lambda x - \mu y) \\ &= \lambda\mu \exp(-(\lambda - 2\mu)x - \mu(2x + y)) \\ &= \lambda\mu \exp(\theta U(x, y) + \xi T(x, y)) \end{aligned}$$

where $\theta \equiv 2\mu - \lambda$, $U(x, y) \equiv x$, $\xi \equiv -\mu$, and $T(x, y) \equiv 2x + y$. Since $\lambda \geq 2\mu$ is equivalent to $2\mu - \lambda \equiv \theta \leq 0 \equiv \theta_0$, (and $\lambda < 2\mu$ is equivalent to $2\mu - \lambda = \theta > 0 \equiv \theta_0$), our theory for exponential families applies, and the UMP unbiased test of H versus K is given by

$$\phi(X, Y) = \begin{cases} 1 & \text{if } X > c_\alpha(T) \\ \gamma(T) & \text{if } X = c_\alpha(T) \\ 0 & \text{if } X < c_\alpha(T) \end{cases}$$

where $c_\alpha(T)$ and $\gamma(T)$ satisfy $E\{\phi(X, Y)|T\} = \alpha$. In this case the conditional distribution of X given T is Uniform(0, $T/2$), so $2X/T$ is Uniform(0, 1) and independent of T . Hence the UMPU test of H versus K of size $\alpha = .2$ is given by “reject H if $2X/T > .8$ ”.

When we observe X_1, \dots, X_m are i.i.d. Exponential(λ) and Y_1, \dots, Y_n are i.i.d. Exponential(μ), then the distribution of the observations is given by

$$\begin{aligned}
p_{\lambda, \mu}(\underline{x}, \underline{y}) &= \lambda^m \mu^n \exp(-\lambda \sum_{i=1}^m x_i - \mu \sum_{j=1}^n y_j) \\
&= \lambda^m \mu^n \exp((\lambda - \mu) \sum_{j=1}^n y_j - \lambda(\sum_{i=1}^m x_i + \sum_{j=1}^n y_j)) \\
&= \lambda^m \mu^n \exp(\theta U(\underline{x}, \underline{y}) + \xi T(\underline{x}, \underline{y}))
\end{aligned}$$

where $\theta \equiv \lambda - \mu$, $U(\underline{x}, \underline{y}) \equiv \sum y_j$, $\xi \equiv -\lambda$, and $T(\underline{x}, \underline{y}) \equiv \sum x_i + \sum y_j$. This rewrite works for (a) and (b), and a similar rewrite works for (c) with the new $U = \sum x_i$, $T = 2 \sum x_i + \sum y_j$. The form of the tests in (a) - (c) remains the same with the new U and T , and all that remains is to calculate the conditional distributions of U given T . In (a) this density is given by

$$f_{U|T}(u|t) = \frac{u^{n-1}(t-u)^{m-1}e^u}{\int_0^t v^{n-1}(t-v)^{m-1}e^v dv}.$$

In (b) it is easily found that $V \equiv U/T \sim \text{Beta}(n, m)$, and the test can be carried out unconditionally using tables of the Beta distributions. In (c) $2U \equiv \sum 2X_i \sim \text{Gamma}(m, \mu)$ and $V \equiv \sum Y_j \sim \text{Gamma}(n, \mu)$ on the boundary $\lambda = 2\mu$, so $2\mu U \sim \chi_{2m}^2$ and $\mu V \sim \chi_{2n}^2$. Therefore

$$\frac{2U/(2m)}{V/(2n)} = \frac{n}{m} 2 \frac{U}{V} \sim F_{2m, 2n}$$

and since $2U/T = 2U/(2U + V) = (2U/V)(1 + (2U/V))$ is a monotone increasing function of $2U/V$, the UMPU test can be carried out unconditionally using tables of the $F_{2m, 2n}$ distributions.




Figure 1: Plots of fixed θ power approximations for $n = 3, 6, 9, 12, 15$.




Figure 2: Plot of local and fixed θ power approximations, $n = 3$.




Figure 3: Plot of local and fixed θ power approximations, $n = 6$.




Figure 4: Plot of local and fixed θ power approximations, $n = 9$.




Figure 5: Plot of local and fixed θ power approximations, $n = 12$.




Figure 6: Plot of local and fixed θ power approximations, $n = 15$.