

Statistics 582, Problem Set 8

Wellner; 2/23/2011

Reading: Chapter 6, sections 6.1 and 6.2 (through page 19);
Lehmann and Romano, Chapter 3.

Due: Wednesday, March 2, 2010.

1. A random variable X takes on the values 1, 2, 3, 4 with probability distribution $p_0(x)$ or $p_1(x)$ as follows:

| | | | | |
|----------|-----|-----|-----|-----|
| x | 1 | 2 | 3 | 4 |
| $p_0(x)$ | .18 | .06 | .36 | .40 |
| $p_1(x)$ | .36 | .18 | .24 | .22 |

- (a) Find a most powerful test of size $\alpha = .2$ for testing p_0 versus p_1 and determine its power.
 (b) Find a test ϕ which minimizes the sum of risks $a + b$ where $a = E_0\phi$ and $b = E_1(1 - \phi)$.
2. (Problem 3.6, Lehmann and Romano, TSH, page 93.) Suppose that P_0 , P_1 , and P_2 be the probability distributions assigning to the integers 1, . . . , 6 the following probabilities:

| | | | | | | |
|----------|-----|-----|-----|-----|-----|-----|
| x | 1 | 2 | 3 | 4 | 5 | 6 |
| $p_0(x)$ | .03 | .02 | .02 | .01 | 0 | .92 |
| $p_1(x)$ | .06 | .05 | .08 | .02 | .01 | .78 |
| $p_2(x)$ | .09 | .05 | .12 | 0 | .02 | .72 |

Determine whether there exists a level- α test of $H : P = P_0$ which is UMP against the alternatives P_1 and P_2 when:

- (i) $\alpha = .01$; (ii) $\alpha = .05$; (iii) $\alpha = .07$.
3. (Problem 3.7, Lehmann and Romano, TSH, page 94, modified) Suppose that the distribution of X is given by

| | | | | |
|---------------|------------|----------|---------------|-----------------|
| x | 0 | 1 | 2 | 3 |
| $p_\theta(x)$ | $\theta/2$ | θ | $.9 - \theta$ | $.1 - \theta/2$ |

where $0 < \theta < .2$. For testing $H : \theta = .05$ against $\theta > .05$ at level $\alpha = .05$, determine which of the following tests (if any) is UMP:

- (i) $\phi(0) = 1, \phi(1) = \phi(2) = \phi(3) = 0$;
- (ii) $\phi(1) = .5, \phi(0) = \phi(2) = \phi(3) = 0$;
- (iii) $\phi(3) = 1, \phi(0) = \phi(1) = \phi(2) = 0$.

4. Suppose that X_1, \dots, X_n are i.i.d. $N(\theta, \sigma^2)$.

- (a) Suppose that $\sigma = \sigma_0$ is known. Consider testing $H : \theta = \theta_0 = 0$ versus $K : \theta = \theta_1 = 1$. In the spirit of chapter 5, plot $(R(\theta_0, \phi), R(\theta_1, \phi))$ for your favorite family of tests ϕ . Find the entire risk body and plot it.
- (b) What happens to the risk body as n grows or as $\sigma_0 \rightarrow 0$?
- (c) What happens to the risk body as θ_1 decreases toward $\theta_0 = 0$?
- (d) What happens to the risk bodies $\{(R(\theta_0, \phi), R(\theta_{1,n}, \phi)) : n \geq 1\}$ when $\theta_1 \equiv \theta_{1,n} \equiv \theta_0 + cn^{-1/2}$?

5. Consider the logistic distributions with location parameter θ having density $p_\theta(x) = g(x - \theta)$ where

$$g(x) = \frac{\exp(x)}{(1 + \exp(x))^2} = \frac{1}{2(1 + \cosh(x))} = \frac{e^{-x}}{(1 + e^{-x})^2}.$$

- (a) Show that the family $\{p_\theta : \theta \in \mathbb{R}\}$ has monotone likelihood ratio in x .
- (b) Unfortunately the result of (a) does not carry over to a sample of size n . If X_1, \dots, X_n are i.i.d. P_θ with density p_θ as in (a), then there is no $T(\underline{X})$ for which the MLR property holds. Nevertheless we can look for locally best tests. Find the locally best test of $H_0 : \theta = 0$ versus $H_1 : \theta > 0$. How would you carry out this test?

6. **Optional bonus problem 1.** Show that the Cauchy scale family of distributions given by

$$p_\theta(x) = \frac{1}{\pi\theta} \frac{1}{1 + (x/\theta)^2}$$

does not have monotone likelihood ratio, but that the distribution of the sufficient statistic $|X|$ (where $X \sim P_\theta$ with Cauchy density p_θ as in the last display) does have monotone likelihood ratio.

7. **Optional bonus problem 2.** Can you prove anything about monotonicity of the power functions of the class of tests derived in problem 5 (b)?