

## Statistics 583, Problem Set 5 Solutions

Wellner; 5/2/2007

1. Exercise 3.8.1, Wasserman, page 39. [Hint: the formula given by Wasserman, page 29, is not correct.] Under what additional hypotheses can we establish  $\sqrt{n}(T(\mathbb{F}_n) - T(F)) \rightarrow N(0, E_F \psi_F^2(X))$ ? (Here my  $\psi_F$  equals Wasserman's  $L_F$ .)

**Solution:** (a) To find the influence function of  $T(F) = \int (x - \mu)^3 dF(x) / \sigma(F)^3$ , let  $F_t \equiv (1 - t)F + tG$ . Then we need to compute  $(d/dt)T(F_t)|_{t=0}$ . But, by using the calculations in examples 7.4.2 and 7.4.3,

$$\begin{aligned}
 \frac{d}{dt}T(F_t)|_{t=0} &= \frac{d}{dt} \frac{\int (x - \mu(F_t))^3 dF_t(x)}{[\sigma^2(F_t)]^{3/2}} \Big|_{t=0} \\
 &= \frac{\int (x - \mu(F_t))^3 d(G - F)(x)}{[\sigma^2(F_t)]^{3/2}} \Big|_{t=0} \\
 &\quad - \frac{3}{2} \frac{\int (x - \mu(F_t))^3 dF_t(x)}{[\sigma^2(F_t)]^{5/2}} \frac{d}{dt} \sigma^2(F_t) \Big|_{t=0} \\
 &\quad - 3 \frac{\int (x - \mu(F_t))^2 dF_t(x)}{\sigma(F_t)^3} \frac{d}{dt} \mu(F_t) \Big|_{t=0} \\
 &= \int \left( \frac{x - \mu(F)}{\sigma(F)} \right)^3 d(G - F)(x) \\
 &\quad - \frac{3}{2} T(F) \frac{1}{\sigma^2(F)} \left\{ \int (x - \mu(F))^2 dG(x) - \sigma^2(F) \right\} \\
 &\quad - 3 \int \left( \frac{x - \mu(F)}{\sigma(F)} \right) dG(x) \\
 &= \int \left( \frac{x - \mu(F)}{\sigma(F)} \right)^3 dG(x) - T(F) \\
 &\quad - \frac{3}{2} T(F) \int \left\{ \left( \frac{x - \mu(F)}{\sigma(F)} \right)^2 - 1 \right\} dG(x) \\
 &\quad - 3 \int \left( \frac{x - \mu(F)}{\sigma(F)} \right) dG(x).
 \end{aligned}$$

Hence by taking  $G = \delta_x$  we find the influence function of  $T(F)$ :

$$\begin{aligned} \dot{T}(F; \delta_x - F) &= \left( \frac{x - \mu(F)}{\sigma(F)} \right)^3 - T(F) - \frac{3}{2}T(F) \left\{ \left( \frac{x - \mu(F)}{\sigma(F)} \right)^2 - 1 \right\} \\ &\quad - 3 \left( \frac{x - \mu(F)}{\sigma(F)} \right) \\ &\equiv \psi_F(x). \end{aligned} \tag{1}$$

Note that this derivation does not seem to agree with the result stated on page 29 of Wasserman: the third term here does not appear in Wasserman's claimed influence function.

(b) Here is a direct calculation to see the result in (a) another way. Write

$$\begin{aligned} &\sqrt{n}(T(\mathbb{F}_n) - T(F)) \\ &= \frac{1}{\sigma(\mathbb{F}_n)^3} \sqrt{n} \left\{ \int (x - \mu(\mathbb{F}_n))^3 d\mathbb{F}_n(x) - \int (x - \mu(F))^3 dF(x) \right\} \\ &\quad + \int (x - \mu(F))^3 dF(x) \sqrt{n} \left\{ \frac{1}{\sigma(\mathbb{F}_n)^3} - \frac{1}{\sigma(F)^3} \right\} \\ &\equiv A_n + B_n. \end{aligned}$$

To understand  $A_n$ , write

$$\begin{aligned} (x - \mu(\mathbb{F}_n))^3 &= (x - \mu(F) - (\mu(\mathbb{F}_n) - \mu(F)))^3 \equiv (a - b)^3 \\ &= a^3 - 3a^2b + 3ab^2 + b^3 \\ &= (x - \mu(F))^3 - 3(x - \mu(F))^2(\mu(\mathbb{F}_n) - \mu(F)) \\ &\quad + 3(x - \mu(F))(\mu(\mathbb{F}_n) - \mu(F))^2 + (\mu(\mathbb{F}_n) - \mu(F))^3. \end{aligned}$$

Thus we see that

$$\begin{aligned} A_n &= \frac{1}{\sigma(F)^3} \left\{ \sqrt{n} \int (x - \mu(F))^3 d(\mathbb{F}_n(x) - F(x)) \right. \\ &\quad - 3 \int (x - \mu(F))^2 d\mathbb{F}_n(x) \sqrt{n}(\mu(\mathbb{F}_n) - \mu(F)) \\ &\quad \left. + 3 \int (x - \mu(F)) d\mathbb{F}_n(x) \sqrt{n}(\mu(\mathbb{F}_n) - \mu(F))^2 + \sqrt{n}(\mu(\mathbb{F}_n) - \mu(F))^3 \right\} + o_p(1) \\ &= \frac{1}{\sigma(F)^3} \left\{ \sqrt{n} \int (x - \mu(F))^3 d(\mathbb{F}_n(x) - F(x)) \right. \\ &\quad \left. - 3\sigma^2(F) \sqrt{n} \int (x - \mu(F)) d(\mathbb{F}_n(x) - F(x)) \right\} \\ &\quad + o_p(1). \end{aligned}$$

For  $B_n$  we can write, with  $m_3(F) \equiv \int (x - \mu(F))^3 dF(x)$

$$\begin{aligned}
 B_n &= m_3(F) \sqrt{n} \left\{ \frac{1}{\sigma(\mathbb{F}_n)^3} - \frac{1}{\sigma(F)^3} \right\} \\
 &= -\frac{m_3(F)}{\sigma(F)^3 \sigma(\mathbb{F}_n)^3} \sqrt{n} \{ \sigma(\mathbb{F}_n)^3 - \sigma(F)^3 \} \\
 &= -\frac{m_3(F)}{\sigma^2(F)^3} \sqrt{n} \{ \sigma^2(\mathbb{F}_n)^{3/2} - \sigma^2(F)^{3/2} \} + o_p(1) \\
 &= -\frac{m_3(F)}{\sigma^2(F)^3} \frac{3}{2} \sigma(F) \sqrt{n} (\sigma^2(\mathbb{F}_n) - \sigma^2(F)) + o_p(1) \\
 &= -\frac{m_3(F)}{\sigma^3(F)} \frac{3}{2\sigma^2(F)} \sqrt{n} \int \{ (x - \mu(F))^2 - \sigma^2(F) \} d\mathbb{F}_n(x).
 \end{aligned}$$

Putting the  $A_n$  and  $B_n$  pieces together we see that we have complete agreement with the result of the influence function calculation:

$$\sqrt{n}(T(\mathbb{F}_n) - T(F)) = \sqrt{n} \int \psi_F(x) d\mathbb{F}_n(x)$$

where  $\psi_F(x)$  is as given in (1). It is clear (from the Central Limit Theorem) that this is asymptotically normal if  $E_F X^6 < \infty$ .

When I use the influence function derived here to obtain an estimator of the Standard Error of the skewness estimator for the nerve data treated in Wasserman's example 3.10, page 29, I get  $\hat{s}e = .163$  rather than Wasserman's estimate of .18, a slight reduction. The resulting confidence interval for the population skewness is  $1.76 \pm 2(.163) = (1.434, 2.086)$ .

2. (a) Exercise 2.7.9, Wasserman, page 24.  
 (b) What additional hypotheses are needed to show that  $\sqrt{n}(T(\mathbb{F}_n) - T(F))$  is asymptotically normal for this particular functional  $T(F)$ ?  
*Reminder:* This exercise gives the same result as we derived last Fall in Stat 581.

**Solution:** (a) I will take a slightly different tack than Wasserman: write

$$T(F) = \frac{E_F(X - \mu_X)(Y - \mu_Y)}{\sqrt{Var_F(X)Var_F(Y)}} \equiv \rho(F) = g(T_1(F), T_2(F), T_3(F))$$

where

$$\begin{aligned}
g(u, v, w) &\equiv \frac{u}{\sqrt{vw}}, \\
T_1(F) &= E_F(X - \mu_X)(Y - \mu_Y), \\
T_2(F) &= \text{Var}_{F_X}(X), \\
T_3(F) &= \text{Var}_{F_Y}(Y).
\end{aligned}$$

From example 4.3 we know that

$$\begin{aligned}
\dot{T}_2(F; G - F) &= \int \psi_{2,F}(x) dG_X(x), & \psi_{2,F}(x) &= (x - \mu_X)^2 - \sigma_X^2, \\
\dot{T}_3(F; G - F) &= \int \psi_{3,F}(y) dG_Y(y), & \psi_{3,F}(y) &= (y - \mu_Y)^2 - \sigma_Y^2.
\end{aligned}$$

Now

$$\begin{aligned}
\left. \frac{d}{dt} T_1(F_t) \right|_{t=0} &= \left. \frac{d}{dt} \int xy dF_t(x, y) \right|_{t=0} - \left. \frac{d}{dt} \mu_X(F_t) \mu_Y(F_t) \right|_{t=0} \\
&= \int \int xy d(G - F)(x, y) - \mu_Y \int (x - \mu_X) dG_X(x) - \mu_X \int (y - \mu_Y) dG_Y(y) \\
&= \int \int \psi_{1,F}(x, y) dG(x, y)
\end{aligned}$$

where

$$\psi_{1,F}(x, y) = (x - \mu_X)(y - \mu_Y) - E_F(X - \mu_X)(Y - \mu_Y).$$

Since  $\nabla g(u, v, w) = (1, -u/(2v), -u/(2w))/\sqrt{vw}$ , it follows by the chain rule that

$$\begin{aligned}
\dot{T}(F; G - F) &= \int \int \frac{\psi_{1,F}(x, y)}{\sigma_X \sigma_Y} dG(x, y) \\
&\quad - \frac{T(F)}{2} \int \frac{\psi_{2,F}(x)}{\sigma_X^2} dG_X(x) - \frac{T(F)}{2} \int \frac{\psi_{3,F}(y)}{\sigma_Y^2} dG_Y(y) \\
&= \int \int \psi_F(x, y) dG(x, y)
\end{aligned}$$

where

$$\begin{aligned}
\psi_F(x, y) &= \frac{(x - \mu_X)(y - \mu_Y)}{\sigma_X \sigma_Y} - T(F) - \frac{T(F)}{2} \left\{ \frac{(x - \mu_X)^2}{\sigma_X^2} - 1 + \frac{(y - \mu_Y)^2}{\sigma_Y^2} - 1 \right\} \\
&= \tilde{x}\tilde{y} - \rho(F) - \frac{\rho(F)}{2} \{ \tilde{x}^2 - 1 + \tilde{y}^2 - 1 \}.
\end{aligned}$$

Thus I get a “centered” version of Wasserman’s formula (page 21, line 2).

(b) To show that  $\sqrt{n}(T(\mathbb{F}_n) - T(F)) \rightarrow_d N(0, E_F \psi_F^2(X, Y))$ , it is clear from the influence calculation in (a) that we will need to assume that  $E(X^4) < \infty$  and  $E(Y^4) < \infty$ .

3. Suppose that we observe  $X_1, \dots, X_n$  i.i.d.  $P$  on  $\mathbb{R}^+ = [0, \infty)$  and assume that  $P \in \mathcal{P}_0 \equiv \{P_\theta : (dP_\theta/d\lambda) = p_\theta, \theta \in \Theta\}$  where  $\theta = (\alpha, \beta) \in (0, \infty)^2$  and  $p_\theta = p_{\alpha, \beta}$  is the Weibull density given by  $p_\theta(x) = (\beta/\alpha)(x/\alpha)^{\beta-1} \exp(-(x/\alpha)^\beta) 1_{(0, \infty)}(x)$ . From Lehmann and Romano, TPE, Example 6.6.1 (page 468) and problems 6.6.1 - 6.6.3 (page 509), we know that the maximum likelihood estimator  $\hat{\theta}_n = (\hat{\alpha}_n, \hat{\beta}_n)$  exists and is unique if  $0 < X_{(1)} < X_{(n)}$ .

(a) If, in fact,  $P \notin \mathcal{P}_0$ , to what function of  $P$  do you expect  $\hat{\theta}_n = (\hat{\alpha}_n, \hat{\beta}_n)$  converges (in probability)? [Hint: use the development in example 6.6.1 of Lehmann and Romano to show that the solution of the population version of the score equations (with sampling from  $P \notin \mathcal{P}$ ) leads to  $\alpha(P) = \{E_P(X^\beta)\}^{1/\beta}$  where  $\beta$  is the solution of

$$\frac{E_P(X^\beta \log X)}{E_P X^\beta} - \frac{1}{\beta} = E_P(\log X),$$

assuming that  $E_P(X^\beta |\log X|) < \infty$ . ]

(b) Show heuristically that  $\theta(P) = (\alpha(P), \beta(P))$  minimizes  $K(P, P_\theta)$  over  $\Theta$ .

(c) In particular, if  $P$  has Gamma(4, 1) density  $p(x) = (x^3 e^{-x}/3!) 1_{(0, \infty)}(x)$  find  $(\alpha, \beta) = (\alpha(P), \beta(P))$  corresponding to the “best-fitting” member of the Weibull family  $P_{(\alpha(P), \beta(P))}$ . Plot both  $p$  and  $p_{(\alpha(P), \beta(P))}$  as functions of  $x$ .

**Solution:** (a) From our discussions in class we expect that  $\hat{\theta}_n = (\hat{\alpha}_n, \hat{\beta}_n)$  will converge to the solution  $T(P) \equiv \theta(P) = (\alpha(P), \beta(P))$  of

$$P \dot{\mathbf{i}}_\theta(X; \theta(P)) = \int \dot{\mathbf{i}}_\theta(x; \theta(P)) dP(x) = 0 \tag{2}$$

where  $\theta(P) = (\alpha(P), \beta(P)) \in \mathbb{R}^2$  and  $\dot{\mathbf{i}}_\theta(\cdot; \theta)$  is the vector of score functions for the Weibull model. Since

$$\dot{\mathbf{i}}_\theta(x; \theta) = \begin{pmatrix} \frac{\beta}{\alpha} \left( \left( \frac{x}{\alpha} \right)^\beta - 1 \right) \\ \frac{1}{\beta} \left( 1 - \log \left\{ \left( \frac{x}{\alpha} \right)^\beta \right\} \left\{ \left( \frac{x}{\alpha} \right)^\beta - 1 \right\} \right) \end{pmatrix},$$

it follows that  $\theta(P) = (\alpha(P), \beta(P))$  satisfies

$$\int (x/\alpha)^\beta dP(x) = 1, \quad \text{and} \\ \int \left( 1 - \log \left\{ \left( \frac{x}{\alpha} \right)^\beta \right\} \left\{ \left( \frac{x}{\alpha} \right)^\beta - 1 \right\} \right) dP(x) = 0, .$$

The first equation yields

$$\alpha(P) = \{E_P X^\beta\}^{1/\beta},$$

and hence the second equation can be re-written as

$$\begin{aligned} 1 &= E_P \left\{ \beta \log X \left( \left( \frac{X}{\alpha} \right)^\beta - 1 \right) \right\} \\ &= \beta \left\{ \frac{E_P(X^\beta \log X)}{E_P X^\beta} - E_P \log X \right\} \end{aligned}$$

or, equivalently,

$$\frac{E_P(X^\beta \log X)}{E_P X^\beta} - \frac{1}{\beta} = E_P(\log X). \quad (3)$$

Letting

$$h(\beta) \equiv \frac{E_P(X^\beta \log X)}{E_P X^\beta} - \frac{1}{\beta},$$

we have

$$\begin{aligned} h'(\beta) &= \frac{E_P\{X^\beta(\log X)^2\}}{E(X^\beta)} - \frac{[E_P(X^\beta \log X)]^2}{[E(X^\beta)]^2} + \frac{1}{\beta^2} \\ &\geq \text{Var}_Q(\log X) \quad \text{under } Q \text{ with } dQ(x) \equiv \frac{x^\beta}{E_P(X^\beta)} dP(x) \\ &> 0 \end{aligned}$$

unless  $P$  is degenerate. Thus if  $P$  is non-degenerate  $h(\beta)$  is monotone increasing with

$$-\infty = \lim_{\beta \downarrow 0} h(\beta) < E_P(\log X) < \lim_{\beta \rightarrow \infty} h(\beta) = \sup\{\text{support}(P)\}.$$

Thus, in parallel to the finite sample situation as in the problems in Lehmann and Romano, the equation (3) has a unique finite solution in  $(0, \infty)$  if  $P$  is non-degenerate, does not have positive mass at 0 and provided that  $X$  has sufficiently many moments: e.g. if  $E_P X^r < \infty$  for some  $r > \beta(P)$ .

(b) We did this in class. Here is a heuristic argument showing why this should be true: Note that for many cases we have

$$\begin{aligned} \hat{\theta}_n &= \operatorname{argmax}_\theta n^{-1} l_n(\theta) = \operatorname{argmax}_\theta \mathbb{P}_n(\log p_\theta) \\ &\rightarrow_p \operatorname{argmax}_\theta P(\log p_\theta) = \operatorname{argmax}_\theta \int \log p_\theta(x) dP(x). \end{aligned}$$

Now

$$\begin{aligned}
P(\log p_\theta) &= P(\log p) + P \log \left( \frac{p_\theta}{p} \right) \\
&= P(\log p) - P \log \left( \frac{p}{p_\theta} \right) \\
&= P(\log p) - K(P, P_\theta).
\end{aligned}$$

Thus

$$\operatorname{argmax}_\theta \int \log p_\theta(x) dP(x) = \operatorname{argmin}_\theta K(P, P_\theta) \equiv \theta(P).$$

If we can interchange differentiation and integration it follows that

$$\nabla_\theta K(P, P_\theta) = \int p(x) \dot{\mathbf{i}}_\theta(x; \theta) d\mu(x) = \int \dot{\mathbf{i}}_\theta(x; \theta) dP(x),$$

so the relation (2) is obtained by setting this gradient vector equal to 0.

(c) When  $P$  has density  $p$  given by  $p(x) = (x^3 e^{-x}/3!)1_{(0,\infty)}(x)$  we compute

$$E_P X^\beta = \frac{1}{6} \int_0^\infty x^{\beta+3} e^{-x} dx = \frac{1}{6} \Gamma(\beta + 4), \quad (4)$$

$$E_P(\log X) = \frac{1}{6} \int_0^\infty x^3 \log x e^{-x} dx = (11 - 3\gamma)/6, \quad \text{and} \quad (5)$$

$$E_P(X^\beta \log X) = \frac{1}{6} \int_0^\infty x^{\beta+3} (\log x) e^{-x} dx \quad (6)$$

$$= \frac{1}{6} \Gamma(4 + \beta) \psi(4 + \beta) \quad (7)$$

where  $\gamma = 0.577216\dots$  is Euler's constant and  $\psi(x) = (\log \Gamma(x))'$  is the digamma function. Thus (3) becomes, using  $\Gamma(1/2) = \sqrt{\pi}$ ,

$$\frac{\Gamma(4 + \beta) \psi(4 + \beta)}{\Gamma(4 + \beta)} - \frac{1}{\beta} = (11 - 3\gamma)/6,$$

or, equivalently,

$$\psi(4 + \beta) - \frac{1}{\beta} = (11 - 3\gamma)/6.$$

The left side is an increasing function of  $\beta$ , while the right side is a constant function of  $\beta$ , and they have a unique intersection at  $\beta = 2.12043\dots = \beta(P)$ . Then  $\alpha(P) = \{E_P X^\beta\}^{1/\beta} = 4.52813\dots$  by using (4) and  $\beta = 2.12043\dots$ , and the resulting Kullback-Leibler distance is  $K(P, P_{\theta(P)}) = 0.0200882$ . The following two figures (Figures 1 and 2) show the functions  $\alpha \mapsto K(\alpha, \beta(P))$  and  $\beta \mapsto K(\alpha(P), \beta)$ .




Figure 1: The function  $\alpha \mapsto K(\alpha, \beta(P))$




Figure 2: The function  $\beta \mapsto K(\alpha(P), \beta)$

Figure 3 shows  $p$  and the Weibull density  $p(\cdot; \theta(P))$  “closest” to  $p$ .




Figure 3: The density  $p$  (blue) and the Weibull density  $p(\cdot; \theta(P))$  (red) minimizing the Kullback-Leibler distance to  $p$

4. Suppose that  $\hat{\theta}_n$  is the MLE for the Weibull family as in problem 1 above, and that  $P \notin \mathcal{P}_0$ . Heuristically we expect that

$$\sqrt{n}(\hat{\theta}_n - \theta(P)) \rightarrow_d N_2(0, \Sigma(P)) \quad (8)$$

for some covariance matrix  $\Sigma = \Sigma(P)$  as  $n \rightarrow \infty$ .

- (a) What is the form of  $\Sigma$  that you expect in (8)?  
(b) What methods could be used to make these heuristics precise?

**Solution:** (a) From our heuristic computation of the influence function of a  $Z$ -estimator and from Huber's  $Z$ -theorem, we expect  $\Sigma(P)$  to be of the form

$$\Sigma(P) = \dot{\Psi}(P)^{-1} V(P) [\dot{\Psi}(P)^{-1}]^T$$

where  $\dot{\Psi}(P) \equiv \dot{\Psi}(\theta(P))$  is the derivative of  $\Psi(\theta) = P(\dot{\mathbf{I}}_\theta(X, \theta))$  at  $\theta(P)$ , i.e.

$$\dot{\Psi}(\theta(P)) = \int \ddot{\mathbf{I}}_{\theta\theta}(x, \theta(P)) dP(x),$$

and

$$\begin{aligned} V(P) &= E_P(\dot{\mathbf{I}}(X; \theta(P)) \dot{\mathbf{I}}(X; \theta(P))^T) \\ &= \begin{pmatrix} \left(\frac{\beta}{\alpha}\right)^2 E_P \left( \left(\frac{X}{\alpha}\right)^\beta - 1 \right)^2 & -\frac{1}{\alpha} E_P \left\{ \left( \left(\frac{X}{\alpha}\right)^\beta - 1 \right)^2 \log \left(\frac{X}{\alpha}\right)^\beta \right\} \\ -\frac{1}{\alpha} E_P \left\{ \left( \left(\frac{X}{\alpha}\right)^\beta - 1 \right)^2 \log \left(\frac{X}{\alpha}\right)^\beta \right\} & \frac{1}{\beta^2} E_P \left\{ \left( 1 - \log \left(\frac{X}{\alpha}\right)^\beta \left( \left(\frac{X}{\alpha}\right)^\beta - 1 \right) \right)^2 \right\} \end{pmatrix} \end{aligned}$$

The matrix  $\dot{\Psi}(P)$  can be written somewhat more explicitly by first calculating  $\ddot{\mathbf{I}}_{\theta\theta}$ , and then computing the necessary expectations: from the forms of  $\dot{\mathbf{I}}_{\alpha}$  and  $\dot{\mathbf{I}}_{\beta}$  we get

$$\begin{aligned}
\ddot{\mathbf{I}}_{\alpha,\alpha}(x, \theta) &= -\frac{\beta}{\alpha^2} \left( \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right) - \left( \frac{\beta}{\alpha} \right)^2 \left( \frac{x}{\alpha} \right)^{\beta}, \\
\ddot{\mathbf{I}}_{\alpha,\beta}(x, \theta) &= \frac{1}{\alpha} \left( \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right) + \frac{1}{\alpha} \left( \frac{x}{\alpha} \right)^{\beta} \log \left( \frac{x}{\alpha} \right)^{\beta}, \\
\ddot{\mathbf{I}}_{\beta,\beta}(x, \theta) &= -\frac{1}{\beta^2} \left( 1 - \log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} \left\{ \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right\} \right) \\
&\quad + \frac{1}{\beta} \left\{ -\log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} \left( \frac{x}{\alpha} \right)^{\beta} \log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} \right. \\
&\quad \quad \quad \left. - \left\{ \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right\} \log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} \right\} \\
&= -\frac{1}{\beta^2} \left( 1 - \log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} \left\{ \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right\} \right) \\
&\quad + \frac{1}{\beta^2} \left\{ -\log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\}^2 \left( \frac{x}{\alpha} \right)^{\beta} \right. \\
&\quad \quad \quad \left. 1 - \left\{ \left( \frac{x}{\alpha} \right)^{\beta} - 1 \right\} \log \left\{ \left( \frac{x}{\alpha} \right)^{\beta} \right\} - 1 \right\}.
\end{aligned}$$

Thus we compute

$$\dot{\Psi}(P) = \begin{pmatrix} -(\beta/\alpha)^2 & \alpha^{-1} E_P(X/\alpha)^{\beta} \log(X/\alpha)^{\beta} \\ \alpha^{-1} E_P(X/\alpha)^{\beta} \log(X/\alpha)^{\beta} & -\beta^{-2} E_P\{1 + (X/\alpha)^{\beta} (\log(X/\alpha)^{\beta})^2\} \end{pmatrix}$$

where  $(\alpha, \beta) \equiv (\alpha(P), \beta(P))$ .

(b) To make the heuristics in problem 3(a) precise, one can use the monotonicity of the “concentrated” version of the estimating equation for  $\beta$  along the lines of our discussion earlier for existence and uniqueness of the solution for  $\hat{\beta}_n$  together with Lemma 5.10 of van der Vaart (1998), page 47. To make the heuristics of part (a) and the claim (8) precise, one can use the  $Z$ -theorem we discussed in class in Stat 582 (see e.g. Theorem 5.21 of van der Vaart (1998), page 52) by working directly with the scores, or, alternatively, one can first reduce to the “concentrated” score equation for  $\beta$  and use the  $Z$ -theorem for  $\hat{\beta}_n$  separately. For a relevant “ $M$ -theorem”, see van der Vaart (1998), Theorem 5.23, page 53.