

Statistics 583, Problem Set 6

Wellner; 5/6/2009

Reading: Chapter 8, sections 8.1- 8.4, pages 1 - 18;

Wasserman, Chapters 2-3, pages 13-41.

Due: Wednesday, May 13, 2007

Reminder 1: Second (and last) make-up lecture, Friday, May 8, 11:30 AM.

1. Wasserman, example 3.10, page 29.

(a) In the 3rd line of this example, Wasserman says that the jackknife estimate of the standard error of the skewness estimator for the nerve data is .17. Check this. (Show your code or method of calculation.)

(b) I claimed in the solution to problem set #5, problem 1 (see page 3 of the solution set), that the estimated standard error using the delta method is .163 rather than .18 as Wasserman claimed. Check this. (Show your code or method of calculation.)

(c) On page 31, Wasserman claims that the bootstrap based on $B = 10^3$ replications gives .16 as a bootstrap estimate of the standard error. Try it yourself to see what you get. (Show your code or method of calculation.)

The data is posted in two forms at:

<http://www.stat.washington.edu/jaw/COURSES/580s/583/sp09.probsets.html/nrvdat>
and

<http://www.stat.washington.edu/jaw/COURSES/580s/583/sp09.probsets.html/nerve.dat>

(d) Plot the empirical distribution function of this data together with the MLE of the distribution function assuming: (i) an exponential distribution; (ii) a Weibull distribution. What alternative methods for (a) and (b) does this suggest?

2. (a) Given n distinct data items, show that the probability that a given data item does not appear in a bootstrap sample is $e_n = (1 - 1/n)^n$

(b) Show that $e_n \rightarrow e^{-1} \approx .368$ as $n \rightarrow \infty$.

(c) Hence show that the probability that each of B bootstrap samples contains an item i is $(1 - e_n)^B$. Evaluate this quantity for $n = 10, 20, 50, 100$ and $B = 10, 20, 50, 100$.

(d) Let $N_n \equiv \sum_{j=1}^n 1_{[M_j=0]}$ where $\underline{M} \equiv (M_1, \dots, M_n) \sim \text{Mult}_n(n, \underline{1}/n)$. Show that $E(n^{-1}N_n) = e_n$ as computed in (a).

3. Suppose that $T(F) = \text{Var}_F(X)$ so that $T_n \equiv T(\mathbb{F}_n) = n^{-1} \sum_{i=1}^n (X_i - \bar{X})^2$. Show that the jackknife estimate of the variance $\sigma_n^2(F) \equiv \text{Var}_F(T_n)$ is

$$\widehat{\text{Var}} = \frac{n^2}{(n-1)^3} (\widehat{\mu}_4 - \widehat{\mu}_2^2)$$

where $\widehat{\mu}_k \equiv n^{-1} \sum_{i=1}^n (X_i - \overline{X})^k$ for $k = 1, 2, \dots$. Hence, assuming that $EX^4 < \infty$, the jackknife estimate of variance is consistent for this T :

$$n\widehat{Var} \rightarrow_p \mu_4 - \mu_2^2 = \mu_2^2 \left\{ 2 + \frac{\mu_4}{\mu_2^2} - 3 \right\} = T_2(F)(2 + \gamma_2).$$

4. (a) Wasserman, problem 3.8.9, page 40.
(b) Wasserman, problem 3.8.13, page 41.
5. **Optional bonus problem:** Show that for linear statistics the jackknife and bootstrap estimates of bias are zero. (A linear statistic corresponds to a functional $T(F)$ of the form $T(F) = \int \psi dF$ for some function ψ for a distribution function F on R , or $T(P) = \int \psi dP$ for a probability distribution P on a general sample space \mathcal{X} .)
6. **Optional bonus problem:**