

Floral volatile alleles can contribute to pollinator-mediated reproductive isolation in monkeyflowers (*Mimulus*)

Kelsey J. R. P. Byers[†], James P. Vela, Foen Peng, Jeffrey A. Riffell and Harvey D. Bradshaw Jr*

Department of Biology, University of Washington, Seattle, WA 98195-1800, USA

Received 17 April 2014; revised 28 September 2014; accepted 6 October 2014; published online 16 October 2014.

*For correspondence (e-mail toby@uw.edu).

Accession numbers for sequences:

KF857265, *M. lewisii* LIMONENE-MYRCENE SYNTHASE gene, complete cds.
KF857264, *M. lewisii* LIMONENE-MYRCENE SYNTHASE mRNA, complete cds.
AH150308, *M. lewisii* LIMONENE-MYRCENE SYNTHASE protein product.
KM659024, *M. cardinalis* LIMONENE-MYRCENE SYNTHASE mRNA, exon3.
KF857262, *M. lewisii* OCIMENE SYNTHASE mRNA, complete cds.
AH150306, *M. lewisii* OCIMENE SYNTHASE protein product.
KF857263, *M. cardinalis* OCIMENE SYNTHASE mRNA, complete cds.
AH150307, *M. cardinalis* OCIMENE SYNTHASE protein product.

[†] Present address: Institute of Systematic Botany, University of Zürich, 8008 Zürich, Switzerland.

SUMMARY

Pollinator-mediated reproductive isolation is a major factor in driving the diversification of flowering plants. Studies of floral traits involved in reproductive isolation have focused nearly exclusively on visual signals, such as flower color. The role of less obvious signals, such as floral scent, has been studied only recently. In particular, the genetics of floral volatiles involved in mediating differential pollinator visitation remains unknown. The bumblebee-pollinated *Mimulus lewisii* and hummingbird-pollinated *Mimulus cardinalis* are a model system for studying reproductive isolation via pollinator preference. We have shown that these two species differ in three floral terpenoid volatiles – α -limonene, β -myrcene, and *E*- β -ocimene – that are attractive to bumblebee pollinators. By genetic mapping and *in vitro* analysis of enzyme activity we demonstrate that these interspecific differences are consistent with allelic variation at two loci, LIMONENE-MYRCENE SYNTHASE (*LMS*) and OCIMENE SYNTHASE (*OS*). *Mimulus lewisii* *LMS* (*MILMS*) and *OS* (*MIOS*) are expressed most strongly in floral tissue in the last stages of floral development. *Mimulus cardinalis* *LMS* (*McLMS*) is weakly expressed and has a nonsense mutation in exon 3. *Mimulus cardinalis* *OS* (*McOS*) is expressed similarly to *MIOS*, but the encoded *McOS* enzyme produces no *E*- β -ocimene. Recapitulating the *M. cardinalis* phenotype by reducing the expression of *MILMS* by RNA interference in transgenic *M. lewisii* produces no behavioral difference in pollinating bumblebees; however, reducing *MIOS* expression produces a 6% decrease in visitation. Allelic variation at the OCIMENE SYNTHASE locus is likely to contribute to differential pollinator visitation, and thus promote reproductive isolation between *M. lewisii* and *M. cardinalis*. OCIMENE SYNTHASE joins a growing list of ‘speciation genes’ (‘barrier genes’) in flowering plants.

Keywords: terpene synthase, floral volatiles, speciation, pollination, *Mimulus lewisii*, *Mimulus cardinalis*.

INTRODUCTION

The rapid diversification of the world’s estimated 275 000 species of flowering plants has often been attributed to their specialized association with different animal pollinators (Grant, 1949; Stebbins, 1970). Flowering plants use a variety of signals to advertise the presence (or illusion) of a reward to their associated pollinators; the association between pollinator type and suites of signals gives rise to the concept of pollination syndromes (Fenster *et al.*, 2004). Perhaps the most well-known and easily studied signal is floral color, which has been investigated in a variety of

pollination syndromes (Rauscher, 2008). Other visual signals, such as texture, pattern, orientation, anthesis time, size, and shape have been investigated to some extent in a variety of systems (Harder and Johnson, 2009; Kay and Sargent, 2009; Yuan *et al.*, 2013a).

Floral scent – the amount, relative ratios, and identities of volatile compounds emitted by the flower – is a generally understudied signal, despite the long understanding that it may play a strong role in attracting pollinators (Raguso, 2008a). The recent development of techniques for

studying floral scent, including chemical analysis of floral scent, analysis of the neural activity of pollinators at both the receptor and higher-order processing levels, and genetic and genomic tools, has allowed some progress in this area. However, although floral scent is frequently characterized, and genes responsible for the production of floral volatiles are occasionally identified, a synthesis of floral scent biochemistry, neurobiology, genetics, ecology, and evolution has been lacking. Those systems with well-characterized volatiles that affect pollination are separate from those with well-characterized genetics (Raguso, 2008a,b; Whitehead and Peakall, 2009; Parachnowitsch *et al.*, 2012).

Much of the work discussing speciation involving floral volatiles has been done in extremely specialized systems where scent is crucial to plant–pollinator interactions (Raguso, 2008b), most notably the sexually deceptive orchids in the genera *Chiloglottis* (Schiestl and Peakall, 2005; Peakall *et al.*, 2010) and *Ophrys* (Schiestl and Ayasse, 2002; Mant *et al.*, 2005; Vereecken *et al.*, 2010; Xu *et al.*, 2012), as well as the non-deceptive genus *Gymnadenia* (Huber *et al.*, 2005). Recent work has begun to expand this to non-orchid systems such as *Silene* (Waelti *et al.*, 2008), *Linanthus* (Chess *et al.*, 2008), and *Petunia* (Klahre *et al.*, 2011). While there are a growing number of studies demonstrating the importance of floral volatiles in mediating these largely specialized plant–pollinator interactions, the genetic pathways controlling volatile production in these systems remain unknown.

In contrast, the genes underlying volatile production are known in a diverse range of angiosperm systems (Gang, 2005), including *Clarkia* (Pichersky *et al.*, 1995; Dudareva *et al.*, 1996, 1998; Wang and Pichersky, 1998), *Antirrhinum* (Dudareva *et al.*, 2000, 2003), *Petunia* (Koeduka *et al.*, 2006; Orlova *et al.*, 2006; Dexter *et al.*, 2007), *Silene* (Gupta *et al.*, 2012), *Arabidopsis* (Bohlmann *et al.*, 2000; Chen *et al.*, 2003), and many species of agricultural importance. Although our knowledge of the genetic underpinnings of volatile production and emission has grown as a result of these systems, there is a paucity of research linking floral volatiles and plant speciation with the genetic and molecular basis for those effects.

Petunia is the only well-developed model demonstrating the role that a specific volatile plays in differential attraction of pollinators between sister species (Klahre *et al.*, 2011). The sister species *Petunia axillaris* and *Petunia exserta* differ in their production of methyl benzoate, a volatile attractive to the hawkmoth pollinators of *P. axillaris*. Through quantitative trait locus (QTL) mapping, two regions underlying this difference were identified in the *Petunia* genome on chromosomes II and VII, with the *P. axillaris* allele at the locus on chromosome II being absolutely required for production of methyl benzoate and the locus on chromosome VII substantially contributing to the quantity of methyl benzoate produced. *ODO1*, one of

the genes hypothesized to underlie the locus on chromosome VII, encodes a MYB transcription factor that is differentially expressed in the two species. The hawkmoth *Manduca sexta* is attracted more strongly to near-isogenic lines with high levels of methyl benzoate production, suggesting that a change in volatile production mediated by a single gene can lead to differential pollinator attraction; however, the underlying genetic mechanisms mediating species-specific volatile emission in this system have not been completely described.

The sister species *Mimulus lewisii* and *Mimulus cardinalis* have served as a model system for studying pollinator-mediated reproductive isolation for several decades (Hiesey *et al.*, 1971), and the combination of ecological and genetic resources has led to the discovery of multiple loci affecting differential pollinator attraction between the two species (Bradshaw and Schemske, 2003; Yuan *et al.*, 2013b,c). *Mimulus lewisii* is a bumblebee-pollinated alpine species, while its sister, *M. cardinalis*, is a lower-elevation hummingbird-pollinated species (Hiesey *et al.*, 1971; Schemske and Bradshaw, 1999). Within areas of sympatry, pollinator fidelity is responsible for 98% of reproductive isolation between *M. lewisii* and *M. cardinalis* (Ramsey *et al.*, 2003). Previous work has shown that three floral volatiles produced by *M. lewisii* – α -limonene, β -myrcene, and *E*- β -ocimene – are important for the attraction of bumblebee pollinators, including *Bombus vosnesenskii*, the native pollinator of *M. lewisii* in the central Sierra Nevada mountains of California. Of the three volatiles, *M. cardinalis* produces only α -limonene, released at just 0.9% of the rate of release in *M. lewisii* (Byers *et al.*, 2014).

Therefore, we ask some global questions, which we begin to address in this paper. What are the genetic underpinnings of the differential emission of floral volatiles between *M. lewisii* and *M. cardinalis*? How many genes are responsible, and how do the species differ in gene expression and protein function? What role, if any, do these scent differences play in differential pollinator visitation, and through this, reproductive isolation? In keeping with other discussions of ‘speciation genes’ in plants (Rieseberg and Blackman, 2010), are these genes of large effect or small effect, structural or regulatory genes? *Mimulus*, with its known attractive volatiles, genetic and genomic tools, and well-studied ecology, is an obvious choice for filling in this missing piece of the floral scent–speciation link.

RESULTS

Genetic mapping of species-specific differences in floral volatiles

Construction of an F₁ cross between *M. lewisii* inbred line LF10 and *M. cardinalis* inbred line CE10 revealed patterns of inheritance of loci controlling the emission of α -limonene,

β -myrcene, and E - β -ocimene. The emission rate of α -limonene for the F_1 plants (mean = 47.2 ng flower⁻¹ h⁻¹; $n = 3$) was similar to that of the *M. lewisii* parental inbred line (mean = 55.1 ng flower⁻¹ h⁻¹, $n = 9$; two-tailed Mann–Whitney $P = 0.864$, $U = 15$) and much higher than that of the *M. cardinalis* parental inbred line (mean = 0.5 ng flower⁻¹ h⁻¹, $n = 9$; one-tailed $P = 0.005$, $U = 27$), suggesting that high levels of emission of α -limonene are inherited from the *M. lewisii* parent in a dominant manner. The pattern was similar for β -myrcene (F_1 , mean = 2.6 ng flower⁻¹ h⁻¹; *M. lewisii*, mean = 3.3 ng flower⁻¹ h⁻¹; *M. cardinalis*, mean = 0.0 ng flower⁻¹ h⁻¹; F_1 versus *M. lewisii* two-tailed $P = 0.600$, $U = 17$; F_1 versus *M. cardinalis* one-tailed $P = 0.005$, $U = 27$). For E - β -ocimene, the *M. lewisii* allele appears to be semi-dominant (F_1 , mean = 2.8 ng flower⁻¹ h⁻¹; *M. lewisii*, mean = 7.6 ng flower⁻¹ h⁻¹; *M. cardinalis*, mean = 0.0 ng flower⁻¹ h⁻¹; F_1 versus *M. lewisii* two-tailed $P = 0.036$, $U = 25$; F_1 versus *M. cardinalis* one-tailed $P = 0.005$, $U = 27$). Complete or partial dominance of the *M. lewisii* alleles for these floral volatiles is consistent with other traits that differ between the species (Bradshaw *et al.*, 1998).

When a backcross ($F_1 \times M. cardinalis$) population of 100 plants was scored for the presence or absence of emission of β -myrcene and E - β -ocimene, it segregated approximately 1:1 for both volatiles (0.52:0.48 β -myrcene present:absent; 0.38:0.62 E - β -ocimene present:absent), suggesting that alleles at Mendelian loci might control the difference in emission of these monoterpenes between *M. lewisii* and *M. cardinalis*. Rates of emission of α -limonene and β -myrcene were very highly correlated ($r = 0.975$), but neither was particularly highly correlated with the rate of emission of E - β -ocimene ($r = 0.474$ versus α -limonene, $r = 0.574$ versus β -myrcene). Therefore, we considered a two-locus model for the difference in these three compounds between the two species – one locus controlling the production of α -limonene and β -myrcene, and another, unlinked, locus controlling E - β -ocimene. A larger backcross population ($n = 768$) was constructed to map the two loci with greater precision.

Identification and characterization of a bifunctional LIMONENE-MYRCENE SYNTHASE (LMS) in *Mimulus lewisii* flowers

The locus associated with α -limonene and β -myrcene emission was mapped to a 15 cM interval between markers M02_510K and M02_1500K (Table S4 in Supporting Information), about 5.3 cM from M02_1500K. Using the assembled and annotated *Mimulus guttatus* genome v1.1 as a reference (<http://www.phytozome.net/cgi-bin/gbrowse/mimulus/>), the ortholog of M02_1500K maps to *M. guttatus* scaffold 89 at position 201 kbp. On scaffold 89 between positions 206 and 226 kbp there is a cluster of three terpene synthases/cyclases – excellent candidates for controlling emission of α -limonene and β -myrcene.

Indel markers developed for two of the *M. lewisii*/*cardinalis* candidate genes in the terpene synthase cluster (Table S5) revealed no recombinations (in 768 backcross plants) between themselves or the putative LIMONENE-MYRCENE SYNTHASE (see Experimental Procedures). The very tight linkage among the candidate terpene synthases within the cluster made it impractical to resolve the identity of the α -limonene and β -myrcene synthases by recombination. Reverse transcriptase (RT)-PCR showed that, of the three candidates, only the *M. lewisii* ortholog (KF857265) of the *M. guttatus* terpene synthase gene on scaffold 89 at position 321 kbp (mgv1a003660m) is transcribed in *M. lewisii* flowers. The marker genotype at M02_1500 accounted for 92% of the difference between emissions of α -limonene in *M. lewisii* and *M. cardinalis* and 98% of the difference in β -myrcene emissions, consistent with a single-locus model for α -limonene and β -myrcene production. No transgressive segregation was observed in the backcross population. The predicted AH150308 gene product contains the conserved DDxx(D/E) and (N,D)Dxx(S,T,G)xxxE (NSE/DTE) motifs required for binding Mg²⁺ during the process of terpene synthesis (Nieuwenhuizen *et al.*, 2013), as well as the RR_xW motif required for cyclic terpene formation (Dudareva *et al.*, 2003).

The *M. lewisii* cDNA (KF857264) orthologous to mgv1a003660m, designated *TS321K*, was overexpressed in *Escherichia coli* (as in Bohlmann *et al.*, 2000). A crude lysate from the *E. coli* culture was supplied with geranyl pyrophosphate (GPP) as a substrate, yielding α -limonene and β -myrcene in the same proportions as observed in the authentic headspace collection from *M. lewisii* flowers (Table S1, Figure 2a). This suggests that the high correlation between α -limonene and β -myrcene emission in the backcross mapping population is due to the pleiotropic effect of a bifunctional LIMONENE-MYRCENE SYNTHASE (LMS) encoded by a single *LMS* gene in *M. lewisii* (*MILMS*). This is consistent with the frequent occurrence of multi-product terpene synthases (Dudareva *et al.*, 2004).

The *M. cardinalis* *LMS* (*McLMS*) coding sequence was not expressed *in vitro* because there is a G66T transversion mutation in exon 3 of *McLMS* (KM659024) that results in a nonsense mutation in the *McLMS* protein (G201X, using *MILMS* AH150308 as the reference allele).

Of note, this is not a definitive demonstration that *LMS* is the gene underlying the locus responsible for the difference in emission of α -limonene and β -myrcene between *M. lewisii* and *M. cardinalis*; it is formally possible that a different, tightly linked gene instead might be responsible. Transgenic complementation of the non-functional *M. cardinalis* *LMS* allele by the *M. lewisii* allele would be necessary to show conclusively that *LMS* is the causal gene for the phenotypic difference.

Identification and characterization of OCIMENE SYNTHASE (OS) in *Mimulus lewisii* flowers

The locus associated with emission of *E*- β -ocimene was mapped to a 7.5-cM interval midway between markers sc4_2325K and M13_2620 (Table S4). The marker genotype at sc4_2325 accounted for 98% of the difference in emissions of *E*- β -ocimene between *M. lewisii* and *M. cardinalis*, consistent with a single-locus model for production of *E*- β -ocimene. No transgressive segregation was observed in the backcross population. The orthologous region of the *M. guttatus* genome lies in a 484-kbp interval (2325–2809 kbp) on scaffold 4. There is a cluster of five terpene biosynthesis genes on scaffold 4 at position 2538–2577 kbp. The gene at position 2538 kbp (mgv1a020487m) is annotated as a terpene synthase, while the other four genes are annotated as sesquiterpene cyclases (Figure 1). An indel marker developed for the *M. lewisii*/*cardinalis* ortholog of mgv1a003660m, designated *TS2538* (Table S5), revealed no recombinations (in 768 backcross plants) with the putative OCIMENE SYNTHASE (see Experimental Procedures).

When overexpressed in *E. coli* and supplied with GPP as a substrate, the *M. lewisii* *TS2538* cDNA (KF857262) encodes a functional OCIMENE SYNTHASE (MIOS, AH150306) (Figure 2a, Table S1). However, under the same conditions the *M. cardinalis* *TS2538* cDNA (KF857263) does not encode an enzyme (McOS, AH150307) capable of synthesizing any monoterpene that we could detect. McOS differs from MIOS at 19 amino acid residues, including

insertion of a leucine residue at position 238 in McOS and deletion of an arginine residue at position 308 in McOS (Figure S1). Both sequences contain the same DDxx(D/E) and NSE/DTE Mg²⁺-binding motifs, as well as the RR_xW cyclase motif, which are unaltered by the 19 non-synonymous amino acid substitutions (Figure S1).

As with *LMS*, in the absence of a transgenic complementation test this is not a strict demonstration that *MIOS* is the gene underlying the locus responsible for the difference in *E*- β -ocimene emission between *M. lewisii* and *M. cardinalis*.

LMS and OS expression *in vivo*

Using RT-PCR with six different stages of flowering tissue from early bud (8 days prior to anthesis, 5 mm) to open flower (see Yuan *et al.*, 2013c), we found that both *MILMS* and *MIOS* are expressed in the last 3 days prior to anthesis (15 and 20 mm) of floral development, as well as in the open flower (Figure 2b). *McLMS* is weakly expressed in late-stage floral buds but not in open flowers. *McOS* is expressed similarly to *MIOS*.

Construction of RNA interference knockdowns of *MILMS* and *MIOS* in stably transformed *Mimulus lewisii*

Using RNA interference (RNAi) via *Agrobacterium tumefaciens*-mediated *in planta* transformation of hairpin RNAi constructs into *M. lewisii* (Yuan *et al.*, 2013c), we were able to knock down the expression of both *MILMS* and *MIOS* to produce much lower levels of floral volatiles, comparable




Figure 1. *Mimulus lewisii* and *Mimulus cardinalis* and their terpene synthases.

(a) *Mimulus lewisii* and *M. cardinalis*.

(b) Genetic maps of *M. lewisii* LIMONENE-MYRCENE SYNTHASE (*MILMS*) and *M. lewisii* OCIMENE SYNTHASE (*MIOS*) and the homologous regions in *Mimulus guttatus*. Positions on the lower half of each are from the *M. guttatus* genome scaffolds; annotations are from queries of the *M. guttatus* transcripts with BLASTx. Putative terpene synthases are highlighted. *M. lewisii* LMS is homologous to a terpene synthase at 319 982 bp on *M. guttatus* scaffold 89, and *M. lewisii* ocimene synthase is homologous to a terpene synthase at 2 538 727 bp on scaffold 4.




Figure 2. Terpene synthase activity *in vitro* and *in vivo*.

(a) Products of terpene synthases using a bacterial overexpression system and *in vitro* enzyme assay. Using geranyl pyrophosphate (GPP, the common monoterpene precursor), the *M. lewisii* LIMONENE-MYRCENE SYNTHASE (MILMS) enzyme produces α -limonene (b) and β -myrcene (a) in the same relative proportion as in the floral volatile emission. *Mimulus lewisii* OCIMENE SYNTHASE (MIOS) produces *E*- β -ocimene (c), but *Mimulus cardinalis* OS does not. All graphs are scaled to the maximum peak in the region, with the exception of *McOS* which is scaled to the maximum peak for *MIOS*.

(b) Temporal expression of terpene synthases *in vivo*. *MILMS* is expressed just prior to flowering and in open flowers, but *McLMS* is not expressed in open flowers. *MIOS* shows a similar expression pattern to *MILMS*, and *McOS* is expressed at the same stages, despite producing no terpenoid volatile that we could detect. Developmental staging is according to bud size of *M. lewisii*, and the corresponding stage of *M. cardinalis* is pictured for *McLMS*, *McOS*, and *McUBC*. UBC, ubiquitin conjugating enzyme.

to those produced in plants homozygous for the *M. cardinalis* alleles at *LMS* and *OS* (Table S2). This allowed us both to verify *LMS* and *OS* gene function *in vivo* and to determine the effect of decreased emission of specific floral volatiles on pollinating bumblebees.

We recovered 24 *M. lewisii* (inbred line LF10) T_1 plants carrying the *MILMS*-RNAi transgene and assayed each transgenic plant in triplicate for production of floral volatiles. All T_1 plants had lower emission rates of α -limonene and β -myrcene relative to the wild-type *M. lewisii* LF10 (α -limonene, range 1.2–56.1%, mean 10.2%; β -myrcene, range 4.1–50.0%, mean 12.8%). Interestingly, most T_1 plants showed a decrease in emission of terpinolene (range 0.0–132.6%, mean 18.0%), indicating that *MILMS* may be responsible for synthesizing an additional minor compound in *M. lewisii*. One of these T_1 transgenics (LMS321K-8) was selfed as the parent of T_2 plants used for pollinator studies (for data on three other T_2 lines from independent T_1 transgenics see Table S3). The original T_1 LMS321K-8 had very low rates of emission of α -limonene and β -myrcene, with a mean production of 2.8% α -limonene and 9.1% β -myrcene relative to the *M. lewisii* LF10 T_0 parent. Notably, LMS321K-8 had an increase in *E*- β -ocimene of 452.8% compared with the *M. lewisii* T_0 parent (Table 1, Figure 3). All other T_1 plants had a similar increase in *E*- β -ocimene production relative to the wild-type parent (range 190.4–493.9%, mean 383.4%).

A total of 71 T_2 plants from the self-pollinated progeny of T_1 LMS321K-8 were assayed using headspace collection of floral volatiles to select the greenhouse population for the bumblebee pollinator behavioral experiment. The 24 individuals selected for the experiment produced much less α -limonene and β -myrcene than the wild-type *M. lewisii* ancestor (α -limonene, range 0.1–2.4%, mean 1.9%; β -myrcene, range 0.0–4.8%, mean 0.6%) and more *E*- β -ocimene than the *M. lewisii* wild-type ancestor (range 93.0–510.5%, mean 247.2%). The levels of α -limonene and β -myrcene were similar to those found in *M. cardinalis* (α -limonene, range 0.03–2.8%, mean 0.9% of wild-type *M. lewisii*; β -myrcene is absent from *M. cardinalis*).

Only two T_1 plants carrying the *MIOS*-RNAi transgene were recovered, but both had the desired *E*- β -ocimene knockdown phenotype relative to *M. lewisii* LF10 (*E*- β -ocimene, range 0.8–2.9%, mean 1.8%; α -limonene, range 39.6–58.9%, mean 49.2%; β -myrcene, range 28.3–41.2%, mean 34.8%). The T_1 plant TS2538-1, which was self-pollinated to create a T_2 population for pollinator studies, had a much lower rate of emission of *E*- β -ocimene (0.8%) relative to *M. lewisii* LF10, as well as lower rates of emission of α -limonene (39.6%) and β -myrcene (28.3%) (Figure 3). The T_1 plant TS2538-2 flowered substantially later than TS2538-1, so T_2 lines were not created from this plant. A total of 80 T_2 plants were produced from TS2538-1, and these produced similar amounts of α -limonene and

Table 1 Volatile production in transgenic (T₁ parent plants of greenhouse experiment lines) and wild-type *Mimulus lewisii* and *Mimulus cardinalis*. Values are an average of two to three independent headspace samples. Emission values in parentheses represent the 90% confidence interval. See 2 for complete data on all volatiles produced by *M. lewisii*

Volatile	LF10 (ng h ⁻¹)	CE10 (ng h ⁻¹)	MILMS-RNAi (ng h ⁻¹)	MIOS-RNAi (ng h ⁻¹)	MILMS-RNAi (%LF10)	MILMS-RNAi (%CE10)	MIOS-RNAi (%LF10)	MIOS-RNAi (%CE10)
β-myrcene	2.837 (2.056, 3.793)	Absent (0.000, 0.000)	0.257 (0.194, 0.339)	1.116 (1.047, 1.185)	9.06	n.a.	39.34	n.a.
D-limonene	43.228 (35.539, 50.708)	1.024 (0.649, 1.419)	1.216 (0.765, 1.757)	23.820 (23.285, 24.354)	2.81	118.75	55.10	2326.17
E-β-ocimene	5.563 (4.320, 7.049)	Absent (0.000, 0.000)	25.125 (20.591, 30.120)	0.054 (0.042, 0.074)	451.64	n.a.	0.97	n.a.

n.a., not applicable.

β-myrcene as the *M. lewisii* LF10 ancestor (D-limonene, range 61.3–127.4%, mean 83.6%; β-myrcene, range 57.3–144.5%, mean 88.7%) but much less E-β-ocimene (range 0.9–3.9%, mean 1.9%).




Figure 3. Results of terpene synthase RNAi knockdown in *M. lewisii*. Stable RNA interference (RNAi) knockdowns of *M. lewisii* LIMONENE-MYRCENE SYNTHASE (MILMS) and *Mimulus lewisii* OCIMENE SYNTHASE (MIOS) in *M. lewisii* produce plants with low emission levels of D-limonene (b)/β-myrcene (a) and E-β-ocimene (c), respectively.

Effects of MILMS and MIOS knockdowns on bumblebee pollinator behavior

Two experiments, one for the MILMS-RNAi transgenics and one for the MIOS-RNAi transgenics, were performed to assay the impact of reduced monoterpene production on bumblebee (*Bombus impatiens*) visitation in a captive greenhouse setting. During each experiment, both preference (expressed as the proportion of total visits to each flower type) and constancy (expressed as the tendency of an individual bumblebee to deviate from random choices, exclusive of preference, see Waser, 1986) were measured.

A total of 1682 visits were observed to flowers in the MILMS-RNAi knockdown experiment. Visits were defined as observable contact with the sexual organs of the flower, i.e. the bumblebee entered the flower fully, as required to effect pollination. Of 1682 visits, 833 (49.52%) were to the wild-type *M. lewisii* and 849 (50.48%) were to the *M. lewisii* MILMS-RNAi transgenic plants, showing no significant difference ($\chi^2 = 0.15$, $P = 0.70$, Figure 4). Bumblebees appeared to show no overall qualitative behavioral difference towards either flower type.

A total of 39 bumblebee foraging bouts were assayed for constancy, with an average Bateman's index of -0.0114 (-1 indicates complete inconstancy – regular switching between types; 0 indicates random visitation patterns; +1 indicates complete constancy, always within types). To determine whether this constancy was significantly different from random visitation, the same bumblebee foraging bouts were used with 100 000 simulated runs of randomly permuted plant locations, resulting in an average Bateman's index of -0.1141. A total of 96 648 simulations had more divergent Bateman's index values than the actual data, showing that bumblebees demonstrated no constancy when presented with these flowers ($P = 0.97$).

For the MIOS-RNAi knockdown experiment, a total of 2202 visits were observed. Of these visits, 1166 (52.95%) were to wild-type *M. lewisii* and 1036 (47.05%) were to the




Figure 4. Greenhouse experiments with *Bombus impatiens* and *Mimulus lewisii* wild-type and transgenic lines.

(a) Image of a typical bumblebee visit.

(b) Response of bumblebees to *M. lewisii* LIMONENE-MYRCENE SYNTHASE (*MILMS*) RNA interference (RNAi) knockdowns, *M. lewisii* OCIMENE SYNTHASE (*MIOS*) RNAi knockdowns, and the wild-type parent. Bumblebees show the same visitation response to *MILMS* knockdown transgenics as to wild-type *M. lewisii*. Bumblebees preferentially visit wild-type *M. lewisii* over *MIOS* knockdown transgenics.

M. lewisii *MIOS*-RNAi plants, showing a significant preference for the wild-type *M. lewisii* flowers ($\chi^2 = 7.67$, $P = 0.0056$, Figure 4). Bumblebees approaching the *MIOS*-RNAi flowers were noted to frequently wave their antennae and contact the flower with their antennae prior to aborting a potential visit, suggesting that *E*- β -ocimene may operate as a near-field olfactory cue, but this behavior was not noted for the wild-type flowers in this experiment nor for either flower type in the *MILMS*-RNAi experiment. Constancy was also absent in the *MIOS*-RNAi experiment, with a total of 46 bumblebee foraging bouts showing an average Bateman's index of 0.0149; the simulation described above was repeated using these foraging bouts, with an average Bateman's index of -0.1142 ($P = 0.95$).

DISCUSSION

Mimulus lewisii produces three floral volatiles with significant neurophysiological and behavioral effects on bumblebees – α -limonene, β -myrcene, and *E*- β -ocimene – while *M. cardinalis* produces only α -limonene at much lower levels (0.9% of *M. lewisii*) (Byers *et al.*, 2014). These differences are probably due to mutations in two genes, LIMONENE-MYRCENE SYNTHASE (*MILMS*, *McLMS*) and OCIMENE SYNTHASE (*MIOS*, *McOS*). In quantitative genetic terms, allelic variation at loci containing *LMS* and *OS* accounts for 92–98% of the phenotypic difference between *M. lewisii* and *M. cardinalis* in floral emission of α -limonene, β -myrcene, and *E*- β -ocimene. The very low level of volatile emission from *M. cardinalis* flowers can be explained at the molecular genetic level; *McLMS* is a null allele due to a nonsense mutation in exon 3, while *McOS* has multiple coding sequence differences that eliminate its ability to produce *E*- β -ocimene.

The RNAi knockouts show that the loss-of-function *LMS* and *OS* alleles can recapitulate the *M. cardinalis* volatile emission phenotypes, and that functional copies of both genes are necessary to produce α -limonene, β -myrcene, and *E*- β -ocimene *in vivo*. However, there remains the formal possibility that the allelic variants producing differences in floral volatile emissions between *M. lewisii* and *M. cardinalis* are not in *LMS* or *OS*, but in genes very tightly linked to them. To show that the *M. lewisii* alleles of *LMS* and *OS* are sufficient (since we have shown that they are necessary) to produce α -limonene, β -myrcene, and *E*- β -ocimene, we would have to transform *M. cardinalis* with constructs containing the *M. lewisii* alleles. However, *M. cardinalis* is very difficult to transform, so we have not performed these definitive experiments. Although we lack conclusive evidence that these are the genes underlying these loci, we present strong circumstantial evidence (the nonsense mutation in *McLMS* and the lack of product from *McOS* activity *in vitro*) that is consistent with this, and RNAi knockouts show that the loss-of-function *LMS* and *OS* alleles can recapitulate the *M. cardinalis* volatile emission phenotypes necessary to test for differential pollinator visitation.

Surprisingly, despite the high level of production of α -limonene and β -myrcene in *M. lewisii* flowers, substantially knocking down emission of these two compounds produces no significant effect on bumblebee visitation in the greenhouse. In contrast, knocking down emission of *E*- β -ocimene results in a modest (6%) but significant decrease in bumblebee visitation, suggesting that alternative alleles of OCIMENE SYNTHASE can contribute to reproductive isolation between the bumblebee-pollinated *M. lewisii* and the hummingbird-pollinated *M. cardinalis*. Although 6% is a modest effect size in molecular genetic terms, in evolutionary genetic terms a selection coefficient (*s*) of 0.06 (130 more visits to the wild-type plant out of

2202 total visits) would sweep the beneficial allele to fixation very quickly in natural populations (Hartl and Clark, 1997), so we designed our pollinator visitation experiments to detect a difference in visitation as small as 5%. Assuming an infinite population size, the probability of fixation of the allele is $2s$, or 12%; an effective population size greater than five individuals would allow selection to exceed drift as an evolutionary force at this locus ($N_e = 1/4s = 4.17$).

Why does the loss of α -limonene and β -myrcene have no effect on bumblebee visitation? First, the T_2 plants used in the greenhouse experiment had surprisingly high levels of E - β -ocimene, perhaps due to re-routing of a common pool of the shared precursor GPP. Terpene synthesis is a flexible but complex process, and buildups of precursors can be utilized by alternate metabolic pathways (Gang, 2005). Given the much higher emission of E - β -ocimene in the *MILMS*-RNAi transgenic plants, *M. lewisii* may be prone to this effect. As the RNAi technique used here is an analogous (but weaker) representation of the phenotypes resulting from a loss-of-function mutation in a wild population, fluctuations in volatile production as found here are reflective of the system's physiology and the effects that might occur in a natural setting.

It is also possible that the high production of α -limonene and β -myrcene in *M. lewisii* serves another function within the plant, such as defense against herbivores, nectar robbers, or disease (Kessler *et al.*, 2013), as these volatiles are known anti-herbivory compounds (Levin, 1976). Although the three volatiles have similar physical properties, α -limonene and β -myrcene may serve to mediate long-distance attraction at the patch level rather than at the level of the individual flower; long-distance attraction has been shown to be important for honeybee (*Apis mellifera*) navigation (Bogdany and Taber, 1979). The high production of α -limonene and β -myrcene may be a remnant of some previous pollination syndrome, environmental context, or merely the byproduct of some other metabolic process within the plant. Similarly, although a significant effect on bumblebee visitation was seen with the loss of E - β -ocimene, it is possible that the main role of this volatile may lie elsewhere (Kessler *et al.*, 2013), for example in herbivory defense (Arimura *et al.*, 2004), with a secondary role in the attraction of bumblebee pollinators. Data on herbivory, florivory, or pathogen infestation in wild populations of *M. lewisii* and *M. cardinalis* are currently lacking, limiting our ability to speculate on these possibilities. Future field experiments will increase our understanding of the multiple roles these volatiles may be playing in *M. lewisii* and *M. cardinalis*.

Finally, it is possible that these effects differ from those that would be found with wild *B. vosnesenskii*. However, both species are generalist floral visitors, and the *M. lewisii* scent elicits similar olfactory responses in both bee species. Moreover, *B. impatiens* has been used as a model for bumblebee–flower interactions in other systems, including

those involving *B. vosnesenskii* (Bodbyl Roels and Kelly, 2011), thus we feel that *B. impatiens* is an excellent model for these experiments (see Appendix S1 for a full explanation). Although these results differ in detail from those we found in previous behavioral experiments with artificial and extracted floral scents, in which all three monoterpenes were required for maximum bumblebee response (Byers *et al.*, 2014), the greenhouse experiments offer a more realistic assay for the effect of scent on pollinators by allowing them to integrate multiple floral cues.

What role does scent play in pollinator interaction within this system? Many studies have shown that scent plays a strong role in landing decisions by diurnal pollinators such as bumblebees and honeybees (Butler, 1951; Galen and Kevan, 1980, 1983; Lunau, 1992; Majetic *et al.*, 2009; Dötterl and Vereecken, 2010) – the initial approach may be guided by patch-level visual signals, followed by a visually guided approach to an individual flower. At that point, the final landing decision may be influenced by floral scent, especially in relatively weakly scented flowers such as *M. lewisii* (Dötterl and Vereecken, 2010; Parachnowitsch *et al.*, 2012). Therefore, even in the densely flowered greenhouse experiments, signals such as the presence or absence of E - β -ocimene may play a significant role in final landing decisions. Additionally, densities in the greenhouse experiments were similar to those found in wild populations of *M. lewisii*, which grows along montane streambeds in large clusters, so the dense greenhouse conditions are a better indicator of the potential effect of a single change in scent in a wild population.

How might a loss-of-function allele of *OS* promote a pollinator switch from bumblebees to hummingbirds? Hummingbirds have a very limited sense of smell (Ioalé and Papi, 1989) and retain scent information very poorly (Goldsmith and Goldsmith, 1982), so the loss of scent in a hummingbird-pollinated flower such as *M. cardinalis* (an 'anti-bee' but not 'pro-bird' shift, to use the language in Castellanos *et al.*, 2004) would be likely to have no fitness cost, and might even increase fitness by discouraging bumblebee visitors from transferring heterospecific pollen to the stigma and carrying away nectar or pollen. In combination with the difference in visual signals and mechanical access found between *M. lewisii* and *M. cardinalis*, such a loss of E - β -ocimene might serve to reinforce visitation behavior. Whether these changes in floral volatiles evolved in allopatry or as reinforcement during secondary contact is unclear; investigating the volatile profiles and orthologous terpene synthase genes of other species in *Mimulus* section *Erythranthe* may provide some insight into this question.

The fact that the *OS* polymorphism between *M. lewisii* and *M. cardinalis* is in a structural gene contradicts the current thinking that genes involved in pre-zygotic reproductive isolation – often referred to as 'speciation genes' (Coyne, 1992) or 'barrier genes' (Noor and Feder, 2006) – are nearly

always regulatory genes (Rieseberg and Blackman, 2010). However, the limited number of genes with known effects in pre-zygotic reproductive isolation should preclude any general conclusions from being drawn about this process. We would suggest, however, that the potential for structural genes to contribute to this process should not be ignored.

Although many systems used in the study of floral volatiles have relatively strong scents that are detectable by the human nose, scent can also be a factor in reproductive isolation in systems where it is easily missed, as in *Mimulus*. The role of strong emissions of floral volatiles in attracting nighttime pollinators from a distance is well documented (Raguso and Willis, 2003). The potential role of changes in floral scent in pollinator-mediated reproductive isolation involving generalist, daytime pollinators such as bumblebees is largely unknown, and no examples integrating floral scent genetics and pollinator reproductive isolation in sister species with generalist pollinators have been reported. Some authors have commented that the role of floral scent in reproductive isolation is questionable in generalist cases, as floral scents thus serve less as 'private channels' and pollinators are attracted to multiple floral scent profiles (Schiestl and Ayasse, 2002).

Floral scent should be considered as an attractive factor even in generalist systems, along with more easily measured visual signals such as floral color and pattern. Here, the sister species *M. lewisii* and *M. cardinalis* can be used as a model for the study of reproductive isolation involving floral volatiles – one can begin by looking at species-specific differences, identifying critical volatiles within a complex mixture via electrophysiological and behavioral assays (Riffell *et al.*, 2013; Byers *et al.*, 2014). Then, studies can proceed by determining the genetic basis of these phenotypic differences, creating high-resolution genetic materials (near-isogenic lines, transgenics), and, finally, assaying of the results of these genetic changes in ecologically relevant greenhouse or field settings. Nearly all previous studies of the role of floral volatiles in speciation have only answered a subset of these questions, but our work with *Mimulus*, an emerging model system, shows that a comprehensive, integrative study is possible.

EXPERIMENTAL PROCEDURES

Quantitative trait loci and fine mapping

Volatiles were first assayed in triplicate in an F₁ cross of *M. lewisii* inbred line LF10 and *M. cardinalis* inbred line CE10 (LF10 × CE10) and compared with previous results for the parent lines (nine samples each; see Byers *et al.*, 2014) using a Mann–Whitney *U*-test. A coarse mapping population consisting of 100 individuals of a cross between LF10 and CE10, backcrossed to CE10 [(LF10 × CE10) × CE10], was then constructed. Headspace volatiles were collected in the manner described in Byers *et al.* (2014) (see also Appendix S1 and below) and assayed for emission rates of α -limonene, β -myrcene, and *E*- β -ocimene. Pearson correlation

coefficients were calculated pairwise for the three scents to investigate potential linkage or pleiotropy. A subset of 24 backcross plants with the two most divergent phenotypes (high α -limonene/ β -myrcene and low *E*- β -ocimene; low α -limonene/ β -myrcene and high *E*- β -ocimene) were screened at 34 indel markers evenly spaced across the genome (Table S4) with the intent of creating a low-resolution QTL map. However, it was clear from inspection of the genotypic and phenotypic data that the emission of α -limonene/ β -myrcene and *E*- β -ocimene were, to a first approximation, segregating as Mendelian traits.

A larger backcross population ($n = 768$) was constructed and screened with markers flanking the putative *LIMONENE-MYRCENE SYNTHASE* (M02_510 and M02_1500) and the putative *OCIMENE SYNTHASE* (sc4_2325K and M13_2620) (Table S4). Markers used in the mapping process were developed from *M. lewisii* and *M. cardinalis* genome sequences, and amplify co-dominant markers in the backcross. To reduce the effort required to score flowers for volatile production, only those backcross plants with informative recombinations between markers flanking *LMS* ($n = 107$) or *OS* ($n = 52$) were phenotyped for scent, using a direct extraction assay from flowers rather than the more labor- and time-intensive headspace collection method. For further details, see Appendix S1.

The *Mimulus guttatus* genomic region corresponding to the *M. lewisii* region containing *LMS* or *OS* was examined, and candidate genes were identified based upon their map position relative to the flanking molecular markers and the *M. guttatus* annotation. For the *LMS* locus controlling emission of α -limonene and β -myrcene, primers were designed to amplify indel polymorphisms in two of the terpene synthases/cyclases on *M. guttatus* scaffold 89. The candidate genes were designated LC250K and TS306K (see Table S5 for all primers). No recombination events were observed among the two candidate genes and the putative *LMS*, defining a candidate region of <0.1 cM. For the *OS* locus controlling emission of *E*- β -ocimene, primers were designed to amplify an indel polymorphism in a terpene synthase designated TS2538 on *M. guttatus* scaffold 4. No recombination events were observed between TS2538 and *OS*, defining a candidate region of <0.1 cM.

In vitro assay for terpene synthase activity

For details see Appendix S1 and Fäldt *et al.* (2003).

Terpene synthase expression *in vivo*

Total RNA was extracted from flower buds collected at 5-, 8-, 10-, 15-, and 20-mm stages in *M. lewisii* and equivalent stages in *M. cardinalis*. Total RNA was extracted from open flowers of both species. Complementary DNA was prepared from total RNA extracts using the SuperScript III First-Strand Synthesis System (Invitrogen, <http://www.invitrogen.com/>). Reverse transcriptase-PCR for both loci for both species was performed with these cDNA, using *MIUBC* as a control for background expression levels as described in Yuan *et al.* (2013c). The following primers were used: *MILMS* RNAi forward/reverse for *MILMS*, *McLMS* forward/reverse for *McLMS*; *MIOS* RNAi sense forward/reverse for *MIOS* and *McOS*; and *MIUBC* forward/reverse for *MIUBC*.

Construction of *MILMS* and *MIOS* RNAi transgenic *Mimulus lewisii*

Transgenesis was done in the *M. lewisii* background, as insect pollination is inferred to be the ancestral state in this clade (Beardsley *et al.*, 2003). Hairpin RNAi transgenes targeted to knock down the expression of *MILMS* or *MIOS* were constructed in pFGC5941

(Kerschen *et al.*, 2004; Arabidopsis Biological Resource Center, CD3-447) as described in Yuan *et al.* (2013c). In each case, target specificity of the RNAi fragment was assured by BLAST search against the *M. lewisii* LF10 genome sequence. For *MILMS*, a 106-bp fragment of *M. lewisii* cDNA was amplified and directionally cloned into the pFGC5941 *NcoI/Ascl* (sense) and *BamHI/XbaI* (antisense) sites. For *MIOS*, a 289-bp sense fragment was amplified and directionally cloned into the *NcoI/Ascl* site of pFGC5941. A 180-bp antisense fragment (entirely within the 289-bp *NcoI/Ascl* amplicon) was amplified and directionally cloned into the *BamHI/XbaI* site. Constructs were verified by sequencing, then electroporated separately into *A. tumefaciens* strain GV3101 and used for *in planta* transformation of *M. lewisii* LF10 following Yuan *et al.* (2013c).

Greenhouse experiments

For details of experimental design see Appendix S1. The layout of the greenhouse experiments, including randomized plant positions, is depicted in Figure S2.

Observations of bumblebee behavior were recorded for the first 6 h of the first 3 days by two observers using voice recorders, each following one or two bumblebees at a time. At the start of each day prior to the first bumblebee activity, old flowers were removed and newly opened flowers were counted and equalized between the two types of plants (wild type and RNAi transgenic) to ensure that bumblebees had an equal chance of encountering a given flower of each type on each day.

Data were transcribed and analyzed for preference (proportion of total visits) and constancy. For constancy, visits were 'collapsed' to the plant level, i.e. multiple visits to one plant in sequence were reduced to a single visit, since flower numbers were unequal between plants and flowers were often tightly clustered. Bumblebees were only used for constancy analysis if they visited 10 or more plants in a foraging bout. Constancy was calculated using Bateman's method (described in Waser, 1986), which is independent of preference; equalizing flowers at the start of the day gave pollinators equal access to each type, as required by this metric. To determine if observed constancy was different from the null expectation, the same foraging data were used in a permutation test with shuffled plant identities, repeated 100 000 times; the fraction of the simulations with a greater than observed deviation from zero (complete randomness) was used to estimate the *P*-value.

ACKNOWLEDGEMENTS

We thank D. Ewing, P. Beeman, E. Forbush, J. Milne, N. Kurashige, M. Kovic, and B. Watson for greenhouse plant care, and M. Sargent for assistance in greenhouse experiments. The manuscript was improved greatly by helpful and thorough critiques from D. Tholl and two anonymous referees. This work was supported by a National Institutes of Health grant to HDB (5R01GM088805), National Science Foundation grants to HDB (FIBR 0328636; IOS 1209340), JAR (IOS 0822709; DBI 1121692), and KJRPB (IOS 1209340), and a NSF Graduate Research Fellowship to KJRPB (DGE 0718124). KJRPB is also supported by an ARCS Foundation Fellowship from the Seattle Chapter of the Achievement Rewards for College Scientists Foundation and the Melinda Denton Writing Fellowship.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. Protein sequences of MIOS and McOS with differences highlighted.

Figure S2. Schematic of the experimental greenhouse setup.

Table S1. Products of the *in vitro* terpene synthase assays.

Table S2. Volatile production in the two to four best T₁ plants recovered from RNA interference experiments.

Table S3. Volatile production in T₂ plants from four separate T₁ parents recovered from RNA interference knockdown of *MILMS*.

Table S4. Molecular markers used during quantitative trait locus and fine mapping.

Table S5. Sequences of oligonucleotide primers used.

Appendix S1. Description of volatile headspace collection and analysis, volatile extraction collection and scoring, *in vitro* assays for terpene synthase activity, and greenhouse experiments.

REFERENCES

- Arimura, G., Ozawa, R., Kugimiya, S., Takabayashi, J. and Bohlmann, J. (2004) Herbivore-induced defense response in a model legume. Two-spotted spider mites induce emission of (E)- β -ocimene and transcript accumulation of (E)- β -ocimene synthase in *Lotus japonicas*. *Plant Physiol.* **135**(4), 1976–1983.
- Beardsley, P.M., Yen, A. and Olmstead, R.G. (2003) AFLP phylogeny of *Mimulus* section *Erythranthe* and the evolution of hummingbird pollination. *Evolution*, **57**(6), 1397–1410.
- Bodbyl Roels, S.A. and Kelly, J.K. (2011) Rapid evolution caused by pollinator loss in *Mimulus guttatus*. *Evolution*, **65**(9), 2541–2552.
- Bogdany, F.J. and Taber, S. III (1979) The significance of odor for bees orienting across a canyon. *Apidologie*, **10**(1), 55–62.
- Bohlmann, J., Martin, D., Oldham, N.J. and Gershenzon, J. (2000) Terpenoid secondary metabolism in *Arabidopsis thaliana*: cDNA cloning, characterization, and functional expression of a myrcene/(E)- β -ocimene synthase. *Arch. Biochem. Biophys.* **375**(2), 261–269.
- Bradshaw, H.D. Jr and Schemske, D.W. (2003) Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature*, **426**, 176–178.
- Bradshaw, H.D. Jr, Otto, K.G., Frewen, B.E., McKay, J.K. and Schemske, D.W. (1998) Quantitative trait loci affecting differences in floral morphology between two species of monkeyflower (*Mimulus*). *Genetics*, **149**, 367–382.
- Butler, C.G. (1951) The importance of perfume in the discovery of food by the worker honeybee (*Apis mellifera* L.). *Proc. R. Soc. Lond. B Biol. Sci.* **138**, 403–413.
- Byers, K.J.R.P., Bradshaw, H.D. Jr and Riffell, J.A. (2014) Three floral volatiles contribute to differential pollinator attraction in monkeyflowers (*Mimulus*). *J. Exp. Biol.* **217**(4), 614–623.
- Castellanos, M.C., Wilson, P. and Thomson, J.D. (2004) 'Anti-bee' and 'pro-bird' changes during the evolution of hummingbird pollination in *Penstemon* flowers. *J. Evol. Biol.* **17**(4), 876–885.
- Chen, F., Tholl, D., D'Auria, J.C., Farooq, A., Pichersky, E. and Gershenzon, J. (2003) Biosynthesis and emission of terpenoid volatiles from *Arabidopsis* flowers. *Plant Cell*, **15**, 481–494.
- Chess, S.K.R., Raguso, R.A. and LeBuhn, G. (2008) Geographic divergence in floral morphology and scent in *Linanthus dichotomus* (Polemoniaceae). *Am. J. Bot.* **95**(12), 1652–1659.
- Coyne, J.A. (1992) Genetics and speciation. *Nature*, **355**, 511–515.
- Dexter, R., Qualley, A., Kish, C.M., Ma, C.J., Koeduka, T., Nagegowda, D.A., Dudareva, N., Pichersky, E. and Clark, D. (2007) Characterization of a petunia acetyltransferase involved in the biosynthesis of the floral volatile iso Eugenol. *Plant J.* **49**(2), 265–275.
- Dötterl, S. and Vereecken, N.J. (2010) The chemical ecology and evolution of bee-flower interactions: a review and perspectives. *Can. J. Zool.* **88**, 668–697.
- Dudareva, N., Cseke, L., Blanc, V.M. and Pichersky, E. (1996) Evolution of floral scent in *Clarkia*: novel patterns of S-linalool synthase gene expression in the *C. breweri* flower. *Plant Cell*, **8**, 1137–1148.
- Dudareva, N., D'Auria, J.C., Nam, K.H., Raguso, R.A. and Pichersky, E. (1998) Acetyl-CoA:benzylalcohol acetyltransferase – an enzyme involved in floral scent production in *Clarkia breweri*. *Plant J.* **14**(3), 297–304.

- Dudareva, N., Murfitt, L.M., Mann, C.J., Gorenstein, N., Kolosova, N., Kish, C.M., Bohman, C. and Wood, K. (2000) Developmental regulation of methyl benzoate biosynthesis and emission in snapdragon flowers. *Plant Cell*, **12**(6), 949–961.
- Dudareva, N., Martin, D., Kish, C.M., Kolosova, N., Gorenstein, N., Fäldt, J., Miller, B. and Bohlmann, J. (2003) (*E*)- β -Ocimene and myrcene synthase genes of floral scent biosynthesis in snapdragon: function and expression of three terpene synthase genes of a new terpene synthase subfamily. *Plant Cell*, **15**(5), 1227–1241.
- Dudareva, N., Pichersky, E. and Gershenzon, J. (2004) Biochemistry of plant volatiles. *Plant Physiol.* **135**, 1893–1902.
- Fäldt, J., Martin, D., Miller, B., Rawat, S. and Bohlmann, J. (2003) Traumatic resin defense in Norway spruce (*Picea abies*): methyl jasmonate-induced terpene synthase gene expression, and cDNA cloning and functional characterization of (+)-3-carene synthase. *Plant Mol. Biol.* **51**, 119–133.
- Fenster, C.B., Armbruster, W.S., Wilson, P., Dudash, M.R. and Thomson, J.D. (2004) Pollination syndromes and floral specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**, 375–403.
- Galen, C. and Kevan, P.G. (1980) Scent and color, floral polymorphisms and pollination biology in *Polemonium viscosum* Nutt. *Am. Midl. Nat.* **104**(2), 281–289.
- Galen, C. and Kevan, P.G. (1983) Bumblebee foraging and floral scent dimorphism: *Bombus kirbyellus* Curtis (Hymenoptera:Apidae) and *Polemonium viscosum* Nutt. (Polemoniaceae). *Can. J. Zool.* **61**, 1207–1213.
- Gang, D.R. (2005) Evolution of flavors and scents. *Annu. Rev. Plant Biol.* **56**, 301–325.
- Goldsmith, K.M. and Goldsmith, T.H. (1982) Sense of smell in the Black-chinned Hummingbird. *Condor*, **84**, 237–238.
- Grant, V. (1949) Pollination systems as isolating mechanisms in angiosperms. *Evolution*, **3**, 82–97.
- Gupta, A.K., Akhtar, T., Widmer, A., Pichersky, E. and Schiestl, F.P. (2012) Identification of white campion (*Silene latifolia*) guaiacol O-methyltransferase involved in the biosynthesis of veratrole, a key volatile for pollinator attraction. *BMC Plant Biol.* **12**, 158.
- Harder, L.D. and Johnson, S.D. (2009) Darwin's beautiful contrivances: evolutionary and functional evidence for floral adaptation. *New Phytol.* **183**(3), 530–545.
- Hartl, D.L. and Clark, A.G. (1997) *Principles of Population Genetics*, 4th edn. Sunderland, MA: Sinauer Associates Inc.
- Hiesey, W.M., Nobs, M.A. and Björkman, O. (1971) *Experimental Studies on the Nature of Species. V. Biosystematics, Genetics, and Physiological Ecology of the Erythranthe Section of Mimulus*. Washington, DC: Carnegie Inst. Washington Publ. 628, pp. 1–213.
- Huber, F.K., Kaiser, R., Sauter, W. and Schiestl, F.P. (2005) Floral scent emission and pollinator attraction in two species of *Gymnadenia* (Orchidaceae). *Oecologia*, **142**(4), 564–575.
- Ioalé, P. and Papi, F. (1989) Olfactory bulb size, odor discrimination and magnetic insensitivity in hummingbirds. *Physiol. Behav.* **45**, 995–999.
- Kay, K.M. and Sargent, R.D. (2009) The role of animal pollination in plant speciation: integrating ecology, geography, and genetics. *Annu. Rev. Ecol. Evol. Syst.* **40**, 637–656.
- Kerschen, A., Napoli, C.A., Jorgensen, R.A. and Müller, A.E. (2004) Effectiveness of RNA interference in transgenic plants. *FEBS Lett.* **566**, 223–228.
- Kessler, D., Diezel, C., Clark, D.G., Colquhoun, T.A. and Baldwin, I.T. (2013) Petunia flowers solve the defence/apparency dilemma of pollinator attraction by deploying complex floral blends. *Ecol. Lett.* **16**(3), 299–306.
- Klahre, U., Gurba, A., Hermann, K., Saxenhofer, M., Bossolini, E., Guerin, P.M. and Kuhlmeier, C. (2011) Pollinator choice in *Petunia* depends on two major genetic loci for floral scent production. *Curr. Biol.* **21**, 730–739.
- Koeduka, T., Fridman, E., Gang, D.R. et al. (2006) Eugenol and isoeugenol, characteristic aromatic constituents of spices, are biosynthesized via reduction of a coniferyl alcohol ester. *Proc. Natl Acad. Sci. USA*, **103**(26), 10128–10133.
- Levin, D.A. (1976) The chemical defenses of plants to pathogens and herbivores. *Annu. Rev. Ecol. Syst.* **7**, 121–159.
- Lunau, K. (1992) Innate recognition of flowers by bumble bees: orientation of antennae to visual stamen signals. *Can. J. Zool.* **70**, 2139–2144.
- Majetic, C.J., Raguso, R.A. and Ashman, T.-L. (2009) The sweet smell of success: floral scent affects pollinator attraction and seed fitness in *Hesperis matronalis*. *Funct. Ecol.* **23**, 480–487.
- Mant, J., Peakall, R. and Schiestl, F.P. (2005) Does selection on floral odor promote differentiation among populations and species of the sexually deceptive orchid genus *Ophrys*? *Evolution*, **59**(7), 1449–1463.
- Nieuwenhuizen, N.J., Green, S.A., Chen, X., Bailleul, E.J.D., Matich, A.J., Wang, M.Y. and Atkinson, R.G. (2013) Functional genomics reveals that a compact terpene synthase gene family can account for terpene volatile production in apple. *Plant Physiol.* **161**, 787–804.
- Noor, M.A.F. and Feder, J.L. (2006) Speciation genetics: evolving approaches. *Nat. Rev. Genet.* **7**, 851–861.
- Orlova, I., Marshall-Colón, A., Schnepf, J. et al. (2006) Reduction of benzenoid synthesis in petunia flowers reveals multiple pathways to benzoic acid and enhancement in auxin transport. *Plant Cell*, **18**(12), 3458–3475.
- Parachnowitsch, A.L., Raguso, R.A. and Kessler, A. (2012) Phenotypic selection to increase floral scent emission, but not flower size or colour in bee-pollinated *Penstemon digitalis*. *New Phytol.* **195**, 667–675.
- Peakall, R., Ebert, D., Poldy, J., Barrow, R.A., Francke, W., Bower, C.C. and Schiestl, F.P. (2010) Pollinator specificity, floral odour chemistry and the phylogeny of Australian sexually deceptive *Chiloglottis* orchids: implications for pollinator driven speciation. *New Phytol.* **188**(2), 437–450.
- Pichersky, E., Lewinsohn, E. and Croteau, R. (1995) Purification and characterization of S-linalool synthase, an enzyme involved in the production of floral scent in *Clarkia breweri*. *Arch. Biochem. Biophys.* **316**(2), 803–807.
- Raguso, R.A. (2008a) Start making scents: the challenge of integrating chemistry into pollination ecology. *Entomol. Exp. Appl.* **128**(1), 196–207.
- Raguso, R.A. (2008b) Wake up and smell the roses: the ecology and evolution of floral scent. *Annu. Rev. Ecol. Evol. Syst.* **39**, 549–569.
- Raguso, R.A., Willis, M.A. (2003) Hawkmoth pollination in Arizona's Sonoran Desert: behavioral responses to floral traits. In *Evolution and Ecology Taking Flight, Butterflies as Model Systems* (Boggs, C.L., Watt, W. B., and Ehrlich, P. R., eds). Chicago: University of Chicago Press, pp. 43–65.
- Ramsey, J., Bradshaw, H.D. Jr and Schemske, D.W. (2003) Components of reproductive isolation between the monkeyflowers *Mimulus lewisii* and *M. cardinalis* (Phrymaceae). *Evolution*, **57**(7), 1520–1534.
- Rausher, M.D. (2008) Evolutionary transitions in floral color. *Int. J. Plant Sci.* **169**(1), 7–21.
- Rieseberg, L.H. and Blackman, B.K. (2010) Speciation genes in plants. *Ann. Bot.* **106**(3), 439–455.
- Riffell, J.A., Lei, H., Abrell, L. and Hildebrand, J.G. (2013) Neural basis of a pollinator's buffet: olfactory specialization and learning in *Manduca sexta*. *Science*, **339**(6116), 200–204.
- Schemske, D.W. and Bradshaw, H.D. Jr (1999) Pollinator preference and the evolution of floral traits in monkeyflowers (*Mimulus*). *Proc. Natl Acad. Sci. USA*, **96**(21), 11910–11915.
- Schiestl, F.P. and Ayasse, M. (2002) Do changes in floral odor cause speciation in sexually deceptive orchids? *Plant Syst. Evol.* **234**, 111–119.
- Schiestl, F.P. and Peakall, R. (2005) Two orchids attract different pollinators with the same floral odour compound: ecological and evolutionary implications. *Funct. Ecol.* **19**(4), 674–680.
- Stebbins, G.L. (1970) Adaptive radiation of reproductive characteristics in angiosperms, I, pollination mechanisms. *Annu. Rev. Ecol. Syst.* **1**, 307–326.
- Vereecken, N.J., Cozzolino, S. and Schiestl, F.P. (2010) Hybrid floral scent novelty drives pollinator shift in sexually deceptive orchids. *BMC Evol. Biol.* **10**(1), 103.
- Waelti, M.O., Muhlemann, J.K., Widmer, A. and Schiestl, F.P. (2008) Floral odour and reproductive isolation in two species of *Silene*. *J. Evol. Biol.* **21**(1), 111–121.
- Wang, J. and Pichersky, E. (1998) Characterization of S-adenosyl-L-methionine:(isoeugenol O-methyltransferase involved in floral scent production in *Clarkia breweri*. *Arch. Biochem. Biophys.* **349**(1), 153–160.
- Waser, N.M. (1986) Flower constancy: definition, cause, and measurement. *Am. Nat.* **127**(5), 593–603.
- Whitehead, M.R. and Peakall, R. (2009) Integrating floral scent, pollination ecology and population genetics. *Funct. Ecol.* **23**, 863–874.
- Xu, S., Schlüter, P.M., Grossniklaus, U. and Schiestl, F.P. (2012) The genetic basis of pollinator adaptation in a sexually deceptive orchid. *PLoS Genet.* **8**(8), e1002889.

Yuan, Y.-W., Byers, K.J.R.P. and Bradshaw, H.D. Jr (2013a) The genetic control of flower-pollinator specificity. *Curr. Opin. Plant Biol.* **16**(4), 422–428.

Yuan, Y.-W., Sagawa, J.M., Di Stilio, V.S. and Bradshaw, H.D. Jr (2013b) Bulk segregant analysis of an induced floral mutant identifies a *MIX-TA*-like R2R3 *MYB* controlling nectar guide formation in *Mimulus lewisii*. *Genetics*, **194**(2), 523–528.

Yuan, Y.-W., Sagawa, J.M., Young, R.C., Christensen, B.J. and Bradshaw, H.D. Jr (2013c) Genetic dissection of a major anthocyanin QTL contributing to pollinator-mediated reproductive isolation between sister species of *Mimulus*. *Genetics*, **194**(1), 255–263.