

vation process, on a molecular layer basis, holds the possibility for many exciting future experiments. An obvious one is to probe how electric fields alter the ion's ability to penetrate the water-oil interfacial barrier. Another possibility takes advantage of the much faster net motion of ions relative to neutrals at these field strengths: We can epitaxially create structured liquids and measure ion motion or trapping in them. In this way we can closely mimic complex interfaces in electrochemistry and in cell membranes.

References and Notes

1. D. A. Doyle *et al.*, *Science* **280**, 69 (1998).
2. R. K. Sen, E. Yeager, W. E. O'Grady, *Annu. Rev. Phys. Chem.* **26**, 287 (1975).
3. C. M. Starks, C. L. Liotta, M. Halpern, *Phase-Transfer Catalysis—Fundamentals, Applications and Industrial Perspectives* (Chapman & Hall, New York, 1994).

4. W. Mitchell Jr., *Fuel Cells* (Academic Press, New York, 1963).
5. R. Noyes, *Nuclear Waste Cleanup Technology and Opportunities* (Noyes, Park Ridge, NJ, 1995).
6. H. E. Allen, *Metal Contaminated Aquatic Sediments* (Ann Arbor Press, Chelsea, MI, 1995).
7. M. Born, *Z. Phys.* **1**, 45 (1920).
8. E. Conway, *J. Electroanal. Chem.* **65**, 491 (1975).
9. I. Benjamin, *Annu. Rev. Phys. Chem.* **48**, 407 (1997).
10.  $D_3O^+$  motion follows the same pattern in 3MP films regardless of the temperature (30 to 125 K) at which they are annealed or grown.
11. A. C. Ling and J. E. Willard, *J. Phys. Chem.* **72**, 1918 (1968).
12. J. P. Cowin, A. A. Tsekouras, M. J. Iedema, K. Wu, G. B. Ellison, *Nature* **398**, 405 (1999).
13. J. P. Biessecker *et al.*, *Rev. Sci. Instrum.* **69**, 485 (1998).
14. A complete, bulk ice-like layer of water has molecules in two slightly separated planes and is traditionally termed a "bilayer" rather than a monolayer. See N. Materer *et al.*, *Surf. Sci.* **381**, 190 (1997).
15. D. R. Lide, *CRC Handbook of Chemistry and Physics* (CRC Press, Boca Raton, FL, 1997), pp. 6–104.
16. M. J. Iedema *et al.*, *J. Phys. Chem. B* **103**, 9203 (1998).

17. R. S. Smith, C. Huang, E. K. L. Wong, B. D. Kay, *Phys. Rev. Lett.* **79**, 909 (1997).
18. A. A. Tsekouras, M. J. Iedema, J. P. Cowin, *J. Chem. Phys.* **111**, 2222 (1999).
19. L. X. Dang, *J. Phys. Chem.* **103**, 8195 (1999).
20. I. Benjamin, *Science* **261**, 1558 (1993).
21. The  $V_f$  versus  $T$  curve for  $D_3O^+$  on thick amorphous water-ice film (>50 bilayers) grown at 30 K showed that the ions did not move until 150 K, although  $V_f$  did drop below 120 K because of the turning-on of the dielectric constant of the amorphous ice. See, for example, A. A. Tsekouras, M. J. Iedema, J. P. Cowin, *Phys. Rev. Lett.* **80**, 5798 (1998).
22. Supported by the U.S. Department of Energy (DOE) Basic Energy Sciences, and performed at the Wiley Environmental Molecular Sciences Laboratory, sponsored by the DOE Office of Biological and Environmental Research. Pacific Northwest National Laboratory is operated by Battelle (contract DE-AC06-76RLO 1830). We thank A. A. Tsekouras for assistance in the early stages of the experiment. K.W. dedicates this paper to Professor Xiexian Guo (1925–1998).

9 September 1999; accepted 9 November 1999

# Oligotrophy and Nitrogen Fixation During Eastern Mediterranean Sapropel Events

Julian P. Sachs<sup>1\*</sup> and Daniel J. Repeta<sup>2</sup>

Nitrogen isotopic measurements in fossil chlorophyll from late Pleistocene organic-rich sediments (sapropels) in the eastern Mediterranean Sea provide geochemical evidence for stratified, nutrient-depleted surface water and extensive nitrogen fixation. This evidence is reconciled with previous indications of high productivity by invoking a model of sapropel formation in which increased river discharge facilitates development of a specialized phytoplankton population whose annual mass sinking provides the organic flux to generate sapropels. This interpretation is consistent with the widespread occurrence of mat-forming diatoms that thrive in stratified water and can harbor diazotrophic bacterial symbionts, but does not support eutrophication of surface waters by enhanced river runoff or a circulation reversal.

The eastern Mediterranean Sea is a well-ventilated, nutrient-depleted basin (1) characterized by low primary productivity (2) and organic-deficient sediments (3). However, during the late Pleistocene, a series of organic-rich sequences, sapropels, were deposited under what must have been dramatically different depositional conditions than those of today (4). These green-brown to black deposits with organic carbon concentrations of 2 to 5% are interspersed between gray nannofossil and foraminiferal marl oozes with organic carbon concentrations of 0.1 to 0.3% (3, 5). They are typically 1 to 30 cm thick (4), were formed over periods of 1000 to 10,000 years, and appear to be basin-wide events at water depths below 300 m (6). Seven

sapropels, numbered S7 through S1, were deposited during the last 200,000 years, the most recent of which occurred in the early Holocene (4). The cause of these sedimentary layers is most likely enhanced primary productivity and fluxes of organic carbon to the seafloor (3, 7, 8), improved preservation rates of organic matter in oxygen-deficient water (9), or a combination of the two (6, 10). We conclude from nitrogen isotopic ratios in chlorophyll derivatives (chlorins) that surface waters were oligotrophic, deep waters were anoxic, and nitrogen fixation was widespread during these events. By invoking micropaleontological-based models of sapropel formation (6, 7, 11) in which increased river runoff induces shoaling of intermediate waters, the formation of a deep algal community, and diminished deep-water ventilation, we reconcile these findings with existing evidence for high export production.

Large fluctuations in sedimentary nitrogen isotopic ratios occur when sapropel layers are present. Whole-sediment  $\delta^{15}N$  (12) values within sapropels S2 through S7 are nearly con-

stant at  $-0.1 \pm 0.5\text{‰}$  ( $\pm 1\sigma$ ;  $n = 26$ ) and significantly lower than non-sapropel values between those events (Fig. 1, B and C) that average  $5.3 \pm 1.0\text{‰}$  ( $n = 27$ ), irrespective of sample age or core location. There are two fundamentally different interpretations of these data. One is that the non-sapropel  $\delta^{15}N$  value of  $5.3\text{‰}$  represents the sinking flux of nitrogen, or new production, under normal, oligotrophic conditions, and the sapropel  $\delta^{15}N$  value of  $-0.1\text{‰}$  indicates new production from nutrient-replete surface waters (5). As demonstrated in a variety of oceanographic settings, high nutrient concentrations in surface waters lead to the production of organic matter with low  $\delta^{15}N$  values (13) because faster uptake kinetics cause preferential assimilation of  $^{14}N$  (relative to  $^{15}N$ ) when nutrients are abundant (14). Our alternative interpretation of the sedimentary  $\delta^{15}N$  record is that the low values in sapropels are typical of eastern Mediterranean new production both today and throughout the late Pleistocene, and the high values in organic-deficient marl sediments result from extensive diagenetic alteration of nitrogen isotopic ratios in the presence of oxygen.

Support for our interpretation comes from nitrogen isotopic measurements of contemporary and fossil chlorins and dissolved nitrate. These measurements demonstrate that the contemporary eastern Mediterranean is characterized by very low isotopic values of phytoplankton and deep-water nitrate (Fig. 1A) that are similar to phytoplankton and sedimentary  $\delta^{15}N$  values in all six sapropels studied (Fig. 1, B and C). The deposition of recent (non-sapropel) sediments with a  $\delta^{15}N$  value of  $4.3\text{‰}$  (Fig. 1A) is an enigma that must be reconciled with the very low isotopic values of reactive nitrogen reservoirs within the eastern Mediterranean.

We determined the nitrogen isotopic composition of living and ancient phytoplankton by measuring the  $\delta^{15}N$  value of chlorins (15) and correcting for an empirically determined isotopic difference between chlorophyll and total

<sup>1</sup>Department of Environmental Science, Barnard College, Columbia University, 3009 Broadway, New York, NY 10027, USA. <sup>2</sup>Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA.

\*To whom correspondence should be addressed. E-mail: jpsachs@barnard.columbia.edu

## REPORTS

algal nitrogen (16). Chlorophyll, a photosynthetic pigment synthesized by all oxic photoautotrophs, is an excellent biomarker for total algal biomass. Biosynthetic partitioning of nitrogen within microalgae yields chlorophyll that is isotopically depleted by  $5.1 \pm 1.1\%$  (95% confidence interval;  $n = 8$  species) relative to whole algae (16). This result was corroborated by measurements of the isotopic difference between chlorophyll and suspended particles in a variety of marine settings (16) that averaged  $5.3 \pm 1.5\%$  ( $n = 6$ ). Although early diagenesis results in some structural changes to the chlorophyll molecule, transformations that

remove chlorins from our analytical window, such as macrocycle cleavage, or that involve functional groups on the macrocycle periphery are not expected to alter chlorin  $\delta^{15}\text{N}$  values because they occur at least two bond lengths from all nitrogen atoms.


Pyropheophorbide *a* (17), the most abundant chlorin in sapropels S2 through S7, had a nearly constant  $\delta^{15}\text{N}$  value of  $-5.0 \pm 0.4\%$  ( $n = 7$ ) (Fig. 1, B and C), which was close to the isotopic composition of chlorophyll from living phytoplankton collected in the eastern Mediterranean in May and June 1996 of  $-6.4 \pm 1.4\%$  ( $n = 3$ ) (Fig. 1A). Adjusting for the isotopic

depletion of chlorophyll relative to total algal nitrogen yields mean  $\delta^{15}\text{N}$  values for phytoplankton of  $0.0 \pm 1.4\%$  in the sapropels, and  $-1.4 \pm 1.9\%$  in the contemporary eastern Mediterranean. The similar or slightly higher  $\delta^{15}\text{N}$  value of phytoplankton during sapropel events compared with today is evidence for oligotrophic surface waters, an inference that is supported by micropaleontological evidence (6, 7) for stratified, nutrient-impoorished surface waters at those times. If nutrient supply had exceeded demand by phytoplankton, then lower isotopic values than those measured under nutrient-depleted (that is, contemporary) conditions should be observed (13).

Although our phytoplankton samples were collected during a single spring-time cruise, the nitrogen isotopic composition of deep-water nitrate,  $-0.7 \pm 0.1\%$  ( $n = 2$ ) (Fig. 1A), provides support for very low mean annual  $\delta^{15}\text{N}$  values in phytoplankton. Nitrate in the deep eastern Mediterranean is the principal reservoir of reactive nitrogen in the basin (1). With a residence time of 50 to 80 years (18), it represents a spatial and temporal integration of new production over the entire basin for many decades. Thus, the time-averaged  $\delta^{15}\text{N}$  value of material sinking from the euphotic zone must approximate this value in the steady state. The remarkably light isotopic value of abyssal nitrate [versus the global deep-ocean average of  $5.7 \pm 0.7\%$  (19)] is good evidence for very low mean annual  $\delta^{15}\text{N}$  values of phytoplankton in the contemporary eastern Mediterranean. Also consistent with the low isotopic values of phytoplankton and deep nitrate is the  $\delta^{15}\text{N}$  value of suspended particulate material in the euphotic zone,  $-0.6 \pm 0.1\%$  ( $n = 3$ ) (Fig. 1A). This mixture of algal biomass and detritus is expected to have a longer residence time than living phytoplankton, and therefore to have an isotopic composition similar to that of deep nitrate.

The extremely low isotopic values of contemporary phytoplankton, nitrate, and suspended particles require that nitrogen fixation supply a substantial fraction of new nitrogen to the Mediterranean Sea. Microbial nitrogen fixation imparts an isotopic fractionation to atmospheric  $\text{N}_2$  ( $\delta^{15}\text{N} = 0\%$ ), the nitrogenous substrate, producing organic matter with a mean  $\delta^{15}\text{N}$  value of  $-2.6 \pm 1.3\%$  (20). When this biomass is remineralized it yields isotopically depleted nitrate. Other ocean basins where extensive nitrogen fixation occurs, such as the Gulf of Mexico, the southern Arabian Sea, and the subtropical Pacific, are characterized by low nitrogen isotopic ratios (21, 22).

Extensive  $\text{N}_2$  fixation in the Mediterranean Sea has been inferred from nutrient budgets and elevated N/P ratios in deep waters (that is, N/P = 22 to 29 versus the marine average of 15) (23–25). Whereas the budget calculations imply a 7 to 41% contribution of biological  $\text{N}_2$  fixation to the Mediterranean Sea reactive nitrogen inventory (25), the isotopic data indicate



**Table 1.** Nitrogen isotopic alteration during early diagenesis.

| Bottom-water $\text{O}_2$ | Location              | $\Delta\delta^{15}\text{N}_{\text{sed/trap}}^*$ | $\Delta\delta^{15}\text{N}_{\text{sed/susp}}$ | $\Delta\delta^{15}\text{N}_{\text{sed/phyt}}$ |
|---------------------------|-----------------------|---|---|---|
| Well-oxygenated           | Southern Ocean        | 7.5   | 4.5   |   |
|                           | Eastern Mediterranean |   | 4.9   | 5.7   |
|                           | Equatorial Pacific    |   | 6   |   |
| Low oxygen                | Arabian Sea           | 2–4   |   |   |
|                           | Peru Margin           |   | 2   | -0.5  |
| Anoxic                    | Black Sea             |   | -1  | 0.7   |
|                           | Cariaco Trench        |   | -1.5  |   |
|                           | Framvaren Fjord       | -0.4  | -2  |   |

\*The isotopic difference between surface sediments and either deep-water sediment trap material (trap), suspended particles in overlying surface waters (susp), or phytoplankton (phyt). Phytoplankton  $\delta^{15}\text{N}$  values were calculated from chlorin  $\delta^{15}\text{N}$  values by adding 5.1 per mil (16). Additional nitrogen isotopic data are from (13, 29).

that diazotrophy supplies 46 to 70% of eastern basin new nitrogen (26). Large increases of fixed nitrogen in phosphorous addition experiments with surface water, and extreme phosphorous limitation of primary production provide additional evidence for high rates of diazotrophy in the eastern Mediterranean (24). That widespread nitrogen fixation has been overlooked by field studies is perhaps not surprising in light of recent evidence (22, 23) for dramatically higher global fixation rates than previously reported. In addition to contemporary populations of picocyanobacteria, *Synechococcus* (cyanobacterium), and *Posidonia* (sea grass) (24, 25), extensive rhizosolenid diatom mats living during sapropel episodes (7) may have supported nitrogen-fixing bacterial symbionts (27). Other sources of nitrogen to the Mediterranean Sea, such as rivers, rain, and north Atlantic seawater, have isotopic values between 1 and 6‰ (19, 28). Altering the contribution of any of these nitrogen sources cannot produce an eastern Mediterranean nitrogen inventory with an isotopic value less than 0‰; nitrogen fixation must be invoked.

With nitrogen isotopic values of 0‰ or lower in phytoplankton and the large abyssal nitrate reservoir, it remains to be explained why sediments between sapropel layers and in core-tops have high δ<sup>15</sup>N values of 4 to 7‰ (Fig. 1). We hypothesize that decomposing organic matter becomes enriched in <sup>15</sup>N in the presence of oxygen. The severity of the isotopic enrichment appears to be inversely related to bottom-water oxygen concentrations. When bottom waters are well oxygenated, as they are in the Southern Ocean, the central equatorial Pacific, and the eastern Mediterranean, sediments are enriched in <sup>15</sup>N by 4 to 6‰ relative to sinking and suspended particles (13, 29), and phytoplankton (Table 1). Under low bottom-water oxygen conditions, such as in the Arabian Sea, surface sediments are 2 to 4‰ enriched in <sup>15</sup>N relative to deep-sinking particles (29). When bottom waters are anoxic, such as in the Peru Margin, the Black Sea, the Cariaco Trench, and the Framvaren Fjord, surface sediments are within 2‰ of sinking and suspended particles (29) and phytoplankton (Table 1). Because phytoplankton δ<sup>15</sup>N values in the sapropels are nearly identical to the associated whole-sediment δ<sup>15</sup>N values, bottom waters were likely anoxic during those events (30), corroborating faunal (4) and geochemical (31) evidence for bottom-water oxygen deficiency.

Our nitrogen isotopic measurements support a model of sapropel generation in which lowered salinities (4) in regions where intermediate waters form cause pycnocline shoaling and diminished ventilation of eastern basin deep water (6, 32). A shallower pycnocline would allow the incursion of nutrient-enriched intermediate water into the euphotic zone to fuel new production by a deep-dwelling algal community (6), while diminished ventilation rates would

facilitate deep-water anoxia (6) and the preservation of organic matter (9). Whereas the contemporary eastern Mediterranean pycnocline lies below the euphotic depth, numerical simulations and micropaleontological studies indicate that it shoaled to 50 to 120 m during sapropel events (6, 32). This would account for the occurrence in sapropel layers of rhizosolenid diatoms (7) and certain calcareous plankton species (6, 11) adapted to a deep nutrient source (33). High abundances of rhizosolenids in sapropels are consistent with nitrogen isotopic evidence for surface-water nutrient depletion because these algae live in stratified, oligotrophic water (27).

Enhanced discharge of east African monsoonal rains through the Nile River and increased runoff from Eurasian rivers (34) likely triggered the incursion of nutrient-rich intermediate waters into the euphotic zone (6, 32) but, contrary to previous inferences (3), did not supply appreciable quantities of nitrogen for phytoplankton growth. Global mean isotopic values of freshwater and estuarine total nitrogen are 4.3 ± 2.7‰ (n = 64) and 4.6 ± 2.0‰ (n = 199), respectively (28), and contemporary groundwater nitrate in the lower Nile region is between 2.9 to 14.5‰ (35). Phytoplankton δ<sup>15</sup>N values of 0‰ in the sapropels thus preclude any dramatic increase in the flux of freshwater nitrogen. Two additional lines of evidence arguing against appreciable nutrient loading by runoff are observations of efficient biological removal of nutrients in estuaries (36), and mass balance calculations indicating that a flooding Nile could increase new nitrogen fluxes by no more than 10% (37).

Nitrogen isotopic measurements in fossil chlorophyll provide a new tool for understanding the oceanographic conditions during eastern Mediterranean sapropel events. Stratified, nutrient-impooverished surface water supported high rates of nitrogen fixation that contributed substantially to reactive nitrogen pools. Nitrogen from enhanced river runoff was insignificant compared with diazotrophy as a new nitrogen source. In contrast to the contemporary eastern Mediterranean, where well-oxygenated deep water facilitates the diagenetic enrichment of <sup>15</sup>N in decomposing organic matter, anoxic bottom waters during sapropel deposition preserved nitrogen isotopic ratios.

References and Notes

1. J. P. Bethoux, *Mar. Chem.* **10**, 141 (1981); *Deep-Sea Res.* **36**, 769 (1989).
2. R. C. Dugdale and F. P. Wilkerson, *Oceanol. Acta* **9**, 179 (1988).
3. S. E. Calvert, *Oceanol. Acta* **6**, 255 (1983).
4. M. B. Cita et al., *Quat. Res.* **8**, 205 (1977); R. B. Kidd, M. B. Cita, W. B. F. Ryan, *Init. Rep. Deep Sea Drill. Prog.* **42**, 421 (1978).
5. S. E. Calvert, B. Nielsen, M. R. Fontugne, *Nature* **359**, 223 (1992).
6. E. J. Rohling and W. W. C. Gieskes, *Paleoceanography* **4**, 531 (1989); E. J. Rohling, *Mar. Geol.* **122**, 1 (1994).
7. A. E. S. Kemp, R. B. Pearce, I. Koizumi, J. Pike, S. J. Rance, *Nature* **39**, 57 (1999).

8. B. J. H. Van Os, L. J. Lourens, F. J. Hilgen, G. J. De Lange, *Paleoceanography* **9**, 601 (1994).
9. R. Cheddadi and M. Rossignol-Strick, *Paleoceanography* **10**, 301 (1995); H. E. Hartnett, R. G. Keil, J. I. Hedges, A. H. Devol, *Nature* **391**, 572 (1998).
10. G. J. de Lange and H. L. ten Haven, *Nature* **305**, 797 (1983); J. A. Jenkins and D. F. Williams, *Mar. Micropaleontol.* **8**, 521 (1983/84); R. C. Thunell and D. F. Williams, *Nature* **338**, 493 (1989); M. W. Howell and R. C. Thunell, *Mar. Geol.* **103**, 461 (1992).
11. D. Castradori, *Paleoceanography* **8**, 459 (1993); E. J. Rohling, F. J. Jorissen, C. Vergnaud-Grazzini, W. J. Zachariasse, *Mar. Micropaleontol.* **21**, 191 (1993).
12. Standard delta notation is used for reporting stable isotopic ratios of nitrogen:

$$\delta^{15}\text{N} = \left[ \frac{^{15}\text{N}/^{14}\text{N}_{\text{sample}}}{^{15}\text{N}/^{14}\text{N}_{\text{standard}}} - 1 \right] \times 1000$$

and the isotopic standard is atmospheric N<sub>2</sub> with δ<sup>15</sup>N = 0‰.

13. M. A. Altabet, in *Particle Flux in the Ocean*, V. Ittekkot, P. Schafer, S. Honjo, J. Depetris, Eds. (Wiley, London, 1996), pp. 155–184.
14. E. Wada and A. Hattori, *Geomicrobiol. J.* **1**, 85 (1978).
15. J. P. Sachs and D. J. Repeta, *Org. Geochem.*, in press.
16. J. P. Sachs, D. J. Repeta, R. Goericke, *Geochim. Cosmochim. Acta* **63**, 1431 (1999).
17. Pyropheophorbide a is the trivial (Fischer) name for 13<sup>2</sup>-demethoxycarbonyl-pheophorbide a.
18. J. P. Bethoux, B. Gentili, J. Raunet, D. Tailliez, *Nature* **347**, 660 (1990); R. Schlitzer et al., *Deep-Sea Res.* **38**, 1531 (1991).
19. K.-K. Liu and I. R. Kaplan, *Limnol. Oceanogr.* **34**, 820 (1989).
20. T. C. Hoering and H. T. Ford, *J. Am. Chem. Soc.* **82**, 376 (1960); C. C. Delwiche and P. L. Steyn, *Environ. Sci. Technol.* **4**, 929 (1970); M. Minagawa and E. Wada, *Mar. Chem.* **19**, 245 (1986); S. A. Macko, M. L. F. Estep, P. E. Hare, T. C. Hoering, *Chem. Geol.* **65**, 79 (1987); E. J. Carpenter, H. R. Harvey, B. Fry, D. G. Capone, *Deep-Sea Res.* **44**, 27 (1997).
21. S. A. Macko, L. Entzeroth, P. L. Parker, *Naturwissenschaften* **71**, 374 (1984); D. G. Capone et al., *Mar. Ecol. Prog. Ser.* **172**, 281 (1998).
22. D. Karl et al., *Nature* **388**, 533 (1997).
23. N. Gruber and J. L. Sarmiento, *Global Biogeochem. Cycles* **11**, 235 (1997).
24. D. J. Bonin, M. C. Bonin, T. Berman, *Aquat. Sci.* **21**, 131 (1989); M. D. Krom, N. Kress, S. Brenner, L. I. Gordon, *Limnol. Oceanogr.* **36**, 424 (1991).
25. J. P. Bethoux and G. Copin-Montegut, *Limnol. Oceanogr.* **31**, 1353 (1986); J. P. Bethoux, P. Morin, C. Madec, B. Gentili, *Deep-Sea Res.* **39**, 1641 (1992).
26. The fraction x of eastern basin reactive nitrogen derived from N<sub>2</sub> fixation can be estimated by mass balance: δ<sub>fix</sub> × x + δ<sub>other</sub> × (1 - x) = δ<sub>nitrate</sub>, where δ<sub>fix</sub> is the δ<sup>15</sup>N value of fixed nitrogen (-2.6‰), δ<sub>other</sub> is the weighted average δ<sup>15</sup>N of all other new nitrogen sources, and δ<sub>nitrate</sub> is the δ<sup>15</sup>N of deep-water nitrate (-0.7‰), the principle reactive nitrogen pool. By considering a range of plausible values for δ<sub>other</sub>—which includes rivers, Western Mediterranean surface water, and precipitation—of 1 to 4‰ (28), it is calculated that N<sub>2</sub> fixation accounts for 46 to 70% of eastern basin new nitrogen sources. Although this fraction is larger than that derived from nitrogen budgets of the entire Mediterranean Sea (25), it is consistent with the west-east gradient in δ<sup>15</sup>N values of suspended particles and deep-water nitrate across the two basins. <sup>15</sup>N enrichments in the western (S. Pantoja, D. J. Repeta, J. P. Sachs, D. M. Sigman, unpublished data) compared with the eastern basin indicate a larger diazotrophic contribution to the reactive nitrogen pool of the eastern Mediterranean.
27. R. R. L. Guillard and P. Kilham, in *The Biology of Diatoms*, D. Werner, Ed. (Univ. of California Press, Berkeley, CA, 1977), vol. 13, pp. 372–469; E. J. Carpenter, in *Nitrogen in the Marine Environment*, E. J. Carpenter and D. C. Capone, Eds. (Academic Press, New York, 1983), pp. 65–103; T. A. Villareal, *Mar. Ecol. Prog. Ser.* **76**, 201 (1991).
28. N. J. P. Owens, *Adv. Mar. Biol.* **24**, 390 (1987).
29. Supporting nitrogen isotopic data are from the fol-

lowing sources: E. Wada, M. Terazaki, Y. Kabaya, T. Nemoto, *Deep-Sea Res.* **34**, 829 (1987) (Southern Ocean); P. Schafer and V. Ittekkot, *Naturwissenschaften* **80**, 511 (1993) (Arabian Sea); S. M. Libes and W. G. Deuser, *Deep-Sea Res.* **35**, 517 (1988) (Peru margin suspended particles); B. Fry *et al.*, *Deep-Sea Res.* **38**, S1003 (1991) (Peru margin, Black Sea, and Cariaco Trench suspended particles); D. J. Velinsky and M. L. Fogel, *Mar. Chem.* **67**, 161 (Framvaren Fjord).

30. Microbial nitrate reduction (denitrification) in the absence of oxygen is associated with a large kinetic isotope effect that causes isotopic enrichment of the remaining nitrate pool [F. A. Richards and B. B. Benson, *Deep-Sea Res.* **7**, 254 (1961); J. D. Cline and I. R. Kaplan, *Mar. Chem.* **3**, 271 (1975)], a signal transferrable to phytoplankton and sediments [T. Saino and A. Hattori, *Deep-Sea Res.* **34**, 807 (1987); M. A. Altabet, R. Francois, D. W. Murray, W. L. Prell, *Nature* **373**, 506 (1995); R. S. Ganeshram, T. F. Pedersen, S. E. Calvert, J. W. Murray, *Nature* **376**, 755 (1995)]. However, denitrification will lead to an increase in  $\delta^{15}\text{N}$  only if the rate of nitrate replenishment is greater than the rate of denitrification. If the rate of denitrification exceeds the rate of nitrate supply, the likely case during eastern Mediterranean sapropel events when deep-water ventilation was reduced (32) and thermohaline circulation was similar or slowed (32) [R. Zahn, M. Sarnthein, M. Erlenkeuser, *Paleoceanography* **2**, 543 (1987)] compared with today, no net isotopic fractionation is expressed. In addition, if eastern basin nitrate was isotopically enriched through denitrification during sapropel events, it should be expressed in both chlorophyll and sediments. Instead, we observe that chlorophyll  $\delta^{15}\text{N}$  remains constant whereas sediment  $\delta^{15}\text{N}$  decreases in sapropels compared with contemporary and non-sapropel values.

31. A. Mangini and J. Dominik, *Sediment. Geol.* **23**, 113 (1979); H. F. Passier *et al.*, *Nature* **397**, 146 (1999).

32. P. G. Myers, K. Haines, E. J. Rohling, *Paleoceanography* **13**, 586 (1998).

33. The incursion of nutrient-rich intermediate waters into the euphotic zone would not lead to excess supply of nutrients and ensuing nitrogen isotopic fractionation (5). When the pycnocline lies within the euphotic zone of a stratified basin, a deep algal community develops and autotrophic nutrient uptake causes the retreat of the nutricline below the euphotic depth (6). Implicit in this retreat is expansion of the algal population to meet the increased supply of nutrients. Thus, a nutrient excess within the euphotic zone would not occur, and kinetic fractionation would not be expressed (14).

34. M. Rossignol-Strick, W. Nesteroff, P. Olive, C. Vergnaud-Grazzini, *Nature* **295**, 105 (1982); H. F. Shaw and G. Evans, *Mar. Geol.* **61**, 1 (1984); M. Rossignol-Strick, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **49**, 237 (1985); *Paleoceanography* **2**, 333 (1987).

35. A. I. M. Aly, M. A. Mohamed, E. Hallaba, in *Stable Isotopes*, H.-L. Schmidt, H. Förstel, K. Heinzinger, Eds. (Elsevier, Amsterdam, 1982), pp. 475–481.

36. T. D. Jickells, *Science* **281**, 217 (1998).

37. Estimates of the water flux from a flooding Nile are about 2.5 times the modern (pre-Aswan Dam) flow (34), or 100 to 200 km<sup>3</sup> year<sup>-1</sup> [W. F. Wadie, *Acta Adriatica* **25**, 29 (1984)]. Nitrogen concentrations in desert rivers are very low, with total dissolved nitrogen being 3  $\mu\text{M}$  in a representative unpolluted river (38). A conservative assumption is that Nile River dissolved nitrogen concentrations increased during its flood, possibly to the high values associated with tropical rivers, which are between 2 to 20  $\mu\text{M}$  nitrate (38). Assuming the high value of 20  $\mu\text{M}$  [compared with total dissolved nitrogen concentrations of 8.3  $\mu\text{M}$  in average world river water (38)], and twice the high estimate of predam discharge rates, nitrogen input from the Nile during flood is estimated at  $0.06 \times 10^{12}$  g N year<sup>-1</sup>. Estimates of new production from nutrient and oxygen budgets in the contemporary eastern Mediterranean Sea range from 6 to 12 g C m<sup>-2</sup> year<sup>-1</sup> (7). Assuming a molar C/N ratio of 6.6 gives nitrogen new production fluxes of 1.1 to 2.3 g N m<sup>-2</sup> year<sup>-1</sup>, similar to a previous estimate of 1.5 g N m<sup>-2</sup> year<sup>-1</sup> (2) derived from measured primary production. Given an area of  $1.66 \times 10^{12}$  m<sup>2</sup>, today's integrated new nitrogen production for the entire

eastern basin is  $1.9$  to  $3.8 \times 10^{12}$  g N year<sup>-1</sup>. Finally, comparing our conservative estimate of dissolved inorganic nitrogen discharge from a flooding Nile ( $0.06 \times 10^{12}$  g N year<sup>-1</sup>) to total eastern Mediterranean new nitrogen production today ( $1.9$  to  $3.8 \times 10^{12}$  g N year<sup>-1</sup>) demonstrates that nutrient delivery from the Nile was unlikely to have increased new nitrogen production by more than 1.6 to 3.2% over contemporary rates. Increasing the Nile discharge rate beyond 2.5 times the modern value, and adding a contribution from Eurasian rivers such that riverine runoff was four times that of the predam Nile discharge would only increase new nitrogen production by 5 to 10%.

38. M. Meybeck, *Am. J. Sci.* **282**, 401 (1982).

39. D. M. Sigman *et al.*, *Mar. Chem.* **57**, 227 (1997).

40. We thank B. Fry and D. Sigman for isotopic analyses. P. Rimbault, H. Claustra, and the crew of R/V *Le Suroit* provided ship time and assistance with sampling. The Ocean Drilling Program provided sediment samples. Funding came from NSF Chemical Oceanography (D.J.R.), an Office of Naval Research Graduate Fellowship (J.P.S.), an American Chemical Society Petroleum Research Fund Grant (D.J.R.), and a Woods Hole Oceanographic Institution Ocean Ventures Fund Grant (J.P.S.).

29 June 1999; accepted 5 November 1999

## Heavy Nitrogen in Carbonatites of the Kola Peninsula: A Possible Signature of the Deep Mantle

Nicolas Dauphas<sup>1\*</sup> and Bernard Marty<sup>1,2</sup>

Nitrogen and argon isotopes were measured in carbonatites and associated rocks from the Kola Peninsula in Russia. The Kola mantle source, which is thought to be located in the deep mantle, is enriched in heavy nitrogen (+3 per mil relative to air) as compared to Earth's surface (atmosphere and crust, +2 per mil) and the shallow mantle (-4 per mil). Recycling of oceanic crust (+6 per mil) or metal-silicate partitioning may account for the nitrogen isotopic composition of the deep mantle.

Determining the structure and composition of the mantle is necessary in order to understand current and past mantle dynamics and mantle-to-crust interactions. For example, some have argued that part of the mantle has been isolated from mantle convection for most of Earth's history, allowing the preservation of a primitive component that is only sampled by plumes. Evidence in support of this view arises mainly from noble gases, because plume-derived magmas often show lower radiogenic/primitive isotope ratios than do mid-ocean ridge basalts (MORBs) (1). At variance with these models are geophysical (2), experimental (3), and geochemical (4) lines of evidence that suggest that some of the subducting slabs sink through the 670-km-deep seismic discontinuity, implying global stirring of the mantle.

The nitrogen isotopic composition of the shallow mantle that feeds MORs, which is expressed as per mil (‰) deviation relative to the composition of air [in  $\delta^{15}\text{N}$  notation (5)], is estimated to be  $\sim -4\text{‰}$  (6–13). This signature

may be a remnant of the nitrogen isotopic composition of Earth-forming planetesimals, which later evolved as a result of (i) addition of meteoritic (14, 15) and cometary (15) volatiles, or (ii) fractional loss of atmospheric volatiles (16), or both. Biologic activity leads to a fractionation of nitrogen isotopes and gives sedimentary rocks a specific composition [ $\delta^{15}\text{N} \sim +6\text{‰}$  (17)]. Nitrate is used by denitrifying bacteria as the terminal electron acceptor in energy generation when oxygen is unavailable. Associated with denitrification is a kinetic isotope effect that enriches the residual nitrate in  $^{15}\text{N}$  (18). Because  $\text{NO}_3^-$  is the main nitrogen-bearing nutrient, marine organisms and sediments are enriched in  $^{15}\text{N}$  relative to the atmosphere (17). The oceanic crust thus enriched in  $^{15}\text{N}$  is subducted back into Earth at convergent plate margins, which makes nitrogen a potentially powerful tracer of volatile recycling in the mantle.

Mantle plumes, which are assumed to be fixed relative to plate motion, sample a deeper region than that feeding MORs (19). The isotopic composition of nitrogen in plumes has not been documented (20). Low  $^4\text{He}/^3\text{He}$  ratios and a steep  $^{20}\text{Ne}/^{22}\text{Ne}$ - $^{21}\text{Ne}/^{22}\text{Ne}$  correlation (21) indicate that a mantle plume contributed to the 370-million-year-old (22) ultrabasic-alkaline-carbonatite magmatism in the Kola Peninsula in Russia (the eastern segment of the Baltic Shield). The Kola rocks crystallized at depth (23), which prevented them from being extensively outgassed, minimizing any subsequent

<sup>1</sup>Centre de Recherches Pétrographiques et Géochimiques, CNRS UPR 9046, 15 rue Notre-Dame des Pauvres, Boîte Postale 20, 54501 Vandoeuvre-lès-Nancy Cedex, France. <sup>2</sup>Ecole Nationale Supérieure de Géologie, rue du doyen Marcel Roubault, Boîte Postale 40, 54501 Vandoeuvre-lès-Nancy Cedex, France.

\*To whom correspondence should be addressed. E-mail: dauphas@crpg.cnrs-nancy.fr