



Hylid frogs are some of the most charismatic and endangered anurans in Mexico. During a time where amphibian extinctions are becoming a common occurrence, it is important to celebrate rediscoveries. With rediscoveries of endangered species also comes an urgent need to acquire basic natural history information and population data. Pictured here is an individual of *Duellmanohyla ignicolor* found near La Esperanza, Oaxaca, Mexico. © Itzue W. Caviedes Solis



Missing in action for nearly 50 years: rediscovery of the Mexican treefrog *Duellmanohyla ignicolor* (Anura: Hylidae)

ROSS FURBUSH^{1,2}, ITZUE W. CAVIEDES-SOLIS¹, FAUSTO R. MÉNDEZ-DE LA CRUZ³, AND ADAM D. LEACHÉ¹

¹Department of Biology and Burke Museum of Natural History and Culture, University of Washington, Box 351800, Seattle, Washington 98195, United States. Email: itzuecs@uw.edu (Corresponding author)

²Current: Department of Science, Principia Upper School, 132301 Clayton Rd, St. Louis, Missouri 63131, United States.

³Laboratorio de Herpetología, Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, C.P. 04510, Apartado Postal 70–153, México, D.F., Mexico.

ABSTRACT: The genus *Duellmanohyla* currently is comprised of eight species distributed from eastern Veracruz, Mexico, to western Panama. Three species in this genus occur in Mexico (*D. chamulae*, *D. ignicolor*, and *D. schmidtorum*), of which the first two are endemic to the country and are considered Endangered by the IUCN, whereas *D. schmidtorum* is classified as Vulnerable. We conducted surveys at the type localities and surrounding regions of the species of *Duellmanohyla* distributed in Mexico. We found one individual of *D. chamulae* near Rayón Mezcalapa, Chiapas; three individuals of *D. schmidtorum* from San Rafael, Chiapas; and rediscovered three individuals of *D. ignicolor*, a species that has not been found since 1970, near La Esperanza, Santiago Comaltepec, Oaxaca.

Key Words: *Duellmanohyla ignicolor*, *Duellmanohyla chamulae*, *Duellmanohyla schmidtorum*, Mexico, Oaxaca, rediscovery

RESUMEN: El género *Duellmanohyla* está compuesto por ocho especies que se distribuyen del este de Veracruz, México, hasta el oeste de Panamá. Tres especies en este género se distribuyen en México (*D. chamulae*, *D. ignicolor* y *D. schmidtorum*), de cual las dos primeras son endémicas al país y están consideradas En Peligro por UICN, mientras que *D. schmidtorum* está clasificada como Vulnerable. El trabajo de campo fue realizado en los alrededores de las localidades tipo de las especies de *Duellmanohyla* que se distribuyen en México. Reportamos un individuo de *D. chamulae* cerca de Rayón Mezcalapa, Chiapas; tres individuos de *D. schmidtorum* de San Rafael, Chiapas; y redescubrimos tres individuos de *D. ignicolor*, una especie que no se había encontrado desde 1970, cerca de La Esperanza, Santiago Comaltepec, Oaxaca.

Palabras Claves: *Duellmanohyla ignicolor*, *Duellmanohyla chamulae*, *Duellmanohyla schmidtorum*, Mexico, Oaxaca, redescubrimiento

Citation: Furbush, R., I. W. Caviedes-Solis, F. R. Méndez-de la Cruz, and A. D. Leaché. 2017. Missing in action for nearly 50 years: rediscovery of the Mexican treefrog *Duellmanohyla ignicolor* (Anura: Hylidae). *Mesoamerican Herpetology* 4: 120–127.

Copyright: Furbush et al. 2017. This work is licensed under a Creative Commons Attribution-No Derivatives 4.0 International License.

Received: 11 November 2016; **Accepted:** 11 January 2017; **Published:** 30 March 2017.

INTRODUCTION

Mexico contains a remarkable amount of amphibian diversity and endemism, as it harbors 376 species of amphibians, of which 60% are endemic to the country (Flores-Villela, 1993; Parra-Olea et al., 2014). Approximately 20% of the world's amphibian diversity is found in Mexico, a country that ranks third with respect to global amphibian endemism (Stuart et al., 2008). Nonetheless, the same anthropomorphic and climatic pressures that are impacting other amphibian populations in Latin America and the world are threatening Mexico's unique amphibian diversity (Young et al., 2001; Lips et al. 2004; Stuart et al., 2008; Frías-Alvarez et al., 2010). In particular, montane stream-breeding frogs in Latin America are susceptible to unexpected population declines, even within protected areas (Pounds et al., 1997; Lips, 1999). Low population density or species rarity can lead to the false assumption that a species has gone extinct, but with comprehensive surveys such species sometimes can be rediscovered (Caviedes-Solis et al., 2015).

The genus *Duellmanohyla* currently is comprised of eight species distributed from eastern Veracruz, Mexico, to western Panama (Aguilar-López et al., 2010; Frost, 2016). Three of the species are distributed in Mexico: *D. chamulae* and *D. ignicolor* were described from Vista Hermosa, Sierra de Juarez, Oaxaca by Duellman (1961); and *D. schmidtorum* was described from Finca el Porvenir, San Marcos, Guatemala by Stuart (1954) (Fig. 1). *Duellmanohyla chamulae* and *D. ignicolor* are endemic to Mexico and considered Endangered by the International Union for Conservation of Nature (IUCN) (Santos-Barrera and Muñoz-Alonso, 2004a; Santos-Barrera et al., 2004), and *D. schmidtorum* is found in Guatemala and Mexico and is classified as Vulnerable (Santos-Barrera and Muñoz-Alonso, 2004b). Of these taxa, these authors noted *D. chamulae* as “not a rare species”; however, *D. schmidtorum* was indicated as “not very common” and *D. ignicolor* as “relatively uncommon and known only from a few localities.” Mexican law protects the three species under the “Special Protection” category (SEMARNAT, 2010).

Whereas individuals of both *D. schmidtorum* and *D. chamulae* have been found during the last seven years (Muñoz-Alonso, 2010), the last known record for *D. ignicolor* corresponds to 24 adult males collected in 1970 (Caldwell, 1974); females of this species remain unknown. Previous surveys for stream-breeding frogs in the Mexican highlands have failed to find *D. ignicolor*; no adults were detected during the years 2000, 2004, 2007, 2010, 2011, or 2014, despite surveys conducted in streams within the expected elevational range of 680–1,850 m at the type locality in the Sierra de Juarez (Lips et al., 2004; Delia et al., 2013; Caviedes-Solis et al., 2015). Thirteen tadpoles found in 2004 tentatively were referred to as *D. ignicolor* (Lips et al., 2004); however, after further inspection they did not match the description and illustrations provided by Duellman (2001) (J. Mendelson III, pers. comm.). Subsequently, these specimens were lost.

MATERIALS AND METHODS

We surveyed localities in the surrounding areas of the type localities of the three species of *Duellmanohyla* distributed in Mexico: Santiago Comaltepec, Oaxaca (3 days), Rayón Mezcalapa (2), and San Rafael, Chiapas (1) (Fig. 1). The collecting effort included two field biologists: Itzue Caviedes-Solis and Ross Furbush. We searched for approximately six hours each night in September of 2015, using visual encounter and acoustic surveys along slow flowing streams. We captured the frogs by hand and euthanized them with an anesthetic, according to the University of Washington IACUC protocol (#4209-01). We collected liver tissue samples and preserved them in ethanol for molecular genetic studies. We fixed the specimens with buffered formalin and preserved them in 70% ethanol. All specimens are deposited at the Burke Museum of Natural History and Culture, University of Washington (Accession #s 2016–22).




Fig. 1. Distribution map of species in the genus *Duellmanohyla* in Mexico.

© Itzue-Caviedes-Solis

RESULTS

On 8 September 2015, we recorded three male individuals of *Duellmanohyla ignicolor* from near La Esperanza, Santiago Comaltepec, Oaxaca (UWBM 9205–9207). We found the three frogs on tree leaves between 1 and 1.5 m above the ground: one was on a branch overhanging a slow flowing stream and the other two were found less than 2 m away from the water. The water source was located among the limits of cloud forest, cow pastures, and riparian vegetation. On 12 September 2015, we found one male *D. chamulae* near Rayón Mezcalapa, Chiapas (UWBM 2904), in a patch of cloud forest downstream from a cow pasture. The individual was calling approximately 5 m upslope from a slow flowing stream, and was found on a leaf approximately 1.5 m from the ground. We also found three individuals of *D. schmidtorum* near San Rafael, Chiapas (UWBM 2908–2910). On 14 September 2015, we found additional individuals of *D. schmidtorum* in cloud forest habitat, on leaves of low vegetation less than 1 m above the ground and less than 3 m from a fast flowing river. We provide color images of all the specimens in Fig. 2, and present their morphometric data in Table 1.

Our specimens match the descriptions of these species provided by Duellman (1961, 2001); however, we also recorded new color variation among adult males of *D. ignicolor*. The dorsum of one specimen (UWBM 9205; Fig. 2 D) was uniform green, whereas the dorsal coloration of the other three (KU 87160, UWBM 2206, 2207; Fig. 2 C, E, and F) contained various brown and dark green irregular markings and cream spots that extend from the dorsum to the arms and legs. The iris color of the holotype was described as pale gold, probably similar to the specimen in Fig. 2 C; however, the iris color of the three specimens we found (Fig. 2D–F) was pale red/pink. The original description of *D. ignicolor* noted red or orange red coloration on the thighs, ventral surfaces of the shanks, anterior surfaces of the tarsi, and upper proximal surfaces of the 1st, 2nd, and 3rd toes (Duellman, 2001); pale pink-orange was present in the same areas in our three specimens (Fig. 2D–F), but it also extended to the four fingers and five toes and toepads. As noted by Duellman (2001), all of our specimens have a white line on their sides, along with a discontinuous brown line below. Interrupted marked bands (Fig. 2C–F) are present on the dorsal parts of the thighs and arms, and one of the males (Fig. 2 F) showed scars on the head. We are not including detailed descriptions of *D. chamulae* and *D. schmidtorum*, but the color pattern of the specimens agrees with the variation noted by previous authors (Campbell and Smith, 1992; Duellman, 2001; Aguilar-López et al., 2010).



Fig. 2. Color variation in specimens of the genus *Duellmanohyla*. *D. chamulae* (A = KU 58062; and B = UWBM 9204); *D. ignicolor* (C = KU 87160; D = UWBM 9205; E = UWBM 9206; and F = UWBM 9207); *D. schmidtorum* (G = KU 58016-29; H = UWBM 9208; I = UWBM 9209; and J = UWBM 9210). A, C, and H correspond to specimens previously collected (AmphibiaWeb, 2016).

© Itzue W. Caviedes-Solis (B, D, E, F, H–J) and William E. Duellman (A, C, G).

Table 1. Morphological measurements (mm) for specimens of *Duellmanohyla*. UWBM = Burke Museum of Natural History and Culture, University of Washington. SVL – snout–vent length.

UWBM	Sex	SVL	Tibia	Head Length	Head Width	Eye Diameter	Tympanum Diameter
<i>D. chamulae</i>							
09204	M	28.21	13.47	9.38	8.71	3.29	1.79
<i>D. ignicolor</i>							
09205	M	35.39	18.75	11.05	10.90	4.68	2.15
09206	M	35.30	18.03	10.61	11.00	5.40	2.12
09207	M	34.71	18.75	10.97	10.95	4.56	2.02
<i>D. schmidtorum</i>							
09208	M	29.48	14.85	9.02	9.09	3.79	1.76
09209	M	30.23	14.91	9.63	9.41	3.38	1.68
09210	M	30.69	14.79	9.12	9.73	3.64	1.95

DISCUSSION

Recently, many amphibian surveys have been conducted in Sierra de Juarez during the rainy season, and all failed to detect *Duellmanohyla*. Our success in finding *D. ignicolor* in this region might have resulted from favorable weather conditions, as well as from surveys conducted in new localities with adequate microhabitats. During 2015 the climate phenomenon “El Niño” caused precipitation anomalies throughout Mesoamerica. Records show a significant precipitation reduction in Mexico during the summer before our surveys in 2015 (Servicio Meteorológico Nacional, 2016), which corresponds to the breeding season for most Mexican treefrogs (Duellman, 2001). Some studies have suggested that an increase in temperature along with a reduction of water availability could cause a displacement of the breeding season for species dependent on temporary ponds for reproduction (Blaustein et al., 2001). *Duellmanohyla ignicolor*, for example, requires slow flowing freshwater streams for breeding (Duellman, 1963). These conditions might explain why a delayed rainy season in the summer 2015 might have produced favorable breeding conditions for *D. ignicolor* in late September.

The genus *Duellmanohyla* was described as a monophyletic stream-breeding lineage from the Central American highlands (Campbell and Smith, 1992), but recent phylogenetic analyses found that *Duellmanohyla* is paraphyletic with respect to *Bromeliohyla* and *Ptychohyla* (Duellman, et al., 2016). These three genera are difficult to distinguish based on their adult morphology, and although their reproductive modes and larvae are distinctly different, “the existing molecular data are unable to resolve their relationships” (Duellman et al., 2016: 19). The new genetic resources that we collected from these new specimens of *D. ignicolor*, *D. chamulae*, and *D. schmidtorum* hopefully will contribute to resolving the phylogeny of this Mesoamerican group of frogs.

Mexican endemic species of *Duellmanohyla* are distributed in two states involved with ongoing conservation efforts. Santiago Comaltepec, in the state of Oaxaca, is a member of the “Payment For Environmental Services” program, which helps protect forested areas. In the same municipality, a project was dedicated to search for rare and endangered treefrogs, which is funded by a Conservation Leadership Fellowship (L. Canseco-Márquez, pers. comm.). The state of Chiapas hosts a program called “Strategy for the Conservation of Critically Threatened Amphibians from Chiapas, Mexico,” which meets annually to discuss local amphibian conservation (Luna-Reyes et al., 2010); it also maintains a compilation of records of the state’s amphibian diversity, including the conservation status for each species. A potential cause of amphibian population declines in Chiapas has been the pathogenic

fungus *Batrachochytrium dendrobatidis*. This pathogen has been reported as abundant in populations of *D. schmidtorum*, and although individuals of *D. chamulae* were not tested or found, the pathogen has been reported from within its potential distribution (Muñoz-Alonso, 2010). *Duellmanohyla ignicolor* might also have been affected by *B. dendrobatidis*, which has been found in Sierra de Juárez (Hernández, 2012).

During a time when amphibian extinctions are becoming more common, it is important to celebrate rediscoveries. With rediscoveries, however, also comes an urgent need to acquire basic natural history and population data for such endangered species as *D. ignicolor*. Although emerging conservation programs are present in the state of Oaxaca, these efforts could be more strategic if we knew the population status of *D. ignicolor* and understood the life history of this species. Treefrogs generally are cryptic and females in this genus do not call, unlike their male counterparts, which might explain why females of *D. ignicolor* have never been found. Females must be discovered to understand the sexual dimorphism, phenology, and breeding behavior of this species. Additional surveys will continue to improve our understanding of *D. ignicolor*, and allow us to promote the importance of endemic treefrogs to scientists, the Mexican government, and local communities.


Acknowledgments.—Itzue W. Caviedes-Solis was funded by a CONACyT fellowship No. 313501, “Becas CONACyT al Extranjero” and the Burke Museum of Natural History and Culture. We thank the local municipality authorities and Norma Hernández from Santiago Comaltepec for their help and hospitality, and the people from the communities of Rayón Mezcalapa and San Rafael for use of their facilities. We also thank the Dirección Nacional de Vida Silvestre, Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT), México, for granting the scientific collecting permits (FAUT 043).

LITERATURE CITED

- AGUILAR-LÓPEZ J. L., E. PINEDA, AND U. O. GARCÍA-VÁZQUEZ. 2010. Ampliación del ámbito geográfico-altitudinal de *Duellmanohyla chamulae* (Amphibia: Hylidae) y primer registro para la anfibiofauna de Veracruz. *Revista Mexicana de Biodiversidad* 81: 579–80.
- AMPHIBIAWEB. 2016. University of California, Berkeley, United States. (www.amphibiaweb.org; accessed 12 September 2016).
- BLAUSTEIN, A. R., L. K. BELDEN, D. H. OLSON, D. M. GREEN, T. L. ROOT, AND J. M. KIESECKER. 2001. Amphibian breeding and climate change. *Conservation Biology* 15: 1,804–1,809.
- CALDWELL, J. P. 1974. Tropical Tree Frog Communities: Patterns of Reproduction, Size, and Utilization of Structural Habitat. Unpublished Ph.D. dissertation, University of Kansas, Lawrence, Kansas, United States.
- CAMPBELL, J. A., AND E. N. SMITH. 1992. A new frog of the genus *Ptychohyla* (Hylidae) from the Sierra de Santa Cruz, Guatemala, and description of a new genus of Middle American stream-breeding tree frogs. *Herpetologica* 48: 153–167.
- CAVIEDES-SOLIS, I. W., L. F. VÁZQUEZ-VEGA, I. SOLANO-ZAVALA, E. PÉREZ-RAMOS, S. M. ROVITO, T. J. DEVITT, P. HEIMES, O. A. FLORES-VILLELA, J. A. CAMPBELL, AND A. NIETO MONTES DE OCA. 2015. Everything is not lost: recent records, rediscoveries, and range extensions of Mexican hylid frogs. *Mesoamerican Herpetology* 2: 230–241.
- DELIA, J. R., J. L. WHITNEY, AND T. BURKHARDT. 2013. Rediscovery of ‘lost’ tree frogs from the Oaxacan highlands of Mexico. *Biodiversity and Conservation* 22: 1,405–1,414.
- DUELLMAN, W. E. 1961. Descriptions of two new species of frogs, genus *Ptychohyla*: studies of American hylid frogs V. University of Kansas Publications, Museum of Natural History 13: 349–357.
- DUELLMAN, W. E. 1963. A review of the Middle American tree frogs of the Genus *Ptychohyla*. University of Kansas Publications, Museum of Natural History 15: 297–349.
- DUELLMAN, W. E. 2001. The Hylid Frogs of Middle America. 2 Volumes. 2nd ed. Contributions to Herpetology, Volume 18, Society for the Study of Amphibians and Reptiles, Ithaca, New York, United States.
- DUELLMAN, W. E., A. B. MARION, AND S. B. HEDGES. 2016. Phylogenetics, classification, and biogeography of the tree frogs (Amphibia: Anura: Arboranae). *Zootaxa* 4,104: 1–109.
- FLORES-VILLELA, O. A. 1993. Herpetofauna of Mexico: distribution and endemism. Pp 253–280 In T. P. Ramamoorthy, R. Bye, J. Lot, and J. Fa (Eds), *Biological Diversity of Mexico: Origins and Distribution*, Oxford University Press, New York, New York, United States.
- FRÍAS-ÁLVAREZ, P., J. J. ZÚÑIGA-VEGA, AND O. FLORES-VILLELA. 2010. A general assessment of the conservation status and decline trends of Mexican amphibians. *Biodiversity and Conservation* 19: 3,699–3,742.
- FROST, D. R. 2016. Amphibian Species of the World: an Online Reference. Version 5.4. American Museum of Natural History, New York, New York, United States. (www.research.amnh.org/vz/herpetology/amphibia/; accessed 15 September 2016).
- HERNÁNDEZ, R. C. 2012. Evaluación de la presencia del hongo *Batrachochytrium dendrobatidis*, en poblaciones de anfibios en área cero extinción (AZE) en Oaxaca y Chiapas, México. *Lacandonia*. 6: 7–16.
- IUCN 2016. The IUCN Red List of Threatened Species. Version 2015-3. (www.iucnredlist.org; accessed 9 September 2016).
- LIPS, K. R. 1999. Mass mortality of the anuran fauna at an upland site in Panama. *Conservation Biology* 13: 117–125.

- LIPS, K. R., J. R. MENDELSON III, A. MUÑOZ-ALONSO, L. CANSECO-MÁRQUEZ, AND D. G. MULCAHY. 2004. Amphibian population declines in montane southern Mexico: resurveys of historical localities. *Biological Conservation* 119: 555-564.
- LUNA-REYES R., A. RAMÍREZ-VELAZQUEZ, O. JIMÉNEZ-LANG, M. A. GARCÍA-JIMÉNEZ, Y. Y. MORALES-BARRALES, E. F. ZÚÑIGA-JUÁREZ, J. P. SÁNCHEZ-ESTRADA, AND J. M. ARANDA-COELLO. 2010. Estrategia para la conservación de anfibios críticamente amenazados en Chiapas, México. (No. CH/597.8097275 E8). Secretaría de Medio Ambiente, Vivienda e Historia Natural, Conservation International, and Critical Ecosystem Partnership Fund. 66 pp.
- MUÑOZ-ALONSO, A. 2010. Riqueza, diversidad y estatus de los anfibios amenazados en el sureste de México; una evaluación para determinar las posibles causas de la declinación de sus poblaciones. Colegio de la Frontera Sur, Chiapas, Mexico. 55 pp.
- PARRA-OLEA, G., O. FLORES-VILLELA, AND C. MENDOZA-ALMERALLA. 2014. Biodiversidad de anfibios en México. *Revista Mexicana de Biodiversidad* 85: 460-466.
- POUNDS, J. A., M. P. FOGDEN, J. M. SAVAGE, AND G. C. GORMAN. 1997. Test of null models for amphibian declines on a tropical mountain. *Conservation Biology* 11: 1,307-1,322.
- SANTOS-BARRERA G., AND A. MUÑOZ-ALONSO. 2004a. *Duellmanohyla chamulae*. The IUCN Red List of Threatened Species 2004: e.T55307A11287549. (www.iucnredlist.org; accessed 12 September 2016).
- SANTOS-BARRERA G., AND A. MUÑOZ-ALONSO. 2004b. *Duellmanohyla schmidtorum*. The IUCN Red List of Threatened Species 2004: e.T55312A11288835. (www.iucnredlist.org; accessed 12 September 2016).
- SANTOS-BARRERA G., A. MUÑOZ-ALONSO, AND L. CANSECO-MÁRQUEZ. 2004. *Duellmanohyla ignicolor*. The IUCN Red List of Threatened Species 2004: e.T55308A11287783. (www.iucnredlist.org; accessed 12 September 2016).
- SEMARNAT (SECRETARÍA DE MEDIO AMBIENTE Y RECURSOS NATURALES) 2010. Norma Oficial Mexicana NOM-059-SEMARNAT-2010, protección ambiental-especies nativas de México de flora y fauna Silvestre-categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-lista de especies en riesgo. Diario Oficial de la Federación. México, Mexico.
- SERVICIO METEOROLÓGICO NACIONAL. 2016: Meteorological data from Oaxaca, Mexico. 2015. doi:10.1594/PANGAEA.861745.
- STUART, L. C. 1954. Descriptions of some new amphibians and reptiles from Guatemala. *Proceedings of the Biological Society of Washington* 67: 159-177.
- STUART S. N., M. HOFFMANN, J. S. CHANSON, N. A. COX, R. J. BERRIDGE, P. RAMANI, AND B. E. YOUNG (Eds.). 2008. *Threatened Amphibians of the World*. Lynx Edicions, Barcelona, Spain; IUCN, Gland Switzerland; and Conservation International, Arlington, Virginia, United States.
- YOUNG, B., K. LIPS, J. REASER, R. IBAÑEZ, A. SALAS, R. CEDEÑO, L. COLOMA, S. RON, E. LAMARCA J. MEYER, A. MUÑOZ, F. BOLAÑOS, G. CHAVEZ, AND D. ROMO. 2001. Population declines and priorities for amphibian conservation in Latin America. *Conservation Biology* 15: 1,213-1,223.





Ross Furbush graduated *cum laude* with interdisciplinary honors from University of Washington in 2015 with B.S.'s in Biology (Evolution, Ecology, and Conservation), Environmental Science, and Resource Management (Wildlife Conservation), and minors in Aquatic and Fishery Sciences and Quantitative Science. Ross' capstones examined the biogeography of Corvids in the Puget Sound Region and the impact of window-strike mortality on migrating avifauna. He currently works as a sustainability and biology teacher at Principia Upper School in St. Louis, Missouri.



Itzue W. Caviedes-Solis is a Mexican herpetologist who earned her undergraduate and Master's degrees at the Universidad Nacional Autónoma de México. For her undergraduate thesis, she worked on a herpetological survey of the municipality of Pluma Hidalgo, Oaxaca. Her Master's thesis focused on the molecular phylogeny of the *Sarcohyala bisticta* group (Anura: Hylidae). Itzue currently is a third year Ph.D. student in the Leache Lab at the University of Washington, and her dissertation research is focused on the systematics and evolution of the frog family Hylidae in Middle America.



Fausto R. Méndez-de la Cruz is a full professor at the Universidad Nacional Autónoma de México, in Mexico City. He received his Ph.D. in 1989 at the Universidad Nacional Autónoma de México, and conducted postdoctoral research at the University of Florida, and a sabbatical at Virginia Polytechnic Institute and Virginia State University, both in the United States. Fausto's research focuses on the evolution of reproductive systems in reptiles, mainly in the evolution of viviparity and parthenogenesis, and more recently in the vulnerability of amphibians and reptiles to climate change in Mexico and South America.



Adam D. Leaché is an Associate Professor at the University of Washington and the Curator of Genetic Resources and Herpetology at the Burke Museum of Natural History and Culture. He received his Ph.D. in 2008 at the University of California, Berkeley. The research in his lab focuses on topics including phylogenetics and species tree inference, statistical phylogeography, and species delimitation.