



## Can All Tiles of a Tiling Have Five-Fold Symmetry?

Ludwig Danzer; Branko Grunbaum; G. C. Shephard

*The American Mathematical Monthly*, Vol. 89, No. 8 (Oct., 1982), 568-570+583-585.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28198210%2989%3A8%3C568%3ACATOAT%3E2.0.CO%3B2-D>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# UNSOLVED PROBLEMS

EDITED BY RICHARD GUY

*In this department the MONTHLY presents easily stated unsolved problems dealing with notions ordinarily encountered in undergraduate mathematics. Each problem should be accompanied by relevant references (if any are known to the author) and by a brief description of known partial results. Manuscripts should be sent to Richard Guy, Department of Mathematics and Statistics, The University of Calgary, Calgary, Alberta, Canada T2N 1N4.*

## CAN ALL TILES OF A TILING HAVE FIVE-FOLD SYMMETRY?\*

LUDWIG DANZER

*Department of Mathematics, Universität Dortmund, 4600 Dortmund 50, West Germany*

BRANKO GRÜNBAUM\*

*Department of Mathematics, University of Washington GN-50, Seattle, WA 98195*

G. C. SHEPHARD

*University of East Anglia, Norwich NR4 7TJ, England*

It is well known that the plane can be tiled by equilateral triangles, by squares, or by regular hexagons, but that there exists no such tiling by regular pentagons. More generally it is not hard to see that the plane cannot be tiled by congruent figures each with 5-fold symmetry, that is, admitting as symmetries rotations about a point through angles of  $72^\circ$ ,  $144^\circ$ ,  $216^\circ$  and  $288^\circ$ . However, if one does not insist that the tiles are congruent, and one allows a variety of different shapes and sizes, then the question of tiling by tiles with 5-fold symmetry becomes much more interesting and leads to several open problems.


We recall that one of the basic results in geometric crystallography is the *crystallographic restriction*, which asserts that in the Euclidean plane no figure with a discrete symmetry group can have more than one center of five-fold rotational symmetry. Similar assertions can be made about  $k$ -fold rotations with  $k \geq 7$ . These facts lead to the well-known result that if a planar figure admits translational symmetries but does not admit arbitrarily short translations as symmetries, then the only possible rotational symmetries are  $k$ -fold rotations with  $k = 2, 3, 4$  or  $6$  (see, for example, Buerger [3, p. 33], Coxeter [5, Sect. 4.5], Fejes Tóth [9, Sect. I.1.4]).

However, the crystallographic restriction does not prove the impossibility of a tiling in which each individual tile has 5-fold symmetry. Fruitless efforts by mathematicians to construct such tilings go back at least to Kepler [11]. Fig. 1 shows some of Kepler's attempts; he notes that in each case a tiling can be obtained only at the price of introducing "monsters" (such as the "fused decagons" in Fig. 1a) which do not have 5-fold symmetry. Explanations of Kepler's tilings and variants of them have been discussed by several authors (Caspar [4, p. 374], Bindel [1], Eberhart [7], Grünbaum & Shephard [10, Sect. 2.5]). Even earlier attempts at constructing tilings with tiles that have 5-fold symmetry are discernible in Islamic art (see, for example, Bourgoïn [2], Critchlow [6], Wade [14], El-Said & Parman [8]).


It is therefore somewhat surprising to find that such tilings can easily be constructed by a simple inductive procedure. In the example indicated in Fig. 2, the tiles used are congruent either to the pentagon A shown in Fig. 2a, or to the nonconvex 20-sided polygon B shown in Fig. 2b, or to a polygon  $3^n B$  similar to B in ratio  $3^n$ , where  $n = 1, 2, 3, \dots$ . The construction of the tiling, indicated in Fig. 2c, is based on the fact that five copies of A yield with B a pentagon  $3A$  similar to A in ratio 3; five of these and  $3B$  yield a pentagon  $3^2 A$ , etc.

---


\*Research supported by the National Science Foundation Grant MCS77-01629 A01 and a Guggenheim Fellowship.



(a)



(b)



(c)

FIG. 1. Some of the patches devised by Kepler in 1619 in attempts to tile the plane by tiles with 5-fold symmetry.

The tiles in this example are not uniformly bounded in diameter. It seems that the answer to the question changes if the diameters of the tiles cannot grow without bounds. To make this problem precise from now on we restrict attention to tilings in which the tiles are closed topological disks of diameter at most 1, covering the plane without gaps or overlaps of their interiors. We conjecture that *the Euclidean plane admits no tiling in which each tile has five-fold symmetry*. If true, this conjecture is precariously balanced against various indications that seem to lend support to the opposite view. For example, tilings by congruent regular pentagons are possible on the sphere, in the elliptic plane and in the hyperbolic plane. The plane can be tiled by affinely regular pentagons, all mutually similar and of only two sizes (see Fig. 3). Many kinds of equilateral convex pentagons—some very close in shape to regular pentagons—can be used to tile




FIG. 2. The construction of a tiling (c) in which each tile is congruent either to the regular pentagon in (a), or to the 20-sided polygon in (b), or to an enlarged version of this polygon.




FIG. 3. A tiling of the plane by affinely regular pentagons, all mutually similar and of only two sizes.

the plane by congruent copies (see the surveys in Schattschneider [12], [13]). Probably even more telling are examples like the one in Fig. 4, which *appears* to show a tiling in which each tile is a regular pentagon. Actually, the pentagons in Fig. 4 form a Cantor-type set, and do not cover the plane; hence they do not form a tiling in the sense considered here.

Large regions of the plane can be covered, in many ways, by *patches* of tiles, in which each tile has 5-fold symmetry or even reflective 5-fold symmetry. In order to compare in a meaningful way the sizes of such patches, various measures can be used. Possibly the most appropriate one is the ratio  $\rho$  of the diameter of the largest circular disk covered by the patch to the diameter of the smallest circular disk that can cover each of the tiles in the patch. By adding around Kepler's patch in Fig. 1b ten of the larger pentagons and fifteen of the smaller ones, a patch with  $\rho \approx 3.1$  is




FIG. 4. A tiling which appears to consist of regular pentagons (the triangular spaces being “filled” by smaller and smaller pentagons). However, it can be shown that the union of the pentagons is a set of Cantor type, which does not cover the plane.

obtained. Similarly, the patch in Fig. 1c can be extended (by adding ten pentagons) to a patch with  $\rho \approx 2.6$ , while deleting from Fig. 1a the tiles which do not have 5-fold symmetry leads to a patch with  $\rho \approx 3.9$ .

The largest known patch made up of tiles which are regular pentagons, decagons, and pentagrams is shown in Fig. 5. Here  $\rho = \sqrt{(83 + 37\sqrt{5})}/8 \approx 4.6$ . Patches with larger values of  $\rho$  can be constructed using the process of “decomposition”—a method which has proved useful in other tiling problems (see, for example, [10, Chapter 10]). This may be explained as follows. Let the edge-length of the small pentagons in Fig. 5 be denoted by  $s$ , and let each vertex of this patch be the center of a regular decagon with edge-length  $s/\tau^3$ , where  $\tau = \frac{1}{2}(1 + \sqrt{5})$  is the golden section ratio. Then the parts of the tiles outside these decagons can be partitioned into smaller tiles, each with 5-fold symmetry. In Fig. 6 we show this process applied to (one tenth of) the patch of Fig. 5. To obtain the whole patch we have to mirror the sector shown in the dotted lines and




FIG. 5. A patch with the largest known  $\rho \approx 4.55157\dots$  among patches consisting of regular pentagons, decagons and pentagrams. The largest circle covered by the patch passes through the point marked by an asterisk and through its nine homologues.

their images. The original (larger) tiles are indicated by thin lines, and the tiles obtained by the decomposition process are shown by thicker lines.

Decomposition increases  $\rho$  for two reasons: the diameters of the tiles are decreased and also the patch can be extended a little using the smaller tiles. The patch in Fig. 6 has  $\rho \approx 8.6$ .

Repeating the process of decomposition leads to patches with larger values of  $\rho$ ; in this way we can obtain patches with up to  $\rho \approx 13$ . Using other starting patches, we can get as high as  $\rho \approx 38$ . We do not know whether this is anywhere near the maximum; indeed, we cannot even prove the existence of a value  $\omega < \infty$  such that  $\rho \leq \omega$  for each patch of the kind under consideration. We conjecture that no such  $\omega$  exists.

Similar problems can be raised concerning tilings and patches in which each tile has  $k$ -fold symmetry for some  $k \geq 7$ , or for any combination of such tiles and tiles with 5-fold symmetry.

#### References

1. E. Bindel, *Harmonien im Reiche der Geometrie in Anlehnung an Keplers "Weltharmonik,"* Verlag Freies Geistesleben, Stuttgart, 1964.
2. J. Bourgoïn, *Les éléments de l'art arabe: le trait des entrelacs,* Firmin-Didot et Cie., Paris, 1879. Paperback reprint of the diagrams: *Arabic Geometrical Pattern and Design,* Dover, New York, 1973.
3. M. J. Buerger, *Elementary Crystallography,* Wiley, New York, 1956.
4. M. Caspar, *Welt-Harmonik,* Oldenbourg, München, 1939.
5. H. S. M. Coxeter, *Introduction to Geometry,* Wiley, New York, 1961.
6. K. Critchlow, *Islamic Patterns, An Analytical and Cosmological Approach,* Schocken Books, New York, 1976.
7. S. Eberhart, *New and old problems,* *Mathematical-Physical Correspondence*, 12(1975) 4–5; *ibid.* 13(1975) 22–24.




FIG. 6. A patch with  $\rho \approx 8.6$  obtained by “decomposition” from the tiling in Fig. 5 and addition of some tiles around the boundary.

8. I. El-Said and A. Parman, *Geometric Concepts in Islamic Art*, World of Islam Festival Publishing Co., London, 1976.

9. L. Fejes Tóth, *Reguläre Figuren*, Akadémiai Kiadó, Budapest, 1965. English translation: *Regular Figures*, Pergamon, New York, 1964.

10. B. Grünbaum and G. C. Shephard, *Tilings and Patterns*, Freeman, San Francisco, 1982.

11. J. Kepler, *Harmonice Mundi*. Lincii, 1619. Reprints: Culture et Civilisation, Bruxelles, 1968; Johannes Kepler Gesammelte Werke, M. Caspar, editor, Band VI, Beck, München, 1940. Annotated German translation: M. Caspar, *Welt-Harmonik*, Oldenbourg, München, 1939. An annotated English translation of Book II appears in: J. V. Field, Kepler's star polyhedra, *Vistas in Astronomy*, 23(1979) 109–141.

12. D. Schattschneider, Tiling the plane with congruent pentagons, *Math. Mag.*, 51(1978) 29–44.

13. ———, In praise of amateurs, *The Mathematical Gardner*, edited by D. A. Klarner, Prindle, Weber & Schmidt, Boston, 1981, pp. 140–166 + 5 plates in color.

14. D. Wade, *Pattern in Islamic Art*, Overlook Press, Woodstock, NY, 1976.

---

### MISCELLANEA

83. Mathematics, like Dialectics, is an organ of the inner, higher mind; in practice, it is an art like eloquence. In both, nothing counts but the form; the content is irrelevant.

—J. W. v. Goethe, *Maximen und Reflexionen*, no. 605.