

Session 19

SILICONE WRINKLING MEMBRANES

Traction Forces

- ◆ Previously Paradigm
 - ◆ First cells cultured inside plasma clots (a/k/a fibrin gels)
 - ◆ Widely held belief that cells cause gels to shrink by dehydration
- ◆ Albert Harris worked under Michael Abercrombie
 - ◆ Cell-gel distortions were “side-effect” of propulsion forces
 - ◆ Sought to develop flexible gels to map cell forces
 - ◆ Difficult to get funding





Albert Harris

Flexible Substrata


- ◆ Particle movement
 - ◆ Carbon black particles (soot) mixed in thin layer of plasma clot
 - ◆ Centripetal movement of carbon particles
 - ◆ Difficult to maintain uniform Young's modulus
 - ◆ Many grant applications rejected
- ◆ Cross-linked silicone fluids
 - ◆ Flame-cured flexible skin on silicone fluid covered coverslip glass
 - ◆ Silicone impervious to hydration/dehydration effects
 - ◆ Received tenure on year before *Science* break-through

Wrinkling

- ◆ Compression folds underneath cell
- ◆ Tension wrinkles radiate outward
- ◆ UV treatment increases wrinkling by weakening cross-links in silicone film






Wrinkling Video




Calibration

- ◆ Pulled glass needle
 - ◆ Spring constant of needle calibrated with hanging weights
 - ◆ Pushing force applied to fixed cells on sheet
 - ◆ Force causes reversible wrinkles
 - ◆ Linear relationship between wrinkle length and applied force



Confirmed Rho/Myosin involvement




LPA
activates
Rho

BDM
inhibits
myosin
ATPase
activity

KT5926
inhibits
MLCK

Calmodulin/Myosin Involvement


- ◆ Transfect cells with cDNA constructs
- ◆ Caldesmon inhibits calmodulin, actin, and myosin activity
- ◆ CD445B is truncated caldesmon without actin, calmodulin, & myosin binding sites
- ◆ GFP construct used as control



Helfman, *et al.* (1999) *Mol Biol Cell*, 10:3097

Myofibroblast Differentiation


- ◆ Fibroblasts expressing α -smooth muscle actin generate large traction forces
- ◆ Contractile differentiation important for wound healing



Hinz, B., et al. (2001) *Mol Biol Cell*, 12:2730

Impact of Harris' work

- ◆ Direct observation of small, weak forces not possible before (and strange to some)
- ◆ Technique is not easily reproduced
- ◆ Not a direct quantitative approach
- ◆ Cell force techniques improve on reproducibility and quantification



(MBOC) Figure 19-50. The shaping of the extracellular matrix by cells. This micrograph shows a region between two pieces of embryonic chick heart (rich in fibroblasts as well as heart muscle cells) that were cultured on a collagen gel for 4 days. A dense tract of aligned collagen fibers has formed between the explants, presumably as a result of the fibroblasts in the explants tugging on the collagen. (From D. Stopak and A.K. Harris, *Dev. Biol.* 90:383–398, 1982)