

Posted: August 28, 2007

Enclosures Posted: Sep 27th, 2007

from Lab to Fab Posted: Sep 27th, 2007

other structures

single electrons

Posted: Sep 27th, 2007

Scientists spot sneaky 'neurodegenerative' iron at the

European synchrotron Posted: Sep 27th, 2007

Sequencing Platfor(...) Posted: Sep 27th, 2007

Association's Sports

Posted: Sep 27th, 2007

Posted: Sep 27th, 2007

Posted: Sep 27th, 2007

multifunctionality Posted: Sep 27th, 2007

Posted: Sep 27th. 2007

nanoparticles

Posted: Sep 27th, 2007

Commercializing Nanotechnology

Using nanotubes to detect and

repair cracks in aircraft wings,

Quantum device traps, detects

and manipulates the spin of

BioNanoatrix and Complete

NIST-ATP Award to Develop

IDGLOBAL Receives Second

Apparel/Merchandising Market

Nanoparticle could help detect

cancer and other diseases early

Carbon nanotubes have room for

...more news

Gold nanorods image tumors

Novel 3-D cell culture model shows selective tumor uptake of

Order from a Major League

Genomics Receive \$8.8 Million

Nanoscale system to measure cellular forces

(*Nanowerk News*) University of Pennsylvania researchers have designed a nanoscale system to observe and measure how individual cells react to external forces.

By combining microfabricated cantilevers and magnetic nanowire technology to create independent, nanoscale sensors, the study showed that cells respond to outside forces and demonstrated a dynamic biological relationship between cells and their environment.

The study also revealed that cells sense force at a single adhesion point that leads not to a local response but to a remote response from the cell's internal forces, akin to tickling the cell's elbow and watching the knee kick.

"The cell senses the force that we apply and adjusts its own internal forces to compensate," Chris Chen, an associate professor in the Department of Bioengineering in the School of Engineering and Applied Science at Penn, said. "This suggests that either the cell's cytoskeleton dictates the reaction or the cell organizes a biochemical response. In either instance, cells are adapting at the microscale."

The findings prove useful to more than just an understanding of the mechanics of single cells. Physical forces play a strong role in how whole tissue grows and functions. Using the Penn system, researchers could monitor for differences in how forces are sensed or generated in normal and diseased cells. This could lead to new therapeutic drug targets and to methods for modifying how cells interact with each other.

To study the cell's biomechanical response to forces, Chen and his team applied force to each cell using microfabricated arrays of magnetic posts containing cobalt nanowires interspersed amongst an array of non-magnetic posts. In the magnetic field, the posts with nanowires applied an external force to cells cultured on the tops of the posts. Nonmagnetic posts acted as sensors in which traction forces in each cell were measured. Recording the traction forces in response to such force stimulation revealed two responses: a sudden loss in contractility that occurred within the first minute of stimulation or a gradual decay in contractility over several minutes.

For both types of responses, the subcellular distribution of loss in traction forces was not confined to locations near the actuated micropost or uniformly across the whole cell but instead occurred at discrete locations along the cell periphery. Together, these data suggest that cells actively adjust their internal tension to mechanical forces arising in their microenvironment and reveal an important dynamic biological relationship between external and internal forces.

Mechanical forces contribute to many cellular functions, including changes in gene expression, proliferation and differentiation. Applying shear or tensile stresses to cells in culture, for example, can induce changes in adhesion regulation, intracellular signaling and cell function much like internal forces do. The similarities in cellular responses to external and internal forces have led to the suggestion that both types of forces may use shared mechanotransduction pathways to convert mechanical stimuli into biochemical signals. While externally applied and internally generated forces may act independently on cells, the University of Pennsylvania team postulated and then showed that they are coupled.

The study was conducted by Chen, Nathan J. Sniadecki, Michael T. Yang and Zhijun Liu of the Department of Bioengineering as well as Alexandre Anguelouch, Corinne M. Lamb, Stuart B. Kirschner, Yaohua Liu and Daniel H. Reich of Johns Hopkins University.

Source: University of Pennsylvania

Army Ringtones

Send Complimentary Ringtones to your cell. BestTones4U.com

Next Generation AFM/SPMs

Advanced atomic force microscopes for nanotechnology applications www.AsylumResearch.com

Magnetic Nanoparticles

Diagnostic Particle Fluids for Bead Development, Separations & Assays www.Ferrotec.com

Ga Tech Nano Research Ctr

Microelectronics, Nanotechnology E-beam litho, Nano-imprinter www.mirc.gatech.edu

Strainsert Load Cells

Load Cells, Load Pins, Force Bolts, Instrumentation and Systems www.strainsert.com **EHS Research on Nanotech**

New ICF Study Frames U.S. Policy Issues & Describes Path Forward www.icfi.com/nano

Strainsert Load Cells Load Cells, Load Pins, Force Bolts, Instrumentation and

Systems www.strainsert.com

Nanoscale Science AFM

Nanoscale Bio, physics applications AFM tools for noncontact metrology www.parkafm.com

Army Ringtones

Send Complimentary Ringtones to your cell. BestTones4U.com

Ads by Google

Privacy Statement | Terms of Use | Contact Us | Nanotechnology Home | Sitemap | Top of Page

The contents of this site are copyright © 2005-2007, Nanowerk LLC. All Rights Reserved