


# 3D Integration

Recall the definition of the definite integral of a function of a single variable:


Let  $f(x)$  be defined on  $[a, b]$  and let  $x_0, x_1, \dots, x_n$  be a partition of  $[a, b]$ . For  $i = 1, 2, \dots, n$ , let  $x_i^* \in [x_{i-1}, x_i]$ . Then




$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x.$$

**Generalizing from one variable to two.**

$f(x)$  on an interval  $[a, b]$   
 $f(x_i^*) \Delta x$  little bit of area




$f(x, y)$  on rectangle  $R = [a, b] \times [c, d] =$   
 $f(x_i^*, y_j^*) \Delta x \Delta y$  little bit of volume




**Double Riemann Sum** If  $f(x, y) \geq 0$  the double Riemann sum approximates the volume under the surface.

$$V \approx \sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \underbrace{\Delta x \Delta y}_{\Delta A}$$


$$\iint_R f(x, y) dA = \lim_{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \Delta A$$



(a)  $n = 16$



(b)  $n = 64$



(c)  $n = 256$

1. Let  $R$  be the rectangle  $1 \leq x \leq 1.2$  and  $2 \leq y \leq 2.4$ . If the values for  $f(x, y)$  are as specified below, find a Riemann sum approximation for  $\iint_R f(x, y) dA$  with  $\Delta x = 0.1$  and  $\Delta y = 0.2$ . Determine if your sum is an over- or under-estimate.

$y \setminus x$	1.0	1.1	1.2
2.0	5	7	10
2.2	4	6	8
2.4	3	5	6

2. Evaluate the integrals

(a)  $\int_0^3 \int_0^4 (4x + 3y) dx dy$

(b)  $\int_0^4 \int_0^3 (4x + 3y) dy dx$

(c)  $\int_1^3 \int_0^4 e^{x+y} dy dx$

(d)  $\int_0^4 \int_1^3 e^{x+y} dx dy$