

Three-particle scattering amplitudes from lattice QCD



Steve Sharpe
University of Washington



Collaborators




Max Hansen (Edinburgh)




Raúl Briceño (Berkeley)




Fernando Romero-López
(Bern)



Tyler Blanton (ex-UW)



Zach Draper (ex-UW)



Sebastian Dawid (Indiana)



Drew Hanlon (CMU)



Ben Hörz (Intel)




Sarah Skinner (ex-CMU)



Colin Morningstar (CMU)




Jorge Baeza Ballesteros
(Berlin)



Hans Bijmens (Lund)



Tomas Husek
(Birmingham/Prague)




Mattias Sjö (Marseille)



Wilder Schaaf (UW)


Overview: present status

Overview: present status



Physical quark masses


E.g.: Wang, Leinweber, Liu, Liu, Sun, Thomas, Wu, Xing, Yu,
2502.03700



$$M_{\pi} = 200 \text{ MeV}$$


Bulava, Hanlon, Hörz, Morningstar, Nicholson, Romero-López
Skinner, Varnas, Walker-Loud, 2208.03867

Overview: present status




Physical quark masses

E.g.: Wang, Leinweber, Liu, Liu, Sun, Thomas, Wu, Xing, Yu,
2502.03700



$$M_{\pi} = 200 \text{ MeV}$$

Bulava, Hanlon, Hörz, Morningstar, Nicholson, Romero-López
Skinner, Varnas, Walker-Loud, 2208.03867




Almost “FLAG-ready”, i.e. fully controlled


Abbott et al. [RBC-UKQCD collaboration], 2004.09440

Δ resonance


Bulava, Hanlon, Hörz, Morningstar, Nicholson, Romero-López, Skinner, Varnas, Walker-Loud, 2208.03867



$M_\pi \approx 200$ MeV
 $M_N \approx 950$ MeV
 $a = 0.063$ fm
 $L^3 \times T = 64^3 \times 128$
Clover fermions (CLS)




Overview: present frontier



Physical quark masses

Dawid, Draper, Hanlon, Hörz, Morningstar, Romero-López, SRS, Skinner, [2502.14348](#) & [2502.17976](#)




$$M_{\pi} = 200 \text{ MeV}$$


Padmanath, Prelovsek, [2202.10110](#)

...

Dawid, Romero-López, SRS, [2409.17059](#)



Formalism: Hansen, Romero-López, SRS, [2509.24778](#)




Formalism: Müller, Rusetsky, [2012.13957](#)

Hansen, Romero-López, SRS, [2101.10246](#)


Müller, Pang, Rusetsky, Wu, [2211.10126](#)

Pang, Bubna, Müller, Rusetsky, Wu, [2312.04391](#)

Overview: present frontier




Formalism: Hansen, Romero-López, SRS, 2003.10974
Alternative formalism & first LQCD results:
Yan, Mai, Garofalo, Meißner, Liu, Liu, Urbach,
2407.16659; Yan, Mai, Garofalo, Feng, Döring, Liu, Liu,
Meißner, Urbach, 2510.0947




Formalism: Hansen, Romero-López, SRS, 2003.10974;
Jackura, 2208.10587
First LQCD results:
Briceño, Hansen, Jackura, Edwards, Thomas, 2510.24894


Overview: present frontier




Formalism: Hansen, Romero-López, SRS, 2003.10974
 Alternative formalism & first LQCD results:
 Yan, Mai, Garofalo, Meißner, Liu, Liu, Urbach,
 2407.16659; Yan, Mai, Garofalo, Feng, Döring, Liu, Liu,
 Meißner, Urbach, 2510.0947



Formalism: Hansen, Romero-López, SRS, 2003.10974;
 Jackura, 2208.10587
 First LQCD results:
 Briceño, Hansen, Jackura, Edwards, Thomas, 2510.24894




Di-neutron bound state?



Is H dibaryon bound?


Overview: long-term dreams

Roper resonance





Formalism: Hansen, Romero-López, SRS, ... in prep.

3 neutron interactions



Formalism: Draper, Hansen, Romero-López, SRS, 2303.10219
Toy application: Schaaf, SRS, 2512.24508

CP violation in D decays



Outline

- Motivations
- The fundamental issue & its resolution
- Recent applications of three-particle formalism
 - Three-particle amplitudes involving pions & kaons at physical quark masses
 - $DD\pi$ scattering, relevant for the T_{cc}^+
 - Progress on $N\pi\pi$ — towards the Roper resonance
 - Dreaming of three neutrons
- Summary & Outlook

Motivations

Underlying motivations

- Determine properties of strong interaction resonances from QCD
 - E.g. exotics such as $T_{cc}(3875)^+ \rightarrow DD^* \rightarrow DD\pi$
 - Long-standing puzzles, e.g. Roper $N(1440) \rightarrow N\pi + N\pi\pi$


Underlying motivations

- Determine properties of strong interaction resonances from QCD
 - E.g. exotics such as $T_{cc}(3875)^+ \rightarrow DD^* \rightarrow DD\pi$
 - Long-standing puzzles, e.g. Roper $N(1440) \rightarrow N\pi + N\pi\pi$
- Determine three particle “forces” for $3n$, 3π , $3K$, ...
 - Needed to understand neutron star EoS, properties of large nuclei, ...

Underlying motivations

- Determine properties of strong interaction resonances from QCD
 - E.g. exotics such as $T_{cc}(3875)^+ \rightarrow DD^* \rightarrow DD\pi$
 - Long-standing puzzles, e.g. Roper $N(1440) \rightarrow N\pi + N\pi\pi$
- Determine three particle “forces” for $3n$, 3π , $3K$, ...
 - Needed to understand neutron star EoS, properties of large nuclei, ...
- Calculate weak decay amplitudes within the Standard Model, in order to search for new physics
 - E.g. $K \rightarrow 2\pi$ (essentially done), $K \rightarrow 3\pi$ (method known), & $D \rightarrow \pi^+\pi^-$, K^+K^- (open question)

Cornucopia of exotics



[I. Danilkin, talk at INT workshop, March 23]

Data from Babar, Belle, BESIII, COMPASS, LHCb

The fundamental issue & its resolution

On the one hand...

- LQCD determines energies and properties of finite-volume eigenstates
 - Obtained by fits to (numerically-evaluated) Euclidean correlation functions:


$$\int_L d^3x e^{-i\vec{P}\cdot\vec{x}} \langle \Omega | \sigma_{3\pi}(\tau, \vec{x}) \sigma_{3\pi}^\dagger(0) | \Omega \rangle_L \propto \sum_n \left| \langle 0 | \sigma_{3\pi}^\dagger(0) | 3\pi, \vec{P}, n \rangle_L \right|^2 e^{-E_n\tau}; \quad (\tau > 0)$$

$\vec{P} = 2\pi\vec{n}/L$
Assuming L^3 box with PBC

$\sigma_{3\pi} \sim 3\pi^+$
Lives on timeslice

Tower of finite-volume states
with quantum numbers of $3\pi^+$,
with momentum \vec{P} , and living
in irreps of cubic group

Energies of said states



On the one hand...


- LQCD determines energies and properties of finite-volume eigenstates
 - Obtained by fits to (numerically-evaluated) Euclidean correlation functions:

$$\int_L d^3x e^{-i\vec{P}\cdot\vec{x}} \langle \Omega | \sigma_{3\pi}(\tau, \vec{x}) \sigma_{3\pi}^\dagger(0) | \Omega \rangle_L \propto \sum_n \left| \langle 0 | \sigma_{3\pi}^\dagger(0) | 3\pi, \vec{P}, n \rangle_L \right|^2 e^{-E_n\tau}; \quad (\tau > 0)$$

- E_n are physical quantities!
 - Can determine 5-10 levels for each choice of quantum numbers (\vec{P} , irrep, ...)
 - Can now begin to calculate with physical quark masses
 - Results come with statistical & systematic errors (e.g. need $a \rightarrow 0$)
 - Today, I assume that the physical E_n are provided by LQCD simulations


...while on the other

- We want infinite-volume scattering amplitudes, e.g.




...while on the other

- We want infinite-volume scattering amplitudes, e.g.



- How do we relate these? A finite-volume QFT problem.



A related question:


- LQCD can also calculate matrix elements between finite-volume states

$${}_L\langle \Omega | \sigma_{3\pi}(\tau_f, \vec{P}) \left[\int_L d^3x \mathcal{H}_W(0, \vec{x}) \right] K^\dagger(\tau_i, \vec{P}) | \Omega \rangle_L \propto \sum_{n', n} c_{n', n} e^{-E_n \tau_f} \underbrace{{}_L\langle 3\pi, \vec{P}, n' | \mathcal{H}_W(0) | K, n \rangle_L}_{\text{A physical quantity if } E_{n'} = E_n} e^{E_{K_n} \tau_i}$$

$\tau_f > 0$ (indicated by a red arrow pointing to τ_f)
 $\tau_i < 0$ (indicated by a red arrow pointing to τ_i)


2-particle formalism

[Lüscher, 1986-91 + many subsequent works]




2-particle formalism

[Lüscher, 1986-91 + many subsequent works]




2-particle formalism

[Lüscher, 1986-91 + many subsequent works]



3-particle formalism

[Hansen & SRS, 2014 & 2015 + many subsequent works]






- Using more data than QC2, but fitting much more challenging
- More quantities to parametrize, with limited phenomenological knowledge for $\mathcal{K}_{df,3}$
- Solving integral equations involves new challenges

QC3

$$\det_{k\ell m} \left[F_3^{-1} + \mathcal{K}_{\text{df},3} \right] = 0$$

- QC3 valid up to corrections $\propto \exp(-M_\pi L)$
- QC3 valid up to first inelastic threshold, e.g. $E^* = 5M_\pi$ for 3 pion system
- Matrix indices are spectator finite-volume momentum k , and pair CM-frame ℓ, m
- RFT formalism includes smooth cutoff in k ; must truncate “by hand” in ℓ

$$F_3 = \frac{1}{2\omega L^3} \left[\frac{F}{3} - F \frac{1}{1/\mathcal{K}_{2,L} + F + G} F \right]$$



Relating $\mathcal{K}_{\text{df},3}$ to \mathcal{M}_3

$$\mathcal{M}_3 = \lim_{L \rightarrow \infty} \mathcal{S} \left\{ \mathcal{D}_L^{(u,u)} + \mathcal{M}_{\text{df},3,L}^{(u,u)} \right\} = \mathcal{D} + \mathcal{M}_{\text{df},3}$$

- \mathcal{D} contains all divergent contributions to \mathcal{M}_3 , but depends on cutoff function $H(\vec{k})$


$$\mathcal{D} = \mathcal{S} \left\{ \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} \overset{G^\infty}{\text{---}} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} \overset{G^\infty}{\text{---}} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} \overset{G^\infty}{\text{---}} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} + \dots \right\}$$

- $\mathcal{M}_{\text{df},3}$ is divergence-free, equals $\mathcal{K}_{\text{df},3}$ at leading order, and is also cutoff-dependent

$$\mathcal{M}_{\text{df},3} = \mathcal{K}_{\text{df},3} + \mathcal{S} \left\{ \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{K}_{\text{df},3} \end{array} \overset{\rho}{\text{---}} \begin{array}{c} \circlearrowleft \mathcal{K}_{\text{df},3} \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \circlearrowleft \mathcal{K}_{\text{df},3} \\ \text{---} \\ \circlearrowleft \mathcal{K}_{\text{df},3} \end{array} \overset{\rho}{\text{---}} \begin{array}{c} \circlearrowleft \mathcal{M}_2 \\ \text{---} \\ \circlearrowleft \mathcal{M}_2 \end{array} + \dots \right\}$$


- “Decorations” ensure that \mathcal{M}_3 is unitary

Status: formalism




[Hansen & SRS 14,15 (RFT); Hammer, Pang, Rusetsky 17 (NREFT); Mai, Döring 17 (FVU)]

[Many subsequent refinements & tests]




[Hansen, Romero-López, SRS (RFT) 20;

Mai, Alexandru, Brett, Culver, Döring, Lee, Sadasivan (FVU) 21; Feng, Gil, Döring, Molina, Mai, Shastry, Szczepaniak (FVU) 24; Yan, Mai, Garofalo, Meißner, Liu, Liu, Urbach (FVU) 24; Yan, Mai, Garofalo, Feng, Döring, Liu, Liu, Meißner, Urbach (FVU) 25]




[Blanton, Sharpe (RFT) 20, 21;


Feng et al. (FVU) 24]




[Draper, Hansen, Romero-López, SRS 23]




[Hansen, Romero-López, SRS 24]



[Draper, SRS 24]




[Briceño, Hansen, SRS 17]




[Hansen, Romero-López, SRS 25]


Status: solving integral equations





[Hansen, Briceño, Edwards, Thomas, Wilson (RFT) 20]




[Jackura, Briceño, Dawid, Islam, McCarty, 20;
Dawid, Islam, Briceño (RFT) 23;
Dawid, Islam, Briceño, Jackura (RFT) 23]



[Mai, Alexandru, Brett, Culver, Döring, Lee, Sadasivan (IVU) 21;
Jackura, Briceño, 23; Briceño, Costa, Jackura (RFT) 24; Jackura, Chambers,
Briceño (RFT) 25; Briceño, Hansen, Jackura, Edwards, Thomas (RFT) 25;
Yan, Mai, Garofalo, Feng, Meißner, Liu, Liu, Urbach (IVU) 24;
Yan, Mai, Garofalo, Feng, Döring, Liu, Liu, Meißner, Urbach (IVU) 25]



[Dawid, Draper, Hanlon, Hörz, Morningstar, Romero-López, SRS, Skinner, 25]



[Dawid, Romero-López, SRS 24, 25]

Recent applications of three-particle formalism


QCD predictions for physical multimeson scattering amplitudes

Sebastian M. Dawid,¹ Zachary T. Draper,¹ Andrew D. Hanlon,² Ben Hörz,³ Colin Morningstar,⁴ Fernando Romero-López,^{5,6} Stephen R. Sharpe,¹ and Sarah Skinner⁴

[2502.14348 (PRL) & 2502.17976 (PRD)]



Three PGBs at maximal isospin



$$3\pi^+, \quad 3K^+, \quad \pi^+\pi^+K^+, \quad K^+K^+\pi^+$$

- Benchmark system, with simple repulsive dynamics
 - First calculation at physical quark masses
 - Study $J^P = 0^-, 1^+, 2^-$, including subchannels with $\ell = 0, 1, 2$
 - Compare to expectations from ChPT
- Use CLS ensembles ($\mathcal{O}(a)$ improved Wilson) + GEVP


	$(L/a)^3 \times (T/a)$	M_π [MeV]	M_K [MeV]	N_{cfg}	$M_\pi L$
N203	$48^3 \times 128$	340	440	771	5.41
N200	$48^3 \times 128$	280	460	1712	4.42
D200	$64^3 \times 128$	200	480	2000	4.20
E250	$96^3 \times 192$	130	500	505	4.05

[Blanton, SRS, et al.,
PRL 2020 & JHEP 2021]
[Draper, SRS, et al.,
JHEP 2023]

New work

$$a \simeq 0.063 \text{ fm}$$

Three PGBs at maximal isospin



$$3\pi^+, \quad 3K^+, \quad \pi^+\pi^+K^+, \quad K^+K^+\pi^+$$

- Benchmark system, with simple repulsive dynamics
 - First calculation at physical quark masses
 - Study $J^P = 0^-, 1^+, 2^-$, including subchannels with $\ell = 0, 1, 2$
 - Compare to expectations from ChPT
- Use CLS ensembles ($\mathcal{O}(a)$ improved Wilson) + GEVP

	$(L/a)^3 \times (T/a)$	M_π [MeV]	M_K [MeV]	N_{cfg}	$M_\pi L$
N203	$48^3 \times 128$	340	440	771	5.41
N200	$48^3 \times 128$	280	460	1712	4.42
D200	$64^3 \times 128$	200	480	2000	4.20
E250	$96^3 \times 192$	130	500	505	4.05


[Blanton, SRS, et al.,
PRL 2020 & JHEP 2021]
[Draper, SRS, et al.,
JHEP 2023]

New work

$$a \simeq 0.063 \text{ fm}$$


$$\text{tr } m_q = 2m_{ud} + m_s \simeq \text{const}$$

Example of spectrum




E250: K^+K^+

Example of spectrum



E250: $K^+K^+\pi^+$

Fits to spectra




channels	# levels	# parameters	χ^2/DOF
$2\pi/3\pi$	34 + 32	6	1.14
$2\pi/\pi K/2\pi K$	24 + 25 + 23	9	1.23
$2K/\pi K/2K\pi$	25 + 40 + 50	10	1.95
$2K/3K$	40 + 53	6	1.49

Parameters in \mathcal{K}_2 and \mathcal{K}_3 : use threshold expansions


2-particle interactions

E250: \sim physical quark masses



Inelastic thresholds

Scattering lengths





$\chi^2/\text{dof} = 12.4/(11 - 2)$

- Simultaneous chiral fit to $\pi\pi$, πK , KK scattering lengths
 - Fit involves two LECs, one of which is determined with 2% stat. errors

Comparing $\mathcal{K}_{df,3}$ to ChPT

- Parametrize $\mathcal{K}_{df,3}$ in a threshold expansion, keeping first two terms
- From fits to spectra, we find non vanishing results for these terms at heavier quark masses
- Compare to NLO Chiral Perturbation theory calculation of $\mathcal{K}_{df,3}$ for $3\pi \rightarrow 3\pi$
[Baeza-Ballesteros, Bijens, Husek, Romero-López, SRS, Sjö, 2303.13206 (JHEP) & 2401.14293 (JHEP)]




Large NLO corrections in ChPT resolve LO inconsistency



Physical 3-particle amplitudes

$$M_\pi^4 |\text{Amplitude}|^2$$

$$J^P = 0^-$$



Divergent at threshold



Pion interactions are chirally suppressed

$$\mathcal{M}_3 = \mathcal{D} + \mathcal{M}_{\text{df},3}$$



$$\mathcal{D} = \mathcal{S} \left\{ \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \right\}$$

$$\mathcal{M}_{\text{df},3} = \mathcal{K}_{\text{df},3} + \mathcal{S} \left\{ \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} \right\}$$

Comparison with ChPT

$$M_\pi^4 |\text{Amplitude}|^2$$

$$J^P = 0^-$$



Observe expected breakdown of ChPT as M_π or p_π increase

Using three-particle methods to study the $T_{cc}^+(3875)$



Incorporating $DD\pi$ effects and left-hand cuts in lattice QCD studies of the $T_{cc}(3875)^+$

Maxwell T. Hansen,^a Fernando Romero-López ^b and Stephen R. Sharpe ^c


[2401.06609, JHEP]



Finite- and infinite-volume study of $DD\pi$ scattering

Sebastian M. Dawid ^a, Fernando Romero-López ^{b,c} and Stephen R. Sharpe ^a

[2409.17059, JHEP]



Comparison of integral equations used to study T_{cc}^+ for a stable D^*

Sebastian M. Dawid ^a, Fernando Romero-López ^b and Stephen R. Sharpe ^a

[2505.05466, JHEP]

+ work in progress

Doubly-charmed tetraquark

NATURE PHYSICS | VOL 18 | JULY 2022 | 751-754 | www.nature.com/naturephysics


Observation of an exotic narrow doubly charmed tetraquark

LHCb Collaboration*


$$T_{cc}^+(3875)$$

$$I = 0, J^P = 1^+$$


$$cc\bar{u}\bar{d}$$



Experiment



► For physical quark masses is a three-body resonance



need three-body formalism!

Adapted from: Fernando Romero-López

Doubly-charmed tetraquark

NATURE PHYSICS | VOL 18 | JULY 2022 | 751-754 | www.nature.com/naturephysics


Observation of an exotic narrow doubly charmed tetraquark

LHCb Collaboration*


$$T_{cc}^+(3875)$$

$$I = 0, J^P = 1^+$$


$$cc\bar{u}\bar{d}$$



Experiment




► For physical quark masses is a three-body resonance




need three-body formalism!

LQCD calculations with heavier-than-physical quarks



► D* is stable for slightly heavier-than-physical quarks



suitable for the two-body finite-volume formalism?

Adapted from: Fernando Romero-López

DD^* scattering

- Several LQCD studies using QC2 in T_{cc} channel

[Chen et al., 2206.06185]



[Lyu et al. (HALQCD), 2302.04505]

[Padmanath & Prelovsek, 2202.10110]

[Whyte, Thomas, Wilson, 2405.15741]

[Several talks at Lattice 2025]

- Use heavier-than-physical quarks
- Find signature of virtual bound state
- But QC2 fails at left-hand cut, below which phase shift becomes complex



- Several solutions to failure have been proposed, by generalizing the QC2


[Du et al (2408.09375); Abolnikov et al. (2407.04649), Bubna et al. (2402.12985);

Meng et al. (2312.01930); Raposo, Hansen (2311.18793); Raposo, Hansen, Briceño, Jackura (2502.19375)]

3-body solution

[Hansen, Romero-López, SRS, 2401.06609, JHEP]


- Left-hand cut is a three-body effect




3-body solution

[Hansen, Romero-López, SRS, 2401.06609, JHEP]

- Left-hand cut is a three-body effect




- Use $DD\pi$ QC3, but include D^* as a bound state in p -wave $D\pi$ channel




3-body solution

[Hansen, Romero-López, SRS, 2401.06609, JHEP]

- Left-hand cut is a three-body effect



- Use $DD\pi$ QC3, but include D^* as a bound state in p -wave $D\pi$ channel




- Finite-volume effects from u -channel pion exchange naturally incorporated

3-body solution

[Hansen, Romero-López, SRS, 2401.06609, JHEP]

- Left-hand cut is a three-body effect



- Use $DD\pi$ QC3, but include D^* as a bound state in p -wave $D\pi$ channel




- Finite-volume effects from u -channel pion exchange naturally incorporated
- Another advantage of QC3 approach:
 - Works for all choices of quark masses, including physical case of unbound D^*
- Successfully implemented in [Dawid, Romero-López, SRS, 2409.17059, JHEP & 2505.05466, JHEP; plus work in progress (talk by André Baião Raposo)]

Application to LQCD data

[Dawid, Romero-López, SRS, 2409.17059, JHEP]

- Proof of concept study to (limited) existing LQCD data with bound D^*
 - Choose p -wave $D\pi$ scattering amplitude to obtain D^* , for given choice of $g_{D^*D\pi}$
 - Choose reasonable s -wave $D\pi$ and DD amplitudes
 - Eyeball fit to results of [Padmanath, Prelovsek, arXiv:2202.10110] requires inclusion of $\mathcal{K}_{df,3}$
 - Solve integral equations, analytically continue to D^* pole, use LSZ to obtain $\mathcal{M}(DD^*)$
 - Result shows appropriate behavior at and below left-hand cut



- Virtual bound-state pole moves into complex plane!
- Looking forward: complete analysis requires more extensive LQCD spectra
 - DD , $D\pi$, $DD\pi$ & DD^* levels

Towards a formalism for the Roper

Finite-volume formalism for $N\pi\pi$ at maximal isospin


Maxwell T. Hansen^a, Fernando Romero-López^b, and Stephen R. Sharpe^c

[2509.24778 (to appear in JHEP)]



A failed hope for studying N(1440)

- Need a formalism that accommodates $N\pi + N\pi\pi$
 - Can one include N as a pole in p -wave $N\pi$ scattering, and use the $N\pi\pi$ formalism?
 - No! Nucleon pole lies below the singularity due to u -channel nucleon exchange




- Formalism for $N\pi\pi$ alone only valid at maximal isospin, which forbids mixing with $N\pi$
 - Interesting dynamics due to $\Delta\pi$ subchannel
 - Provides one 3-particle block of arbitrary isospin $N\pi + N\pi\pi$ formalism

Parameter space in QC3


QC3 for $N\pi\pi$:
$$\det_{i\mathbf{p}lmm_s} \left(1 + \hat{\mathcal{K}}_{\text{df},3}(E^*) \hat{F}_3(E, \mathbf{P}, L) \right) = 0$$

Matrix indices are:
channel, spectator momentum,
pair angular momentum,
lab frame nucleon spin

- A key quantity in derivation is B_3 , $3 \rightarrow 3$ Bethe-Salpeter kernel (3 Particle Irreducible)
 - Must be nonsingular in range of kinematic parameters
 - Otherwise introduce uncontrolled L^{-n} finite-volume effects




- Parameter space (for pion spectator)
 - Energy E , momentum \mathbf{P}



Subthreshold singularities

Parameter space for $P = 0$, $M_\pi/M_N = 0.2$



Dreaming of 3 neutrons

Three relativistic neutrons in a finite volume

Zachary T. Draper,^a Maxwell T. Hansen,^b Fernando Romero-López^c
and Stephen R. Sharpe^a

[2303.10219, JHEP]



Implementation of the three-neutron quantization condition


Wilder Schaaf^a and Stephen R. Sharpe^a

[2512.24508]




Spectrum in realistic setup


- Physical masses, $L \approx 4.3$ fm:
 - $M_N L = 20$, $M_\pi = 0.15 M_N \Rightarrow M_\pi L = 3$



Spectrum in realistic setup



Add leading-order $\mathcal{K}_{df,3}$ in threshold expansion
 Keep one term (out of two) with coefficient c_A



Summary & Outlook

Summary & Outlook

- Major steps have been taken in the three-particle sector
 - Formalism well established & cross checked, and almost complete
 - Several pioneering end-to-end applications to three-particle spectra from LQCD
 - Path to a calculation of $K \rightarrow 3\pi$ decay amplitudes is now open
- Near-term steps in implementation
 - $T_{cc}^+ \rightarrow D^*D \rightarrow DD\pi$ for lighter quark masses
 - Extended calculations for $3\pi(I = 0,1,2)$
 - $N\pi\pi \leftrightarrow \Delta\pi(I = 5/2)$
- Final steps in three-particle formalism
 - $3\pi(I = 0) \leftrightarrow K\bar{K}$ (WZW term)
 - $NNN(I = \frac{1}{2}), N\pi\pi + N\pi$ [for Roper], $NN\pi + NN$
- Very recent review—talk by SRS at Lattice 2025 [2601.04147]

Challenges & Future


- Reducing model-dependence in fitting & analytic continuation
 - [Salg, Romero-López, Jay, [2506.16161](#)] propose using Bayesian reconstruction + Nevanlinna interpolation
- Controlling all systematic errors (when will results be FLAG-ready?)
 - For which processes is precision important?
 - For three-nucleon interaction even a semiquantitative result would be useful
- Comparing formalisms in detail (e.g. RFT vs FVU)
 - Equivalent up to technical details (e.g. smooth vs hard cutoff functions)
 - Share data for cross-checking analysis? Bootcamp?
- Moving to more particles (needed, e.g., for $D \rightarrow \pi\pi, K\bar{K}$ & physical $\pi(1300)$)
 - QC4 (which is not yet known except in threshold expansion) may be practical
 - QCN seems impractical—combine QC2-4 with inclusive methods in some yet-to-developed manner?
 - “Scattering amplitudes from finite-volume spectral functions” [Bulava & Hansen, [1903.11735](#)]
 - Squeeze as much information as we can out of the lattice simulations!
- Investigate alternative approaches
 - “Scattering amplitudes from Euclidean correlators: Haag-Ruelle theory and approximation formulae”, [Patella & Tantalo, [2407.02069](#)]
 - HALQCD approach? [Doi et al., [1106.2276](#): 3 nucleon potentials in NR regime]
- Combine LQCD results with EFT and amplitude-analysis methods (Exohad collab.)

Thank you!
Questions?


Backup slides

Long-term dream

CP violation in D decays




Challenge: finite-volume mixing with $4\pi, 6\pi, \eta\eta, \dots$




4+-particle formalism not yet developed

Some angular dependence





Different partial waves (largest is $J = 0$)


Angle in the triangle:
equilateral ($\alpha = \pi/3$) vs isosceles



Angle between
in/out momenta



Three-meson spectrum



8 / 17

Fernando Romero-López, Uni Bern

3-particle resonances

Emergence of the $\pi(1300)$ Resonance from Lattice QCD

Haobo Yan (燕浩波),^{1,2,*} Maxim Mai,^{3,4,†} Marco Garofalo,^{2,‡} Yuchuan Feng,^{4,§} Michael Döring,^{4,¶}
Chuan Liu (刘川),^{1,5,6,**} Liuming Liu (刘柳明),^{7,8,††} Ulf-G. Meißner,^{2,9,10,‡‡} and Carsten Urbach^{2,§§}

[2510.09476, talk by Maxim Mai]

ω Meson from lattice QCD

Haobo Yan (燕浩波),^{1,2,*} Maxim Mai,^{2,3,4,†} Marco Garofalo,^{2,‡}
Ulf-G. Meißner,^{2,5,6,§} Chuan Liu,^{1,7,8,¶} Liuming Liu,^{9,10,**} and Carsten Urbach^{2,††}

[2407.16659, PRL]

Natural candidate for QC3?

$\pi(1300)$	$I^G(J^{PC}) = 1^-(0^{-+})$	[PDGLive]	
$\pi(1300)$ MASS	[1]	1300 ± 100 MeV	
$\pi(1300)$ WIDTH	[1]	200 to 600 MeV	
$\pi(1300)$ DECAY MODES			
Mode	Fraction (Γ_i / Γ)	Scale Factor/ Conf. Level	P(MeV/c)
Γ_1 $\rho\pi$	seen		404
Γ_2 $\pi(\pi\pi)_{S\text{-wave}}$	seen		
Γ_3 $\gamma\gamma$			650

[1] Our estimate. See the Particle Listings for details.

Pro: s -wave decay to 3π implies strong 3-particle coupling

Con: Lies well above the inelastic threshold for QC3


- $M(\pi(1300)) \approx 10M_\pi \gg 5M_\pi$
- QC3 will apply for heavier than physical quark masses

Simulation details


- Use CLQCD ensembles (tadpole improved Clover), fixed $m_s, a = 0.07746(18)\text{fm}$
 - Rest-frame only and single irrep

ens.	size	$M_\pi(\text{MeV})$	$M_K(\text{MeV})$	$M_\pi L$	# 2π levels	# 3π levels
F32P30	$32^3 \times 96$	303	525	3.8	5	3
F48P30	$48^3 \times 96$	303	524	5.7	7	6
F32P21	$32^3 \times 64$	211	492	2.7	5	0
F48P21	$48^3 \times 96$	207	493	3.9	7	3

- Large basis of operators: $\pi, \rho\pi, \sigma\pi, 3\pi$ for three-particle case
 - Large set of contractions, e.g.




Fits



- Best global fit to both masses and volumes
 - 9 params (3 for 2π , and 6 for 3π) to 36 levels, with $\chi^2/\text{dof} = 1.08$
 - Final results from model averaging with Akaike information criterion

Pole positions

- Solve IVU (=infinite-volume unitarity) form of integral equations
 - Find resonance pole for $M_\pi = 305 \text{ MeV}$!



Pole positions

- Solve IVU (=infinite-volume unitarity) form of integral equations
 - Find resonance pole for $M_\pi = 305 \text{ MeV}$!
- Extrapolate to physical masses using EFT-inspired form

