

Revisiting Jablonski (1993): cladogenesis and range expansion explain latitudinal variation in taxonomic richness

P. R. MARTIN,*†‡ F. BONIER* & J. J. TEWKSBURY*

*Department of Biology, University of Washington, Seattle, WA, USA

†Museo Ecuatoriano de Ciencias Naturales, Quito, Ecuador

‡Yanayacu Biological Station, Cosanga, Napo, Ecuador

Keywords:

biodiversity;
biogeography;
cladogenesis;
latitude;
latitudinal gradients;
range expansion;
taxonomic richness.

Abstract

The increase in diversity towards the equator arises from latitudinal variation in rates of cladogenesis, extinction, immigration and/or emigration of taxa. We tested the relative contribution of all four processes to the latitudinal gradient in 26 marine invertebrate orders with extensive fossil records, examined previously by David Jablonski. Coupling Jablonski's estimates of latitudinal variation in cladogenesis with new data on patterns of extinction and current distributions, we show that the present-day gradient in diversity is caused by higher rates of cladogenesis and subsequent range expansion (immigration) at lower latitudes. In contrast, extinction and emigration were not important in the creation of the latitudinal gradient in ordinal richness. This work represents one of the first simultaneous tests of the role of all four processes in the creation of the latitudinal gradient in taxonomic richness, and suggests that low tropical extinction rates are not essential to the creation of latitudinal diversity gradients.

Introduction

Biodiversity increases dramatically from the poles to the equator (Gaston, 2000; Willig *et al.*, 2003; Hillebrand, 2004). This latitudinal gradient in taxonomic richness is caused by one or more abiotic factors that vary over latitude (Francis & Currie, 2003; Hawkins *et al.*, 2003; Currie *et al.*, 2004; Jetz *et al.*, 2004; Ricklefs, 2004). From a mechanistic perspective, abiotic factors produce the gradient in diversity by creating latitudinal variation in rates of cladogenesis, extinction, immigration and/or emigration, the latter two processes resulting from range expansion, contraction and shifting.

A higher rate of tropical cladogenesis has been suggested by both direct and indirect evidence (Jablonski, 1993; Flessa & Jablonski, 1996; Cardillo, 1999; Martin & McKay, 2004; Cardillo *et al.*, 2005; Goldberg *et al.*, 2005; Allen & Gillooly, 2006; Jablonski *et al.*, 2006); however, extinction rates have been described as both higher and

lower in the tropics (Stanley, 1984; Jablonski, 1985; Jablonski *et al.*, 1985, 2006; Coope, 1995, 2004; Flessa & Jablonski, 1996; Bennett, 1997; Goldberg *et al.*, 2005; see also Gaston & Blackburn, 1996; Chown & Gaston, 2000; Chek *et al.*, 2003; Pimm & Brown, 2004). Few studies have examined latitudinal variation in rates of immigration and emigration outside the cyclic expansion, retraction and shifting of taxa during recurrent Milankovitch (glacial) cycles (Silvertown, 1985; Haskell, 2001; Harrington, 2004; see Jablonski *et al.*, 2006 for a notable exception).

Determining the extent to which each of the four processes (cladogenesis, extinction, immigration, emigration) are involved in creating the latitudinal gradient in taxonomic richness would provide a major advance in our understanding of the causes of latitudinal variation in diversity, as it would distinguish the mechanisms by which abiotic factors cause the gradient. Previous work by Jablonski (1993) on 26 benthic marine invertebrate orders is widely cited as evidence for higher rates of tropical cladogenesis. Here we revisit this data set to examine the role of all four processes in the creation of the present-day latitudinal gradient in these 26 marine invertebrate orders. Given that the latitudinal increase in

Correspondence: Paul Martin, Department of Biology, Queen's University, Kingston, Ontario, Canada K7L 3N6.

Tel.: +001 613 830 4801; fax: +001 613 533 6617; e-mail: martinp@biology.queensu.ca

taxonomic diversity towards the equator is evident across species, genera, families and orders (Willig *et al.*, 2003), the causes of latitudinal variation in ordinal richness are probably generalizable to other taxonomic levels (see also Sepkoski, 1999 for evidence showing coincident patterns of orders and species in the fossil record).

Methods

Present-day distributions

We documented the present-day latitudinal distribution of each of the 26 marine invertebrate orders using published literature (Appendix 1). For each order, we recorded presence or absence within 10° latitudinal belts (nine belts, hemispheres combined). For consistency, we followed the same taxonomy as Jablonski (1993), detailed in Jablonski & Bottjer (1990).

Cladogenesis and extinction

Data on latitudinal variation in cladogenesis come directly from Jablonski (1993) where methodology is detailed. Briefly, Jablonski (1993) recorded the location of first occurrence of each of the 26 well-preserved marine invertebrate orders that appeared since the beginning of the Mesozoic, with respect to nine, 10° paleolatitudinal belts, hemispheres combined. He then estimated latitudinal variation in cladogenesis by examining the latitudinal distribution of first occurrence for these 26 orders after controlling for a temperate-zone bias in sampling effort. Controlling for variation in sampling effort addressed both issues of collection effort and variation in the size of 10° latitudinal belts that could bias the location of first occurrence of orders (Jablonski, 1993).

Extinction is defined here as the global extinction of the order. We estimated the contribution of extinction to the present-day latitudinal gradient in ordinal richness by examining the location of first occurrence of orders that subsequently went extinct, and plotting the latitudinal distribution of those orders after correcting for sampling bias using the same methodology as Jablonski (1993). If orders originated in the tropics, shifted their distributions out of the tropics and into high latitudes, and then went extinct, extinction could contribute to the latitudinal gradient in ordinal richness indirectly. We tested predictions of this hypothesis as well, using distributional summaries of extinct orders from Moore (1966) and Moore & Teichert (1978), specifically looking for evidence of distributional shifts of each order from tropical to high latitudes prior to extinction.

Net immigration/emigration

Immigration is defined here as the movement (i.e. range shift or expansion) of an extant order into a latitudinal

belt; emigration is defined as the movement (i.e. range shift or retraction) of an extant order out of a latitudinal belt, where that order did not become globally extinct. Current levels of ordinal richness in each latitudinal belt (R_b) can be partitioned into the contributions of cladogenesis (c_b), extinction (x_b), immigration (i_b) and emigration (e_b) as:

$$R_b = (c_b - x_b) + (i_b - e_b). \quad (1)$$

The net contribution of immigration and emigration to the present-day gradient can be defined as net movement (M_b) into or out of latitudinal belts. We calculate this value as:

$$M_b = (i_b - e_b) = R_b - (c_b - x_b), \quad (2)$$

where positive, M_b indicates a net gain of orders (i.e. net immigration), where negative, a net loss of orders (i.e. net emigration).

We hypothesized that the present-day pattern of ordinal richness should decline towards the poles if these taxa show a typical latitudinal gradient in richness. If cladogenesis contributed to the present-day gradient, then it should be negatively related to latitude. If extinction contributed to the gradient, then it should be positively related to latitude. If net movement of orders contributed to the gradient, then it should be negatively related to latitude, with greater immigration into lower latitude regions. We tested all these predictions using nonlinear regression, with latitude as the independent variable. We used Hill three-parameter equations to model the data, as these equations best described the distribution of most of the data (e.g. present-day distribution, cladogenesis). We use one-tailed tests of significance as all predictions were *a priori* and directional. All significant results remain significant after adjusting α values for multiple tests (four tests; Rice, 1989).

Results

At the beginning of the Mesozoic (T_0), none of the 26 benthic marine invertebrate orders existed (Fig. 1, T_0 ; Jablonski, 1993). At present (T_1), these orders exhibit a latitudinal decline in diversity towards the poles (Fig. 1, T_1 ; $y = 21.2x^{-4}/(62.1^{-4} + x^{-4})$, $r^2 = 0.975$, $F = 119.4$, one-tailed $P < 0.0001$). As shown previously (Jablonski, 1993), cladogenesis was higher at lower latitudes (Fig. 1, cladogenesis; $y = 7x^{-5.8}/(36.4^{-5.8} + x^{-5.8})$, $r^2 = 0.984$, $F = 242.8$, one-tailed $P < 0.0001$). Only 5 of the 26 orders became extinct since cladogenesis. Their locations of first occurrence (uncorrected for variation in sampling effort) were: Encrinida (21–30°), Hemicidaroida (21–30°), Pygasteroida (11–20°), Oligopygoida (concurrently found at 0–10° and 21–30°) and Disasteroida (31–40°) (Jablonski, 1993). The distribution of locations of first occurrence remained similar after we corrected for variation in sampling effort, with extinction reducing the effects of cladogenesis, and reducing diversity, at lower latitudes




Fig. 1 Net change in richness of 26 benthic marine invertebrate orders that have well-preserved fossil records. Twenty-six orders originated since the beginning of the Mesozoic (T_0) (cladogenesis), with higher estimated rates of cladogenesis in the tropics (Jablonski, 1993). Five of these orders have since gone extinct (extinction). After cladogenesis and extinction, higher range expansion at lower latitudes has dominated net change in ordinal diversity (movement) to create the present-day latitudinal gradient in ordinal richness (T_1).

(Fig. 1, extinction; $y = 1.9x^{-12.4}/(25.0^{-12.4} + x^{-12.4})$, $r^2 = 0.617$, $F = 7.4$, one-tailed $P = 0.98$). Extinction rates were significantly higher at lower latitudes (two-tailed $P = 0.02$), and thus extinction could not have directly caused the latitudinal gradient in ordinal diversity. Furthermore, none of the extinct orders showed distributional shifts poleward prior to extinction (Moore, 1966; Moore & Teichert, 1978), ruling out an indirect contribution of extinction to the latitudinal gradient.

Net movement of orders since cladogenesis and extinction was positive at all latitudes (Fig. 1, movement), but more orders immigrated into lower latitudinal bands more often compared with higher latitudes (Fig. 1, movement; $y = 15.8x^{-5.9}/(72.0^{-5.9} + x^{-5.9})$, $r^2 = 0.938$, $F = 61.3$, one-tailed $P < 0.0001$), resulting in all 21 extant orders occurring at lower latitudes (Fig. 1, T_1). This saturation of extant orders at lower latitudes illustrates that range expansion, rather than range shifting, resulted in the current distributional patterns

over latitude. The slight decline in net movement near the equator (Fig. 1, movement) is an artefact of higher rates of tropical cladogenesis. If we control for rates of cladogenesis and extinction, we find consistently high net movement into tropical belts (0–30°), with a subsequent decline towards the poles (data not shown). Range expansion of orders following cladogenesis had a greater effect on the creation of the present-day latitudinal gradient in ordinal diversity than did cladogenesis itself (cf. Fig. 1).

Results illustrate that the present-day latitudinal gradient in ordinal richness was caused by declines in cladogenesis and range expansion with increasing latitude. Extinction and net emigration played no significant role in the creation of the latitudinal gradient in ordinal richness.

Discussion

Fossil and present-day patterns of the 26 well-preserved, benthic marine invertebrate orders examined here suggest that greater tropical richness is the result of higher rates of cladogenesis and subsequent range expansion at low latitudes (Fig. 1). The roles of both cladogenesis and range expansion are consistent with the idea of two distinct patterns comprising the latitudinal gradient in diversity: higher rates of cladogenesis and a greater number of sympatric species in the tropics (cf. Terborgh, 1992, p. 7; Fjelds , 1994; Martin & McKay, 2004). The latter pattern is strongly correlated with estimates of available energy (Wright, 1983; Currie, 1991; Francis & Currie, 2003; Hawkins *et al.*, 2003; Currie *et al.*, 2004), which may allow more taxa to expand their ranges into sympatry in tropical areas. Causes of latitudinal variation in rates of cladogenesis remain obscure, but possibilities include: (1) reduced energy at higher latitudes that causes higher rates of population extinction and subsequent recolonization, degrading population structure and impeding cladogenesis (Simpson, 1964; Maruyama & Kimura, 1980; but see Whitlock, 1992 and references within), (2) greater seasonality in temperature at higher latitudes that favours broader physiological tolerance and high dispersal, reducing population divergence and cladogenesis (Janzen, 1967; Ghalambor *et al.*, 2006), (3) increased shifting of distributions at higher latitudes during Milankovitch (glacial) cycles that causes population mixing, bottlenecks and sampling bias during recolonization, reducing rates of population divergence and cladogenesis (Dennis *et al.*, 1995; McGlone, 1996; Dynesius & Jansson, 2000; Hewitt, 2000; Jansson & Dynesius, 2002; Jansson, 2003; Coope, 2004), and (4) greater energy at lower latitudes that causes faster rates of evolution and cladogenesis by reducing generation time, increasing metabolic rates and/or increasing exposure to ultraviolet radiation (Rohde, 1992, 1999; Barracough & Savolainen, 2001; Bromham & Cardillo, 2003; Davies *et al.*, 2004; Allen *et al.*, 2006; Wright *et al.*, 2006;

see also Fischer, 1960; Wright *et al.*, 2003; Brown & Pauly, 2005; Pawar, 2005).

Our results provide support for recent work by Jablonski *et al.* (2006) who found evidence for an 'out of the tropics' model for the creation of the latitudinal gradient in genera of marine bivalves, where lineages preferentially originated in the tropics and persisted there as they expanded poleward. The main difference between the bivalve genera and marine invertebrate orders examined here is that estimated extinction rates were higher in extratropical vs. tropical bivalve genera, suggesting that low tropical extinction rates contributed to the present-day latitudinal gradient in bivalve diversity (Jablonski *et al.*, 2006). In the present study, the latitudinal gradient in ordinal diversity originated in spite of higher tropical extinction rates, illustrating that low tropical extinction rates were not required for the evolution of the latitudinal diversity gradient.

Although fossil evidence suggests that most of the 26 marine invertebrate orders examined here arose in tropical waters (Jablonski, 1993), the coarse nature of controlling for sampling bias across latitudes may overstate the slope of decline in cladogenesis towards the poles (Fig. 1, cladogenesis). Intensified fossil sampling in tropical regions would improve the accuracy of our estimates of the location of origin of these orders. In addition, we should note that cladogenesis and levels of endemism, particularly at lower taxonomic levels, vary between polar regions, being notably higher in the Southern Ocean (see Crame, 1997, 2000; Clarke & Johnston, 2003). Such variation between high-latitude regions may provide important insight into the abiotic causes of latitudinal variation in rates of cladogenesis. Finally, although we correct for latitudinal variation in sampling intensity, such a correction does not control for potential bias in taphonomy across latitudes that could result from reduced preservation in warm tropical waters (Kidwell & Baumiller, 1989), or enhanced preservation of larger tropical taxa (Vermeij, 1978; Martin, 1999, p. 132).

This analysis provides one of the first examinations of all four possible mechanisms underlying the latitudinal gradient in taxonomic diversity. Consistency in the latitudinal gradients in diversity across taxa (invertebrates, vertebrates and plants) and environments (marine, freshwater and terrestrial) (Willig *et al.*, 2003; Hillebrand, 2004) suggests that these patterns result from the same causes. If latitudinal variation in the richness of these 26 marine invertebrate orders is representative of other taxonomic levels and groups, then cladogenesis and range expansion may be the critical mechanisms that link abiotic causes with the resulting pattern of latitudinal variation in diversity.

Acknowledgments

We thank S. Cairns, B. David, R. Greene, C. Messing and F. Stiller for assistance with documenting distributions.

We thank T. Bradshaw, A. Clarke, C. del Agua, M. Donaldson, R. Huey, D. Jablonski and B. Swalla for helpful comments on earlier versions of the manuscript. P.R.M. acknowledges support from the Natural Sciences and Engineering Research Council of Canada, Sigma Xi, and an Ed Harrison Memorial Scholarship from the Western Foundation of Vertebrate Zoology.

References

- Allen, A.P. & Gillooly, J.F. 2006. Assessing latitudinal gradients in speciation rates and biodiversity at the global scale. *Ecol. Lett.* **9**: 947–954.
- Allen, A.P., Gillooly, J.F., Savage, V.M. & Brown, J.H. 2006. Kinetic effects of temperature on rates of genetic divergence and speciation. *Proc. Natl. Acad. Sci. U.S.A.* **103**: 9130–9135.
- Barraclough, T.G. & Savolainen, V. 2001. Evolutionary rates and species diversity in flowering plants. *Evolution* **55**: 677–683.
- Bennett, K.D. 1997. *Evolution and Ecology: The Pace of Life*. Cambridge University Press, Cambridge.
- Bromham, L. & Cardillo, M. 2003. Testing the link between the latitudinal gradient in species richness and rates of molecular evolution. *J. Evol. Biol.* **16**: 200–207.
- Brown, J.M. & Pauly, G.B. 2005. Increased rates of molecular evolution in an equatorial plant clade: an effect of environment or phylogenetic nonindependence? *Evolution* **59**: 238–242.
- Cardillo, M. 1999. Latitude and rates of diversification in birds and butterflies. *Proc. R. Soc. Lond. B, Biol. Sci.* **266**: 1221–1225.
- Cardillo, M., Orme, C.D.L. & Owens, I.P.F. 2005. Testing for latitudinal bias in diversification rates: an example using New World birds. *Ecology* **86**: 2278–2287.
- Chek, A.A., Austin, J.D. & Lougheed, S.C. 2003. Why is there a tropical-temperate disparity in the genetic diversity and taxonomy of species? *Evol. Ecol. Res.* **5**: 69–77.
- Chown, S.L. & Gaston, K.J. 2000. Area, cradles and museums: the latitudinal gradient in species richness. *Trends Ecol. Evol.* **15**: 311–315.
- Clarke, A. & Johnston, N.M. 2003. Antarctic marine benthic diversity. *Ocean. Mar. Biol. Ann. Rev.* **41**: 47–114.
- Coope, G.R. 1995. Insect fauna in ice age environments: why so little extinction? In: *Extinction Rates* (J. H. Lawton & R. M. May, eds), pp. 55–74. Oxford University Press, Oxford.
- Coope, G.R. 2004. Several million years of stability among insect species because of, or in spite of, Ice Age climatic instability? *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **359**: 209–214.
- Crame, J.A. 1997. An evolutionary framework for the polar regions. *J. Biogeogr.* **24**: 1–9.
- Crame, J.A. 2000. Evolution of taxonomic diversity gradients in the marine realm: evidence from the composition of recent bivalve faunas. *Paleobiology* **26**: 188–214.
- Currie, D.J. 1991. Energy and large-scale patterns of animal- and plant-species richness. *Am. Nat.* **137**: 27–49.
- Currie, D.J., Mittelbach, G.G., Cornell, H.V., Field, R., Guégan, J.-F., Hawkins, B.A., Kaufman, D.M., Kerr, J.T., Oberdorff, T., O'Brien, E. & Turner, J.R.G. 2004. Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic richness. *Ecol. Lett.* **7**: 1121–1134.
- Davies, T.J., Savolainen, V., Chase, M.W., Moat, J. & Barraclough, T.G. 2004. Environmental energy and evolutionary

- rates in flowering plants. *Proc. R. Soc. Lond., B, Biol. Sci.* **271**: 2195–2200.
- Dennis, R.L.H., Shreeve, T.G. & Williams, W.R. 1995. Taxonomic differentiation in species richness gradients among European butterflies (Papilionoidea, Hesperioidea): contribution of macroevolutionary dynamics. *Ecography* **18**: 27–40.
- Dynesius, M. & Jansson, R. 2000. Evolutionary consequences of changes in species' geographical distributions driven by Milankovitch climate oscillations. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 9115–9120.
- Fischer, A.G. 1960. Latitudinal variation in organic diversity. *Evolution* **14**: 64–81.
- Fjeldså, J. 1994. Geographical patterns for relict and young species of birds in Africa and South America and implications for conservation priorities. *Biodiv. Conserv.* **3**: 207–226.
- Flessa, K.W. & Jablonski, D. 1996. The geography of evolutionary turnover: a global analysis of extant bivalves. In: *Evolutionary Paleobiology* (D. Jablonski, D. H. Erwin & J. H. Lipps, eds), pp. 376–397. University of Chicago Press, Chicago, IL.
- Francis, A.P. & Currie, D.J. 2003. A globally consistent richness-climate relationship for angiosperms. *Am. Nat.* **161**: 523–536.
- Gaston, K.J. 2000. Global patterns in biodiversity. *Nature* **405**: 220–227.
- Gaston, K.J. & Blackburn, T.M. 1996. The tropics as a museum of biological diversity: an analysis of the New World avifauna. *Proc. R. Soc. Lond., B, Biol. Sci.* **263**: 63–68.
- Ghalambor, C., Huey, R.B., Martin, P.R., Tewksbury, J.J. & Wang, G. 2006. Are mountain passes higher in the tropics? Janzen's hypothesis revisited. *Integr. Comp. Biol.* **46**: 5–17.
- Goldberg, E.E., Roy, K., Lande, R. & Jablonski, D. 2005. Diversity, endemism, and age distributions in macroevolutionary sources and sinks. *Am. Nat.* **165**: 623–633.
- Harrington, G.J. 2004. Structure of the North American vegetation gradient during the Late Paleocene/Early Eocene warm climate. *Evol. Ecol. Res.* **6**: 33–48.
- Haskell, J. 2001. The latitudinal gradient of diversity through the Holocene as recorded by fossil pollen in Europe. *Evol. Ecol. Res.* **3**: 345–360.
- Hawkins, B.A., Field, R., Cornell, H.V., Currie, D.J., Guégan, J.-F., Kaufman, D.M., Kerr, J.T., Mittelbach, G.G., Oberdorff, T., O'Brien, E.M., Porter, E.E. & Turner, J.R.G. 2003. Energy, water, and broad-scale geographic patterns of species richness. *Ecology* **84**: 3105–3117.
- Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.
- Hillebrand, H. 2004. On the generality of the latitudinal diversity gradient. *Am. Nat.* **163**: 192–211.
- Jablonski, D. 1985. Causes and consequences of mass extinctions: a comparative approach. In: *Dynamics of Extinction* (D. K. Elliot, ed.), pp. 183–229. Wiley & Sons, New York.
- Jablonski, D. 1993. The tropics as a source of evolutionary novelty through geological time. *Nature* **364**: 142–144.
- Jablonski, D. & Bottjer, D.J. 1990. The origin and diversification of major groups: environmental patterns and macroevolutionary lags. In: *Major Evolutionary Radiations* (P. D. Taylor & G. P. Larwood, eds), pp. 17–57. Clarendon Press, Oxford.
- Jablonski, D., Flessa, K.W. & Valentine, J.W. 1985. Biogeography and paleobiology. *Paleobiology* **11**: 75–90.
- Jablonski, D., Roy, K. & Valentine, J.W. 2006. Out of the tropics: evolutionary dynamics of the latitudinal diversity gradient. *Science* **314**: 102–106.
- Jansson, R. 2003. Global patterns in endemism explained by past climatic change. *Proc. R. Soc. Lond., B, Biol. Sci.* **270**: 583–590.
- Jansson, R. & Dynesius, M. 2002. The fate of clades in a world of recurrent climatic change: Milankovitch oscillations and evolution. *Annu. Rev. Ecol. Syst.* **33**: 741–777.
- Janzen, D.H. 1967. Why mountain passes are higher in the tropics. *Am. Nat.* **101**: 233–249.
- Jetz, W., Rahbek, C. & Colwell, R.K. 2004. The coincidence of rarity and richness and the potential signature of history in centres of endemism. *Ecol. Lett.* **7**: 1180–1191.
- Kidwell, S.M. & Baumiller, T. 1989. Experimental disintegration of regular echinoids: roles of temperature, oxygen, and decay thresholds. *Paleobiology* **16**: 247–271.
- Martin, R.E. 1999. *Taphonomy: A Process Approach*. Cambridge University Press, Cambridge.
- Martin, P.R. & McKay, J.K. 2004. Latitudinal variation in genetic divergence of populations and the potential for future speciation. *Evolution* **58**: 938–945.
- Maruyama, T. & Kimura, M. 1980. Genetic variability and effective population size when local extinction and recolonization of subpopulations are frequent. *Proc. Natl. Acad. Sci. U.S.A.* **77**: 6710–6714.
- McGlone, M.S. 1996. When history matters: scale, time, climate and tree diversity. *Global Ecol. Biogeogr. Lett.* **5**: 309–314.
- Moore, R.C., ed. 1966. *Treatise on Invertebrate Paleontology. Part U. Echinodermata 3*. Geological Society of America, Boulder, CO.
- Moore, R.C. & Teichert, C., eds. 1978. *Treatise on Invertebrate Paleontology. Part T. Echinodermata 2*. Geological Society of America, Boulder, CO.
- Pawar, S.S. 2005. Geographical variation in the rate of evolution: effect of available energy or fluctuating environment? *Evolution* **59**: 234–237.
- Pimm, S.L. & Brown, J.H. 2004. Domains of diversity. *Science* **304**: 831–833.
- Rice, W.R. 1989. Analyzing tables of statistical tests. *Evolution* **43**: 223–225.
- Ricklefs, R.E. 2004. A comprehensive framework for global patterns in biodiversity. *Ecol. Lett.* **7**: 1–15.
- Rohde, K. 1992. Latitudinal gradients in species diversity: the search for the primary cause. *Oikos* **65**: 514–527.
- Rohde, K. 1999. Latitudinal gradients in species diversity and Rapoport's rule revisited: a review of recent work and what can parasites teach us about the causes of the gradients? *Ecography* **22**: 593–613.
- Sepkoski, J.J., Jr. 1999. Rates of speciation in the fossil record. In: *Evolution of Biological Diversity* (A. E. Magurran & R. M. May, eds), pp. 260–282. Oxford University Press, Oxford.
- Silvertown, J. 1985. History of a latitudinal diversity gradient: woody plants in Europe 13,000–1000 years B.P. *J. Biogeogr.* **12**: 519–525.
- Simpson, G.G. 1964. Species density of North American recent mammals. *Syst. Zool.* **13**: 57–73.
- Stanley, S.M. 1984. Marine mass extinctions: a dominant role for temperature. In: *Extinctions* (M. H. Nitecki, ed.), pp. 69–117. University of Chicago Press, Chicago, IL.
- Terborgh, J. 1992. *Diversity and the Tropical Rain Forest*. Scientific American Library, New York.
- Vermeij, G.J. 1978. *Biogeography and Adaptation*. Harvard University Press, Cambridge.
- Whitlock, M.C. 1992. Temporal fluctuations in demographic parameters and the genetic variance among populations. *Evolution* **46**: 608–615.

- Willig, M.R., Kaufman, D.M. & Stevens, R.D. 2003. Latitudinal gradients of biodiversity: pattern, process, scale, and synthesis. *Annu. Rev. Ecol. Evol. Syst.* **34**: 273–309.
- Wright, D.H. 1983. Species-energy theory: an extension of species-area theory. *Oikos* **41**: 496–506.
- Wright, S.D., Gray, R.D. & Gardner, R.C. 2003. Energy and the rate of evolution: inferences from plant rDNA substitution rates in the western Pacific. *Evolution* **57**: 2893–2898.

- Wright, S., Keeling, J. & Gillman, L. 2006. The road from Santa Rosalia: a faster tempo of evolution in tropical climates. *Proc. Natl. Acad. Sci. U.S.A.* **103**: 7718–7722.

Received 6 October 2006; accepted 22 December 2006

Appendix 1 Present distribution of marine invertebrate orders examined in this study. Superscripts denote references for presence/absence data following table.

Phylum	Class	Order	0–10°	10–20°	20–30°	30–40°	40–50°	50–60°	60–70°	70–80°	80–90°
Porifera	Hexactinellida	Lychniscosida	Present ¹⁹	Present ¹⁹	Present ¹⁹	Absent ¹⁹	Absent ¹⁹	Absent ¹⁹	Absent ¹⁹	Absent ¹⁹	Absent ¹⁹
Coelenterata	Hydrozoa	Milleporina	Present ²⁵	Present ²⁵	Present ²⁵	Present ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵
Coelenterata	Hydrozoa	Stylasterina	Present ²	Present ²	Present ²	Present ²	Present ²	Present ²	Present ²	Absent ²	Absent ²
Coelenterata	Anthozoa	Helioporacea	Present ²⁵	Present ²⁵	Present ²⁵	Present ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵
Coelenterata	Anthozoa	Scleractina	Present ²⁵	Present ²⁵	Present ²⁵	Present ²⁵	Present ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵	Absent ²⁵
Mollusca	Gastropoda	Neogastropoda	Present ²⁶	Present ²⁶	Present ²⁶	Present ²⁶	Present ²⁶	Present ²⁶	Present ²⁴	Present ¹	Present ¹
Bryozoa	Gymnolaemata	Cheilostomata	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹⁷	Present ¹¹
Echinodermata	Crinoidea	Isocrinida	Present ¹³	Present ¹³	Present ⁹	Present ⁹	Present ⁹	Absent ²⁰	Absent ²⁰	Absent ²⁰	Absent ²⁰
Echinodermata	Crinoidea	Millericrinida	Present ¹²	Present ²⁰	Present ¹²	Present ⁹	Present ⁴	Present ²⁰	Present ²⁰	Present ²⁰	Absent ²⁰
Echinodermata	Crinoidea	Cyrtocrinida	Present ²⁰	Present ¹³	Present ¹³	Present ⁹	Absent ²⁰	Absent ²⁰	Absent ²⁰	Absent ²⁰	Absent ²⁰
Echinodermata	Crinoidea	Bourgueticrinida	Present ¹²	Present ⁹	Present ¹²	Present ¹²	Present ⁸	Present ⁸	Present ⁸	Present ⁹	Absent ²⁰
Echinodermata	Crinoidea	Encrinida	Extinct ²²	–	–	–	–	–	–	–	–
Echinodermata	Echinoidea	Pedinoida	Present ¹²	Present ¹⁸	Present ¹⁸	Present ¹⁸	Present ¹⁸	Absent ¹⁸	Absent ¹⁸	Absent ¹⁸	Absent ¹⁸
Echinodermata	Echinoidea	Hemicidaroida	Extinct ²¹	–	–	–	–	–	–	–	–
Echinodermata	Echinoidea	Salenoida	Present ⁶	Present ⁶	Present ⁶	Present ⁶	Present ⁶	Absent ⁶	Present ⁶	Absent ⁶	Absent ⁶
Echinodermata	Echinoidea	Phymosomatoida	Present ¹⁸	Present ¹⁸	Present ¹⁸	Present ¹⁸	Present ^{7,18}	Present ^{7,18}	Present ¹⁸	Absent ¹⁸	Absent ¹⁸
Echinodermata	Echinoidea	Temnopleuroida	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹⁸	Present ¹⁸	Present ¹⁸	Absent ¹⁸	Absent ¹⁸
Echinodermata	Echinoidea	Echinoida	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹⁸	Present ¹⁸	Absent ¹⁸
Echinodermata	Echinoidea	Pygasteroida	Extinct ^{21,23}	–	–	–	–	–	–	–	–
Echinodermata	Echinoidea	Holectypoida	Present ¹⁸	Present ¹⁸	Present ¹⁸	Present ^{3,18}	Present ¹⁸	Absent ¹⁸	Absent ¹⁸	Absent ¹⁸	Absent ¹⁸
Echinodermata	Echinoidea	Cassiduloida	Present ¹⁶	Present ¹⁶	Present ¹⁶	Present ^{16,10}	Present ^{16,10}	Present ¹⁰	Absent ¹⁶	Absent ¹⁶	Absent ¹⁶
Echinodermata	Echinoidea	Oligopygoida	Extinct ²³	–	–	–	–	–	–	–	–
Echinodermata	Echinoidea	Clypeasteroida	Present ¹⁵	Present ¹⁵	Present ¹⁵	Present ¹⁵	Present ¹⁵	Present ¹⁵	Present ¹⁵	Present ¹⁵	Absent ¹⁵
Echinodermata	Echinoidea	Disasteroida	Extinct ²³	–	–	–	–	–	–	–	–
Echinodermata	Echinoidea	Holasteroida	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ⁵	Present ¹⁴	Present ¹⁴
Echinodermata	Echinoidea	Spatangoida	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹²	Present ¹⁸	Present ¹⁸	Absent ¹⁸

- Aitken, A.E. & Gilbert, R. 1996. Marine mollusca from expedition Fjord, western Axel Heiberg Island, Northwest Territories, Canada. *Arctic* **49**: 29–43.
- Cairns, S.D. 1992. Worldwide distribution of the Stylasteridae (Cnidaria: Hydrozoa). *Sci. Mar.* **60**: 125–130.
- Clark, H.L. 1946. *The Echinoderm Fauna of Australia*. Carnegie Institute, Washington, D.C.
- Commonwealth of Australia. 2002. *Tasmanian Sea Mount Marine Reserve Management Plan*. Environment Australia, Canberra.
- David, B. 1988. Origins of the deep-sea holasteroid fauna. In: *Echinoderm Phylogeny and Evolutionary Biology* (C.R.C. Paul & A.B. Smith, eds), pp. 331–346. Clarendon Press, Oxford.
- Durkin, M.K. 1980. The Saleniinae in time and space. In: *Echinoderms: Present and Past* (M. Jangoux, ed.), pp. 3–14. A.A. Balkema, Rotterdam, The Netherlands.
- Emlet, R.B. 1995. Developmental mode and species geographic range in regular sea-urchins (Echinodermata, Echinoidea). *Evolution* **49**: 476–489.
- Hansson, H.G., compiler. 1998. North east Atlantic taxa: south Scandinavian marine Echinodermata check-list. Internet pdf, <http://www.tmbi.gu.se>.
- Hansson, H.G., compiler. 1999. European Echinodermata check-list. Internet pdf, <http://www.tmbi.gu.se>.
- Higgins, R.C. 1974. Observations on the biology of *Apatopygus recens* (Echinoidea: Cassiduloida) around New Zealand. *J. Zool (Lond.)* **173**: 505–516.
- Kluge, G.A. 1962. *Bryozoa of the Northern Seas of the USSR*. Akademiya Nauk, Moscow.
- Maluf, L.Y. 1988. *Composition and Distribution of the Central Eastern Pacific Echinoderms*. Natural History Museum of Los Angeles County, Los Angeles.

13. Meyer, D.L., Messing, C.G. & Macurda, D.B., Jr. 1978. Biological results of University of Miami deep-sea expeditions. 129. Zoogeography of tropical western Atlantic Crinoidea (Echinodermata). *Bull. Mar. Sci.* **28**: 412–441.
14. Mironov, A.N. 1995. Holasteroid sea urchins. 1. Morphological diversity of *Pourtalesia jeffreysi*. *Zool. Zhurnal.* **74**: 68–77.
15. Mooi, R. 1989. Living and fossil genera of the Clypeasteroidea (Echinoidea: Echinodermata): an illustrated key and annotated checklist. *Smithsonian Contr. Zool.* **488**: 1–51.
16. Mooi, R. 1990. Living Cassiduloids (Echinodermata, Echinoidea) – a key and annotated list. *Proc. Biol. Soc. Wash.* **103**: 63–85.
17. Morris, P.A. 1980. *The Bryozoan Family Hippothoidae (Cheilostomata – Ascophora) with Emphasis on the Genus Hippothoa*. University of Southern California, Los Angeles.
18. Mortensen, T. 1928–1951. *A Monograph of the Echinodea. I-V*. C.A. Reitzel, Copenhagen.
19. Reiswig, H.M. 2002. Order Lychniscosida Schrammen, 1903. In: *Systema Porifera: A Guide to the Classification of Sponges* (J.N.A. Hooper & R.W.M. Van Soest, eds), pp. 1377–1385. Kluwer Academic/Plenum, New York.
20. Roux, M., Messing, C.G. & Améziane, N. 2002. Artificial keys to the genera of living stalked crinoids (Echinodermata). *Bull. Mar. Sci.* **70**: 799–830.
21. Sepkoski, J.J., Jr. 1982. A compendium of fossil marine families. *Milwaukee Public Mus., Contrib. Biol. Geol.* **51**: 1–125.
22. Simms, M.J. 1999. Systematics, phylogeny and evolutionary history. In: *Fossil Crinoids* (H. Hess, W.I. Ausich, C.E. Brett & M.J. Simms, eds), pp. 31–40. Cambridge University Press, Cambridge.
23. Smith, A. 1984. *Echinoid Palaeobiology*. George Allen & Unwin, London.
24. Stanwell-Smith, D., Hood, A. & Peck, L. 1997. *A Field Guide to the Pelagic Invertebrate Larvae of the Maritime Antarctic*. British Antarctic Survey, Cambridge.
25. Veron, J. 2000. *Corals of the World*, Vols 1–3. Australian Institute of Marine Science, Townsville.
26. Wagner, R.J.L. & Abbott, R.T. 1978. *Standard Catalog of Shells*, 3rd edn. with Supplements. American Malacologists, Greenville, DE.