

## Damage formation and repair data for radiations commonly used in microbeam studies of low-dose radiation biology

During the last several years charged-particle microbeams proved themselves as powerful tools in radiobiological research (for review see [1]). Currently two microbeam facilities are routinely used for radiobiological experiments. These are the [Radiological Research Accelerator Facility \(RARAF\)](#) at Columbia University [2] and the [Gray Cancer Institute](#) (formerly Gray Laboratory) microbeam [3, 4]. The [Special Microbeam Utilization Research Facility \(SMURF\)](#) at Texas A&M University is based on microbeam equipment relocated from Pacific Northwest National Laboratory [5] and is also available for radiobiological studies.

We used a [Monte Carlo excision repair \(MCER\) model](#) [6, 7] to generate damage formation and repair data for particles that have been used in various microbeam studies. The DNA damage portion of the MCER model is based on the [Monte Carlo damage simulation \(MCDS\) algorithm](#) proposed by us [8, 9]. The MCDS algorithm is capable of generating data for protons and  $\alpha$  particles of different energies and for 4.5 keV electrons considered representative of low-LET radiation. The MCDS model parameters for heavy-charged particles (protons and  $\alpha$  particles) are identical with the exception of one parameter,  $n_{seg}$ , which is dependent upon the particle LET (refer to Fig. 5 in [8]). This property allows one to determine the algorithm parameters for other heavy-charged particles of the same LET. Using this approximation, damage formation and repair data for 4 MeV  $\alpha$  particles, which have LET of approximately 100 keV/ $\mu$ m, can be used in place of data for  $^3\text{He}^{2+}$  ions that have been extensively used in studies performed with the Gray Cancer Institute microbeam.

---

### Columbia University

The Columbia University microbeam is designed to deliver helium or hydrogen ions with different energies that cover the range of LET from 30 to 220 keV/ $\mu$ m [2]. To date, all the studies that utilized this microbeam have been performed with 5.3 MeV  $\alpha$  particles that have LET of approximately 90 keV/ $\mu$ m (for examples of studies with this microbeam system, see [10–21]).

- 5.3 MeV  $\alpha$  particles ([input file](#), [output file](#))

---

### Gray Cancer Institute

The Gray Cancer Institute microbeam was initially configured to irradiate cells with protons of energies <3.5 MeV [3]. Later a change has been made to also deliver  $^3\text{He}^{2+}$  ions. The majority of studies performed using the Gray Cancer Institute microbeam used  $^3\text{He}^{2+}$  ions with LET of  $\sim$ 100 keV/ $\mu$ m [22–26]. 1 and 3.2 MeV protons have also been utilized in some experiments [27, 28].

- 4 MeV  $\alpha$  particles; equivalent to 100 keV/ $\mu$ m  $^3\text{He}^{2+}$  ions ([input file](#), [output file](#))
- 1 MeV protons ([input file](#), [output file](#))
- 3.2 MeV protons ([input file](#), [output file](#))

---

### PNNL/Texas A&M University

The Texas A&M University microbeam is capable of producing  $\alpha$  particles and protons with energies up to 6 MeV and up to 4 MeV respectively. Also available is a 100 keV electron microbeam. The predecessor of TAMU's microbeam, the microbeam at Pacific Northwest National Laboratory, has been applied in one study that utilized 3.2 MeV  $\alpha$  particles [29].

- 3.2 MeV  $\alpha$  particles ([input file](#), [output file](#))

---

## References

- [1] Folkard M, Prise KM, Vojnovic B, Gilchrist S, Schettino G, Belyakov OV, Ozols A, Michael BD. The impact of microbeams in radiation biology. *Nucl. Instr. Meth. Phys. Res. B* 181 (2001) 426–430.
- [2] Randers-Pehrson G, Geard CR, Johnson G, Elliston CD, Brenner DJ. The Columbia university single-ion microbeam. *Radiat. Res.* 156 (2001) 210–214. [[PubMed](#)]
- [3] Folkard M, Vojnovic B, Prise KM, Bowey AG, Locke RJ, Schettino G, Michael BD. A charged-particle microbeam: I. Development of an experimental system for targeting cells individually with counted particles. *Int. J. Radiat. Biol.* 72 (1997) 375–385. [[Abstract](#)]
- [4] Folkard M, Vojnovic B, Hollis KJ, Bowey AG, Watts SJ, Schettino G, Prise KM, Michael BD. A charged-particle microbeam: II. A single-particle micro-collimation and detection system. *Int. J. Radiat. Biol.* 72 (1997) 387–395. [[PubMed](#)]
- [5] Braby LA, Brooks AL, Metting NF. Cellular effects of individual high-linear energy transfer particles and implications for tissue response at low doses. *Radiat. Res.* 148 (1997) S108–S114. [[PubMed](#)]
- [6] Semenenko VA, Stewart RD, Ackerman EJ. Monte Carlo simulation of base and nucleotide excision repair of clustered DNA damage sites. I. Model properties and predicted trends. *Radiat. Res.* 164 (2005) 180–193. [[PubMed](#)] [[HomePage](#)]
- [7] Semenenko VA, Stewart RD. Monte Carlo simulation of base and nucleotide excision repair of clustered DNA damage sites. II. Comparisons of model predictions to measured data. *Radiat. Res.* 164 (2005) 194–201. [[PubMed](#)] [[HomePage](#)]
- [8] Semenenko VA, Stewart RD. A fast Monte Carlo algorithm to simulate the spectrum of DNA damages formed by ionizing radiation. *Radiat. Res.* 161 (2004) 451–457. [[PubMed](#)] [[HomePage](#)]
- [9] Semenenko VA, Stewart RD. Fast Monte Carlo simulation of DNA damage formed by electrons and light ions. *Phys. Med. Biol.* 51 (2006) 1693–1706. [[HomePage](#)]
- [10] Hei TK, Wu LJ, Liu SX, Vannais D, Waldren CA, Randers-Pehrson G. Mutagenic effects of a single and an exact number of  $\alpha$  particles in mammalian cells. *Proc. Natl. Acad. Sci. USA* 94 (1997) 3765–3770. [[PubMed](#)]
- [11] Miller RC, Randers-Pehrson G, Geard CR, Hall EJ, Brenner DJ. The oncogenic transforming potential of the passage of single  $\alpha$  particles through mammalian cell nuclei. *Proc. Natl. Acad. Sci. USA* 96 (1999) 19–22. [[PubMed](#)]
- [12] Wu LJ, Randers-Pehrson G, Xu A, Waldren CA, Geard CR, Yu Z, Hei TK. Targeted cytoplasmic irradiation with alpha particles induces mutations in mammalian cells. *Proc. Natl. Acad. Sci. USA* 96 (1999) 4959–4964. [[PubMed](#)]
- [13] Zhou H, Randers-Pehrson G, Waldren CA, Vannais D, Hall EJ, Hei TK. Induction of a bystander mutagenic effect of alpha particles in mammalian cells. *Proc. Natl. Acad. Sci. USA* 97 (2000) 2099–2104. [[PubMed](#)]
- [14] Zhou H, Suzuki M, Randers-Pehrson G, Vannais D, Chen G, Trosko JE, Waldren CA, Hei TK. Radiation risk to low fluences of  $\alpha$  particles may be greater than we thought. *Proc. Natl. Acad. Sci. USA* 98 (2001) 14410–14415. [[PubMed](#)]
- [15] Sawant SG, Randers-Pehrson G, Geard CR, Brenner DJ, Hall EJ. The bystander effect in radiation

oncogenesis: I. Transformation in C3H 10T½ cells *in vitro* can be initiated in the unirradiated neighbors of irradiated cells. Radiat. Res. 155 (2001) 397–401. [\[PubMed\]](#)

[16] Sawant SG, Randers-Pehrson G, Metting NF, Hall EJ. Adaptive response and the bystander effect induced by radiation in C3H 10T½ cells in culture. Radiat. Res. 156 (2001) 177–180. [\[PubMed\]](#)

[17] Sawant SG, Zheng W, Hopkins KM, Randers-Pehrson G, Lieberman HB, Hall EJ. The radiation-induced bystander effect for clonogenic survival. Radiat. Res. 157 (2002) 361–364. [\[PubMed\]](#)

[18] Zhou H, Randers-Pehrson G, Geard CR, Brenner DJ, Hall EJ, Hei TK. Interaction between radiation-induced adaptive response and bystander mutagenesis in mammalian cells. Radiat. Res. 160 (2003) 512–516. [\[PubMed\]](#)

[19] Mitchell SA, Randers-Pehrson G, Brenner DJ, Hall EJ. The bystander response in C3H 10T(1/2) cells: the influence of cell-to-cell contact. Radiat. Res. 161 (2004) 397–401. [\[PubMed\]](#)

[20] Mitchell SA, Marino SA, Brenner DJ, Hall EJ. Bystander effect and adaptive response in C3H 10T(1/2) cells. Int. J. Radiat. Biol. 80 (2004) 465–472. [\[PubMed\]](#)

[21] Ponnaiya B, Jenkins-Baker G, Brenner DJ, Hall EJ, Randers-Pehrson G, Geard CR. Biological responses in known bystander cells relative to known microbeam-irradiated cells. Radiat. Res. 162 (2004) 426–432. [\[PubMed\]](#)

[22] Prise KM, Belyakov OV, Folkard M, Michael BD. Studies of bystander effects in human fibroblasts using a charged particle microbeam. Int. J. Radiat. Biol. 74 (1998) 793–798. [\[PubMed\]](#)

[23] Kadhim MA, Marsden SJ, Goodhead DT, Malcolmson AM, Folkard M, Prise KM, Michael BD. Long-term genomic instability in human lymphocytes induced by single-particle irradiation. Radiat. Res. 155 (2001) 122–126. [\[PubMed\]](#)

[24] Belyakov OV, Malcolmson AM, Folkard M, Prise KM, Michael BD. Direct evidence for a bystander effect of ionizing radiation in primary human fibroblasts. Br. J. Cancer 84 (2001) 674–679. [\[PubMed\]](#)

[25] Belyakov OV, Folkard M, Mothersill C, Prise KM, Michael BD. A proliferation-dependent bystander effect in primary porcine and human urothelial explants in response to targeted irradiation. Br. J. Cancer 88 (2003) 767–774. [\[PubMed\]](#)

[26] Shao C, Stewart V, Folkard M, Michael BD, Prise KM. Nitric oxide-mediated signaling in the bystander response of individually targeted glioma cells. Cancer Res. 63 (2003) 8437–8442. [\[PubMed\]](#)

[27] Schettino G, Folkard M, Prise KM, Vojnovic B, Bowey AG, Michael BD. Low-dose hypersensitivity in Chinese hamster V79 cells targeted with counted protons using a charged-particle microbeam. Radiat. Res. 156 (2001) 526–534. [\[PubMed\]](#)

[28] Tartier L, Spenlehauer C, Newman HC, Folkard M, Prise KM, Michael BD, Menissier-de Murcia J, de Murcia G. Local DNA damage by proton microbeam irradiation induces poly(ADP-ribose) synthesis in mammalian cells. Mutagenesis 18 (2003) 411–416. [\[PubMed\]](#)

[29] Nelson JM, Brooks AL, Metting NF, Khan MA, Buschbom RL, Duncan A, Miick R, Braby LA. Clastogenic effects of defined numbers of 3.2 MeV alpha particles on individual CHO-K1 cells. Radiat. Res. 145 (1996) 568–574. [\[PubMed\]](#)