

Due to Canvas by 11:00pm PDT on the due date, at the latest.

No late papers accepted, so aim to get it in earlier!

To submit, see <https://canvas.uw.edu/courses/1352870/assignments/5301318>

100 points possible.

Open book and notes, and you can use other resources too, except please don't discuss it with other students or anyone else.

If you need clarification on some problem, or you think there's an error or typo, please post it on the Canvas discussion board so everyone has access to the same information.

Problem 1.

In Homework 2 you solved a linear beam equation, valid for small deformations. If the deformations are larger, then the equation must be nonlinear. In some cases an equation of this form can be used:

$$\begin{aligned} au''''(x) - b(u'(x))^2(u''(x))^2 &= f(x) \quad \text{for } 0 \leq x \leq 1 \\ u(0) = \alpha_0, \quad u'(0) = \alpha_1, \quad u(1) = \beta_0, \quad u'(1) = \beta_1. \end{aligned}$$

where $a, b, \alpha_0, \alpha_1, \beta_0, \beta_1$ are all specified constants.

(a) If we discretize with a uniform grid using $h = 1/(m+1)$, suggest a nonlinear system of m equations that could be solved for $[U_1, U_2, \dots, U_m]$ (the interior grid values) to obtain an approximate solution that is second order accurate.

Make sure the boundary conditions are also handled appropriately. Use the "First approach" described in [hw2.solutions.html](#) since this is less messy and should still be second order accurate.

(b) If we wanted to solve this system using Newton's method, we would need the Jacobian matrix for the nonlinear system developed in (a). You don't need to compute the full matrix, but do compute the diagonal elements J_{ii} for $i = 1, 2, \dots, m$.

Problem 2.

Suppose A is a 2×2 singular matrix that is symmetric and has one positive eigenvalue, for example

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$$

is one such matrix. Then A is symmetric positive *semi-definite* ($u^T A u \geq 0$ for all nonzero u) but is not positive definite. If we define the functional

$$\phi(u) = \frac{1}{2} u^T A u - u^T f$$

as in Section 4.3 then levels sets of $\phi(u)$ are no longer ellipses.

(a) What is the geometry of these level sets in general?

(b) Let $z \in \mathbb{R}^2$ be a null vector of A and suppose $f \in \mathbb{R}^2$ is a vector satisfying $z^T f = 0$. Then the system $Au = f$ has infinitely many solutions. Show that the steepest descent method applied to $\phi(u)$ from any initial guess $u^{[0]}$ converges to *some* solution of the linear system in a single iteration. Hint: It might be easier to explain this using the result of (a) than by computing $u^{[1]}$ explicitly. That's fine as long as you give a convincing argument.

(a) Set up a tridiagonal linear system $Au = f$ that could be solved to model this, with the following properties:

- $u = [u_0, u_1, \dots, u_m]$ contains the unknown boundary value u_0 but not the known value $u_{m+1} = \beta$ (assuming as usual that $x_j = jh$ for $j = 0, 1, \dots, m+1$ on a grid with $h = 1/(m+1)$).
- The method is second order accurate.

Follow the strategy of the second approach on page 31 to obtain the first equation in your linear system (i.e. introduce a ghost point u_{-1} and then eliminate it from the two equations that involve this unknown). Write out the matrix and right hand side of your system.

(b) Determine the local truncation error of your method, $\tau = [\tau_0, \tau_1, \dots, \tau_m]$. We expect $\tau_j = C_j h^2 + o(h^2)$ so determine the constants C_j in terms of derivative(s) of the exact solution $u(x)$ (by doing Taylor series expansions, assuming $u(x)$ is sufficiently smooth).

Problem 6. (With corrected boundary conditions)

(a) Implement the method you derived in the previous problem (in Python or Matlab). It is ok to base this on code you have previously written for homework problems, and/or the class Jupyter notebooks.

(b) Test it on the problem

$$\begin{aligned}u''(x) &= 4, & 0 \leq x \leq 1, \\2u(0) + 3u'(0) &= 1, & u(1) = 2,\end{aligned}$$

which has exact solution $u(x) = 2x^2 + x - 1$.

Explain why you expect the exact solution to your linear system to agree with exact solution of the ODE when evaluated at the grid points, and confirm that this is true for your implementation.

(c) Also test it on the problem

$$\begin{aligned}u''(x) &= -18x + 4, & 0 \leq x \leq 1, \\2u(0) + 3u'(0) &= 1, & u(1) = -1,\end{aligned}$$

which has exact solution $u(x) = -3x^3 + 2x^2 + x - 1$. In this case check that your method is second order accurate by producing a log-log plot of the errors (similar to what was done in the notebook BVP1), using $m+1 = 50, 100, 200, 400, 800$.