

28. Diffraction and the Fourier Transform

Diffraction examples


Diffraction by an edge

Arago spot

The far-field

Fraunhofer Diffraction

Some examples




Simeon Poisson
(1781 - 1840)



Francois Arago
(1786 - 1853)



A plane wave incident on a sharp edge




Fresnel diffraction integral, in one dimension:

$$E(x_0) = \int_{x_1=0}^{\infty} \frac{1}{j\lambda z} \exp \left\{ jk \left[z + \frac{(x_0 - x_1)^2}{2z} \right] \right\} dx_1$$

geometrical shadow


exact diffraction result



The irradiance exactly at the edge is 25% of the value far from the edge.


Diffraction by an Edge

Light passing
by an edge



(a)


Electrons passing
by an edge




(b)

An interesting manifestation of diffraction effects: The Spot of Arago

If a beam encounters a circular “stop”, it develops a hole, which fills in as it propagates and diffracts:




Interestingly, the hole fills in from the center first!



The Spot of Arago

Why does it happen?



According to Huygen's principle, every point on the perimeter of the disc radiates a spherical wave.

Obviously, the distance from each spherical wave's point of origin to a point on the central axis of the disc is the same.

→ Constructive interference on the center line!

In 1818, Poisson used Fresnel's theory to predict this phenomenon. He regarded this as proof that Fresnel's wave theory was nonsense, and that light must be a particle and not a wave. But almost immediately, Arago experimentally verified Poisson's prediction.

Simplification of Fresnel diffraction

Recall the Fresnel diffraction result:

$$E(x_0, y_0) \propto \iint \exp \left\{ jk \left[\frac{(-2x_0x_1 - 2y_0y_1)}{2z} + \frac{(x_1^2 + y_1^2)}{2z} \right] \right\} \text{Aperture}(x_1, y_1) dx_1 dy_1$$

Let D be the largest dimension of the aperture: $D^2 = \max(x_1^2 + y_1^2)$.

Our first step, which allowed us to obtain the Fresnel result, was the paraxial approximation:

$$z \gg D \quad \text{or} \quad \frac{D}{z} \ll 1$$

Note that this approximation **does not** contain the wavelength.

A more severe approximation is suggested by noticing that the integral simplifies A LOT if only we could neglect the quadratic terms x_1^2 and y_1^2 .

If $kD^2/2z \ll 1$, then we could do that...

Approximation #2: involving the wavelength

This new approximation, which **does** contain λ , is equivalent to:

$$\frac{D}{z} \ll \frac{2}{kD} = \frac{\lambda}{\pi D}$$

So D/z is not merely required to be less than 1, but is required to be less than $\lambda/\pi D$, which is generally smaller than one for visible light.

Note: if the aperture is a slit of width $D = 2b$, this condition becomes:

$$\frac{2b}{z} \ll \frac{1}{kb} \quad \text{or} \quad \frac{4\pi b^2}{\lambda z} \ll 1$$

Recall our definition of the **Fresnel number** for a slit of width $2b$: $N = \frac{b^2}{\lambda z}$

we see that this new approximation is equivalent to: $4\pi N \ll 1$.

Approximation #2: Fraunhofer diffraction

Apply this approximation: $\frac{kD^2}{2z} \ll 1$ to the Fresnel diffraction result:

$$E(x_0, y_0) \propto \iint \exp \left\{ jk \left[\frac{(-2x_0x_1 - 2y_0y_1)}{2z} + \frac{(x_1^2 + y_1^2)}{2z} \right] \right\} \text{Aperture}(x_1, y_1) dx_1 dy_1$$

In this case, the quadratic terms are tiny, so we can **ignore them**.

As in Fresnel diffraction, we'll typically assume a plane wave incident field, we'll neglect the phase factors, and we'll explicitly write the aperture function in the integral:

$$E(x_0, y_0) \propto \iint \exp \left\{ -\frac{jk}{z} (x_0x_1 + y_0y_1) \right\} \text{Aperture}(x_1, y_1) dx_1 dy_1$$

Fraunhofer Diffraction: when the quadratic terms can be ignored.

Joseph von Fraunhofer

Fraunhofer achieved fame by developing recipes for the world's finest optical glass. He also invented precise methods for measuring dispersion of glass, and discovered more than 500 different absorption lines in sunlight, most due to specific atomic or molecular species at the sun's surface. These are still known as Fraunhofer lines.



Joseph von Fraunhofer
1787-1826

He had almost nothing to do with Fraunhofer diffraction that we're discussing today. He did, however, invent the diffraction grating, which we will discuss next lecture.

He was almost an exact contemporary of Augustin Fresnel. But it is unlikely that they ever met.

The Fraunhofer regime

How far away is far enough? We must have **both** $z \gg D$ and $z \gg \pi D^2/\lambda$.

Example #1:

green light ($\lambda = 0.5 \mu\text{m}$)

If $D = 1$ millimeter, then:

$$z \gg \frac{\pi D^2}{\lambda} = \frac{\pi (1000)^2}{0.5} = 6.3 \text{ meters}$$

If $D = 10$ microns, then:

$$z \gg \frac{\pi D^2}{\lambda} = \frac{\pi (10)^2}{0.5} = 630 \text{ microns}$$

Example #2:

microwaves ($\lambda = 3 \text{ cm}$)

If $D = 10$ centimeters, then:

$$z \gg \frac{\pi D^2}{\lambda} = \frac{\pi (10)^2}{3} = 1 \text{ meter}$$

If $D = 1$ millimeter, then:

$$z \gg \frac{\pi D^2}{\lambda} = \frac{\pi (1)^2}{30} = 0.1 \text{ millimeters}$$

But notice that $z \gg 1 \text{ mm}$ is required for the paraxial approximation...

Fraunhofer diffraction is a Fourier transform

$$E(x_0, y_0) \propto \iint \exp\left\{-\frac{jk}{z}(x_0x_1 + y_0y_1)\right\} \text{Aperture}(x_1, y_1) dx_1 dy_1$$

This is just a Fourier Transform! (actually, two of them, in two variables)


Interestingly, it's a Fourier Transform from position, x_1 , to another position variable, x_0 (in another plane, i.e., a different z position).

Usually, the Fourier “conjugate variables” have reciprocal units (e.g., t & ω , or x & k). The conjugate variables here are really x_1 and kx_0/z , which have reciprocal units.

So, the light in the Fraunhofer regime (the “far field”) is simply the Fourier Transform of the apertured field!

Knowing this makes the calculations a lot easier...

Fraunhofer diffraction pattern for a slit



In this case, the problem is a single Fourier transform (in x), rather than two of them (in x and y):

$$E(x_0) \propto \int \exp\left\{-\frac{jk}{z}(x_0 x_1)\right\} \text{Aperture}(x_1) dx_1$$

The aperture function is simple:

$$\text{Aperture}(x_1) = \begin{cases} 1 & -b < x_1 < b \\ 0 & \text{otherwise} \end{cases}$$

But we know that the Fourier transform of a square function (of width $2b$) is a sinc function:

$$FT[\text{Aperture}(x_1)] \propto \frac{\sin(kx_0 b/z)}{kx_0 b/z} = \frac{\sin(2\pi N x_0/b)}{2\pi N x_0/b}$$



in terms of the
Fresnel number
 $N = b^2/\lambda z$

How very satisfying! This is exactly the answer we found last lecture, for the Fresnel diffraction result in the limit of very large z .


Fraunhofer Diffraction from a Square Aperture

A square aperture (edge length = b) just gives the product of two sinc functions in x and in y . Just as if it were two slits, orthogonal to each other.

$$FT \left[\text{Square Aperture}(x_1, y_1) \right] \propto \frac{\sin(kx_0 b/z)}{kx_0 b/z} \cdot \frac{\sin(ky_0 b/z)}{ky_0 b/z}$$



Diffracted irradiance



Diffracted field



Fraunhofer diffraction from a circular aperture

The 2D Fourier transform of a circular aperture, radius = b , is related to a Bessel function of the first kind:

$$FT [Circular\ aperture(x_1, y_1)] \propto \frac{J_1(k\rho b/z)}{k\rho b/z}$$

where $\rho = \sqrt{x_1^2 + y_1^2}$ is the radial coordinate in the $x_1 - y_1$ plane.


Most of the energy falls in the central region, for values $k\rho b/z < 3.83$



Diffracted E-field plotted in 2D

The Airy pattern

A circular aperture yields a diffracted pattern known as an “Airy pattern” or an “Airy disc”.



$$\text{Diffracted Irradiance} \propto \left| \frac{J_1(k\rho b/z)}{k\rho b/z} \right|^2$$

The central spot contains about 84% of the total energy:


$$\frac{\int_0^{3.83} \left| \frac{J_1(r)}{r} \right|^2 dr}{\int_0^{\infty} \left| \frac{J_1(r)}{r} \right|^2 dr} = 0.838$$



Sir George Biddell Airy
1801-1892

The size of the Airy pattern

How big is the central spot, where most of the energy is found?



Define the size of the spot as ρ_{spot} , the radial distance from the center to the first zero:


$$k \rho_{spot} b / z = 3.83$$
$$\rho_{spot} = \frac{3.83z}{kb} = 3.83 \frac{\lambda z}{2\pi b}$$

The diameter of the aperture is $D = 2b$, so we can write:

$$\rho_{spot} = \frac{3.83}{\pi} \frac{\lambda z}{D} = 1.22 \frac{\lambda z}{D}$$

Diffraction from small and large circular apertures


Small aperture



This is a good illustration of the Scale Theorem!


$$\rho_{spot} \propto \frac{1}{D}$$

Large aperture



Fraunhofer Diffraction: an interesting example

Randomly placed identical holes yield a diffraction pattern whose gross features reveal the shape of the holes.



The Fourier Transform of a random array of identical tiny objects

Define a random array of two-dimensional delta-functions:

$$Rand(x, y) = \sum_{i=1}^n \delta(x - x_i, y - y_i)$$

$$F\{Rand(x, y)\} = \sum_{i=1}^n \exp[-j(k_x x_i + k_y y_i)]$$

Sum of rapidly
varying sinusoids
(looks like noise)



If $OneHole(x, y)$ is the shape of an individual tiny hole, then a random array of identically shaped tiny holes is:

$$Holes(x, y) = Rand(x, y) * OneHole(x, y)$$

The Fourier Transform of a random array of identically shaped tiny holes is:

$$F\{Holes(x, y)\} = F\{Rand(x, y)\} F\{OneHole(x, y)\}$$




rapidly
varying



slowly
varying

Diffraction from multiple slits

Infinitely many equally spaced slits (a Shah function!) yields a far-field pattern which is the Fourier transform; that is, the Shah function.



Laser speckle is a diffraction pattern.



When a laser illuminates a rough surface, it yields a “speckle” pattern. It is a diffraction pattern from the very complex surface.

Don't try to do this Fourier Transform at home.